

Ashland County, Ohio

Cumulative cases of AIDS diagnosed through 2003, by selected characteristics and transmission category

		Cumulative AIDS Diagnoses			
		No.	%		
		No.	%	No.	%
Gender				Transmission category	
Male	10	91%	Male adult or adolescent		
Female	1	9%	Male-to-male sexual contact	6	60%
Unknown	-	-	Injection drug use (IDU)	-	-
			Male-to-male sexual contact and IDU	-	-
Age			Heterosexual contact	-	-
<13	-	-	Blood product	1	10%
13-19	1	9%	Other/unknown	3	30%
20-24	-	-	Subtotal	10	100%
25-29	2	18%	Female adult or adolescent		
30-39	5	45%	Injection drug use	-	-
40-49	2	18%	Heterosexual contact	-	-
50-64	1	9%	Blood product	-	-
65+	-	-	Other/unknown	1	100%
			Subtotal	1	100%
Race/Ethnicity			Child (<13 yrs at diagnosis)		
White, not Hispanic	10	91%	Perinatal	-	-
Black, not Hispanic	-	-	Other	-	-
Hispanic	1	9%	Subtotal	-	-
Asian/Pacific Islander	-	-			
American Indian/Alaska Native	-	-			
Unknown	-	-			
Race/Gender					
White, not Hispanic Male	9	82%			
White, not Hispanic Female	1	9%			
Black, not Hispanic Male	-	-			
Black, not Hispanic Female	-	-			
Other Male	1	9%			
Other Female	-	-			
Unknown	-	-			
Total	11				

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.