

## Auglaize County, Ohio

### Reported persons living with HIV/AIDS as of Dec. 31, 2003, by selected characteristics and transmission category

Reported Persons Living with HIV/AIDS					
	No.	%		No.	%
<b>Current disease status</b>			<b>Race/Gender</b>		
HIV (not AIDS)	8	47%	White, not Hispanic Male	11	65%
AIDS	9	53%	White, not Hispanic Female	4	24%
			Black, not Hispanic Male	-	-
<b>Gender</b>			Black, not Hispanic Female	-	-
Male	12	71%	Other Male	1	6%
Female	5	29%	Other Female	1	6%
Unknown	-	-	Unknown	-	-
<b>Age</b>			<b>Transmission category</b>		
<13	-	-	<b>Male adult or adolescent</b>		
13-19	-	-	Male-to-male sexual contact	6	50%
20-24	-	-	Injection drug use (IDU)	-	-
25-29	-	-	Male-to-male sexual contact and IDU	1	8%
30-39	9	53%			-
40-49	4	24%	Heterosexual contact	-	-
50-64	3	18%	Blood product	1	8%
65+	1	6%	Other/unknown	4	33%
			Subtotal	12	100%
<b>Race/Ethnicity</b>			<b>Female adult or adolescent</b>		
White, not Hispanic	15	88%	Injection drug use	2	40%
Black, not Hispanic	-	-	Heterosexual contact	1	20%
Hispanic	2	12%	Blood product	-	-
Asian/Pacific Islander	-	-	Other/unknown	2	40%
American Indian/Alaska Native	-	-	Subtotal	5	100%
Unknown	-	-	<b>Child (&lt;13 yrs at diagnosis)</b>		
			Perinatal	-	-
			Other	-	-
			Subtotal	-	-
<b>Total</b>	<b>17</b>				

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Auglaize County, Ohio

### Cumulative cases of AIDS diagnosed through 2003, by selected characteristics and transmission category

		Cumulative AIDS Diagnoses			
		No.	%		
		No.	%	No.	%
<b>Gender</b>				<b>Transmission category</b>	
Male	11	73%	<b>Male adult or adolescent</b>		
Female	4	27%	Male-to-male sexual contact	4	36%
Unknown	-	-	Injection drug use (IDU)	1	9%
			Male-to-male sexual contact and IDU	1	9%
<b>Age</b>			Heterosexual contact	-	-
<13	-	-	Blood product	1	9%
13-19	-	-	Other/unknown	4	36%
20-24	1	7%	Subtotal	11	100%
25-29	3	20%	<b>Female adult or adolescent</b>		
30-39	6	40%	Injection drug use	1	25%
40-49	2	13%	Heterosexual contact	2	50%
50-64	2	13%	Blood product	-	-
65+	1	7%	Other/unknown	1	25%
			Subtotal	4	100%
<b>Race/Ethnicity</b>			<b>Child (&lt;13 yrs at diagnosis)</b>		
White, not Hispanic	13	87%	Perinatal	-	-
Black, not Hispanic	-	-	Other	-	-
Hispanic	2	13%	Subtotal	-	-
Asian/Pacific Islander	-	-			
American Indian/Alaska Native	-	-			
Unknown	-	-			
<b>Race/Gender</b>					
White, not Hispanic Male	10	67%			
White, not Hispanic Female	3	20%			
Black, not Hispanic Male	-	-			
Black, not Hispanic Female	-	-			
Other Male	1	7%			
Other Female	1	7%			
Unknown	-	-			
<b>Total</b>	<b>15</b>				

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.