

Belmont County, Ohio

Reported persons living with HIV/AIDS as of Dec. 31, 2003, by selected characteristics and transmission category

Reported Persons Living with HIV/AIDS						
			No.	%		
			No.	%	No.	%
Current disease status					Race/Gender	
HIV (not AIDS)	9	39%	White, not Hispanic Male	18	78%	
AIDS	14	61%	White, not Hispanic Female	1	4%	
			Black, not Hispanic Male	2	9%	
Gender			Black, not Hispanic Female	1	4%	
Male	21	91%	Other Male	1	4%	
Female	2	9%	Other Female	-	-	
Unknown	-	-	Unknown	-	-	
Age			Transmission category			
<13	-	-	Male adult or adolescent			
13-19	1	4%	Male-to-male sexual contact	13	62%	
20-24	1	4%	Injection drug use (IDU)	2	10%	
25-29	-	-	Male-to-male sexual contact and IDU	-	-	
30-39	9	39%	Heterosexual contact	-	-	
40-49	9	39%	Blood product	1	5%	
50-64	2	9%	Other/unknown	5	24%	
65+	1	4%	Subtotal	21	100%	
Race/Ethnicity			Female adult or adolescent			
White, not Hispanic	19	83%	Injection drug use	-	-	
Black, not Hispanic	3	13%	Heterosexual contact	1	50%	
Hispanic	1	4%	Blood product	-	-	
Asian/Pacific Islander	-	-	Other/unknown	1	50%	
American Indian/Alaska Native	-	-	Subtotal	2	100%	
Unknown	-	-	Child (<13 yrs at diagnosis)			
			Perinatal	-	-	
			Other	-	-	
			Subtotal	-	-	
Total	23					

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Belmont County, Ohio

Cumulative cases of AIDS diagnosed through 2003, by selected characteristics and transmission category

		Cumulative AIDS Diagnoses				
		No.	%		No.	%
Gender				Transmission category		
Male	28	97%	Male adult or adolescent			
Female	1	3%	Male-to-male sexual contact	19	68%	
Unknown	-	-	Injection drug use (IDU)	2	7%	
			Male-to-male sexual contact and IDU	3	11%	
Age			Heterosexual contact	-	-	
<13	-	-	Blood product	1	4%	
13-19	1	3%	Other/unknown	3	11%	
20-24	1	3%	Subtotal	28	100%	
25-29	1	3%	Female adult or adolescent			
30-39	16	55%	Injection drug use	-	-	
40-49	8	28%	Heterosexual contact	-	-	
50-64	2	7%	Blood product	-	-	
65+	-	-	Other/unknown	1	100%	
			Subtotal	1	100%	
Race/Ethnicity			Child (<13 yrs at diagnosis)			
White, not Hispanic	26	90%	Perinatal	-	-	
Black, not Hispanic	2	7%	Other	-	-	
Hispanic	1	3%	Subtotal	-	-	
Asian/Pacific Islander	-	-				
American Indian/Alaska Native	-	-				
Unknown	-	-				
Race/Gender						
White, not Hispanic Male	26	90%				
White, not Hispanic Female	-	-				
Black, not Hispanic Male	1	3%				
Black, not Hispanic Female	1	3%				
Other Male	1	3%				
Other Female	-	-				
Unknown	-	-				
Total	29					

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.