

Brown County, Ohio

Reported persons living with HIV/AIDS as of Dec. 31, 2003, by selected characteristics and transmission category

Reported Persons Living with HIV/AIDS					
	No.	%		No.	%
Current disease status			Race/Gender		
HIV (not AIDS)	6	38%	White, not Hispanic Male	14	88%
AIDS	10	63%	White, not Hispanic Female	2	13%
			Black, not Hispanic Male	-	-
Gender			Black, not Hispanic Female	-	-
Male	14	88%	Other Male	-	-
Female	2	13%	Other Female	-	-
Unknown	-	-	Unknown	-	-
Age			Transmission category		
<13	-	-	Male adult or adolescent		
13-19	-	-	Male-to-male sexual contact	7	50%
20-24	-	-	Injection drug use (IDU)	-	-
25-29	-	-	Male-to-male sexual contact and IDU	-	-
30-39	6	38%	Heterosexual contact	2	14%
40-49	6	38%	Blood product	-	-
50-64	4	25%	Other/unknown	5	36%
65+	-	-	Subtotal	14	100%
Race/Ethnicity			Female adult or adolescent		
White, not Hispanic	16	100%	Injection drug use	-	-
Black, not Hispanic	-	-	Heterosexual contact	2	100%
Hispanic	-	-	Blood product	-	-
Asian/Pacific Islander	-	-	Other/unknown	-	-
American Indian/Alaska Native	-	-	Subtotal	2	100%
Unknown	-	-	Child (<13 yrs at diagnosis)		
			Perinatal	-	-
			Other	-	-
			Subtotal	-	-
Total	16				

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Brown County, Ohio

Cumulative cases of AIDS diagnosed through 2003, by selected characteristics and transmission category

		Cumulative AIDS Diagnoses			
		No.	%		
				No.	%
Gender				Transmission category	
Male	12	86%	Male adult or adolescent		
Female	2	14%	Male-to-male sexual contact	6	50%
Unknown	-	-	Injection drug use (IDU)	1	8%
			Male-to-male sexual contact and IDU	-	-
Age			Heterosexual contact	2	17%
<13	-	-	Blood product	1	8%
13-19	-	-	Other/unknown	2	17%
20-24	-	-	Subtotal	12	100%
25-29	2	14%	Female adult or adolescent		
30-39	7	50%	Injection drug use	-	-
40-49	3	21%	Heterosexual contact	1	50%
50-64	-	-	Blood product	1	50%
65+	2	14%	Other/unknown	-	-
			Subtotal	2	100%
Race/Ethnicity			Child (<13 yrs at diagnosis)		
White, not Hispanic	14	100%	Perinatal	-	-
Black, not Hispanic	-	-	Other	-	-
Hispanic	-	-	Subtotal	-	-
Asian/Pacific Islander	-	-			
American Indian/Alaska Native	-	-			
Unknown	-	-			
Race/Gender					
White, not Hispanic Male	12	86%			
White, not Hispanic Female	2	14%			
Black, not Hispanic Male	-	-			
Black, not Hispanic Female	-	-			
Other Male	-	-			
Other Female	-	-			
Unknown	-	-			
Total	14				

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.