

Butler County, Ohio

Reported persons living with HIV/AIDS as of Dec. 31, 2003, by current disease status and selected characteristics

	Current Disease Status						
	HIV/AIDS			HIV (not AIDS)		AIDS	
	Rate ^a	No.	%	No.	%	No.	%
Gender							
Male	102.2	166	83%	81	80%	85	87%
Female	19.4	33	17%	20	20%	13	13%
Unknown	*	-	-	-	-	-	-
Age (as of 12/2003)							
<13	*	1	1%	1	1%	-	-
13-19	*	2	1%	-	-	2	2%
20-24	*	4	2%	2	2%	2	2%
25-29	75.3	16	8%	11	11%	5	5%
30-39	129.2	65	33%	38	38%	27	28%
40-49	135.4	70	35%	30	30%	40	41%
50-64	72.9	35	18%	15	15%	20	20%
65+	16.9	6	3%	4	4%	2	2%
Race/Ethnicity							
White, not Hispanic	48.5	147	74%	76	75%	71	72%
Black, not Hispanic	235.0	43	22%	20	20%	23	23%
Hispanic	104.8	5	3%	2	2%	3	3%
Asian/Pacific Islander	*	-	-	-	-	-	-
American Indian/Alaska Native	*	-	-	-	-	-	-
Unknown	*	4	2%	3	3%	1	1%
Race/Gender							
White, not Hispanic Male	86.6	128	64%	66	65%	62	63%
White, not Hispanic Female	12.2	19	10%	10	10%	9	9%
Black, not Hispanic Male	342.5	30	15%	11	11%	19	19%
Black, not Hispanic Female	136.3	13	7%	9	9%	4	4%
Other Male	*	4	2%	1	1%	3	3%
Other Female	*	1	1%	1	1%	-	-
Unknown	*	4	2%	3	3%	1	1%
Total	59.8	199		101		98	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with HIV/AIDS per 100,000 population calculated using 2000 census data.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Butler County, Ohio

Reported persons living with HIV/AIDS as of Dec. 31, 2003, by current disease status and transmission category

Transmission category	HIV/AIDS		Current Disease Status			
			HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	94	57%	41	51%	53	63%
Injection drug use (IDU)	12	7%	6	8%	6	7%
Male-to-male sexual contact and IDU	4	2%	2	3%	2	2%
Heterosexual contact	4	2%	1	1%	3	4%
Blood product	3	2%	2	3%	1	
Other/unknown	47	29%	28	35%	19	23%
Subtotal	164	100%	80	100%	84	100%
Female adult or adolescent						
Injection drug use	5	16%	3	15%	2	17%
Heterosexual contact	9	28%	5	25%	4	33%
Blood product	1	3%	-	-	1	8%
Other/unknown	17	53%	12	60%	5	42%
Subtotal	32	100%	20	100%	12	100%
Child (<13 yrs at diagnosis)						
Perinatal	2	67%	1	100%	1	50%
Other	1	33%	-	-	1	50%
Subtotal	3	100%	1	100%	2	100%
Total	199		101		98	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Butler County, Ohio

Reported cases of HIV infection, by year of diagnosis and selected characteristics

	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
Gender				
Male	41	87%	25	71%
Female	6	13%	10	29%
Unknown	-	-	-	-
Age				
<13	1	2%	-	-
13-19	1	2%	1	3%
20-24	4	9%	6	17%
25-29	8	17%	2	6%
30-39	20	43%	10	29%
40-49	6	13%	12	34%
50-64	6	13%	3	9%
65+	1	2%	1	3%
Race/Ethnicity				
White, not Hispanic	35	74%	22	63%
Black, not Hispanic	9	19%	9	26%
Hispanic	1	2%	3	9%
Asian/Pacific Islander	-	-	-	-
American Indian/Alaska Native	-	-	-	-
Unknown	2	4%	1	3%
Race/Gender				
White, not Hispanic Male	31	66%	16	46%
White, not Hispanic Female	4	9%	6	17%
Black, not Hispanic Male	7	15%	6	17%
Black, not Hispanic Female	2	4%	3	9%
Other Male	1	2%	2	6%
Other Female	-	-	1	3%
Unknown	2	4%	1	3%
Total	47		35	

Note: HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Butler County, Ohio

Reported cases of HIV infection, by year of diagnosis and transmission category

Transmission category	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
Male adult or adolescent				
Male-to-male sexual contact	16	40%	9	36%
Injection drug use (IDU)	2	5%	2	8%
Male-to-male sexual contact and IDU	-	-	1	4%
Heterosexual contact	1	3%	1	4%
Blood product	-	-	-	-
Other/unknown	21	53%	12	48%
Subtotal	40	100%	25	100%
Female adult or adolescent				
Injection drug use	-	-	-	-
Heterosexual contact	3	50%	1	10%
Blood product	-	-	-	-
Other/unknown	3	50%	9	90%
Subtotal	6	100%	10	100%
Child (<13 yrs at diagnosis)				
Perinatal	1	100%	-	-
Other	-	-	-	-
Subtotal	1	-	-	-
Total	47		35	

Note: HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Butler County, Ohio

Reported cases of AIDS, by year of diagnosis and selected characteristics

	Year of AIDS Diagnosis				Cumulative	
	1998 - 2000		2001 - 2003			
	No.	%	No.	%	No.	%
Gender						
Male	24	100%	17	77%	160	91%
Female	-	-	5	23%	16	9%
Unknown	-	-	-	-	-	-
Age						
<13	-	-	-	-	1	1%
13-19	1	4%	1	5%	3	2%
20-24	1	4%	2	9%	9	5%
25-29	3	13%	2	9%	21	12%
30-39	12	50%	5	23%	80	45%
40-49	4	17%	10	45%	37	21%
50-64	3	13%	2	9%	23	13%
65+	-	-	-	-	2	1%
Race/Ethnicity						
White, not Hispanic	18	75%	13	59%	140	80%
Black, not Hispanic	4	17%	8	36%	31	18%
Hispanic	1	4%	1	5%	3	2%
Asian/Pacific Islander	-	-	-	-	1	1%
American Indian/Alaska Native	-	-	-	-	-	-
Unknown	1	4%	-	-	1	1%
Race/Gender						
White, not Hispanic Male	18	75%	9	41%	132	75%
White, not Hispanic Female	-	-	4	18%	8	5%
Black, not Hispanic Male	4	17%	7	32%	23	13%
Black, not Hispanic Female	-	-	1	5%	8	5%
Other Male	1	4%	1	5%	4	2%
Other Female	-	-	-	-	-	-
Unknown	1	4%	-	-	1	1%
Total	24		22		176	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Butler County, Ohio

Reported cases of AIDS, by year of diagnosis and transmission category

Transmission category	Year of AIDS Diagnosis				Cumulative	
	1998 - 2000		2001 - 2003			
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	15	63%	6	35%	113	71%
Injection drug use (IDU)	-	-	2	12%	7	4%
Male-to-male sexual contact and IDU	-	-	-	-	7	4%
Heterosexual contact	-	-	1	6%	4	3%
Blood product	-	-	-	-	5	3%
Other/unknown	9	38%	8	47%	23	14%
Subtotal	24	100%	17	100%	159	100%
Female adult or adolescent						
Injection drug use	-	-	-	-	4	27%
Heterosexual contact	-	-	-	-	3	20%
Blood product	-	-	-	-	-	-
Other/unknown	-	-	5	100%	8	53%
Subtotal	-	-	5	100%	15	100%
Child (<13 yrs at diagnosis)						
Perinatal	-	-	-	-	1	50%
Other	-	-	-	-	1	50%
Subtotal	-	-	-	-	2	100%
Total	24		22		176	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.