

Champaign County, Ohio

Reported persons living with HIV/AIDS as of Dec. 31, 2003, by selected characteristics and transmission category

Reported Persons Living with HIV/AIDS					
	No.	%		No.	%
Current disease status			Race/Gender		
HIV (not AIDS)	9	43%	White, not Hispanic Male	12	57%
AIDS	12	57%	White, not Hispanic Female	7	33%
			Black, not Hispanic Male	1	5%
Gender			Black, not Hispanic Female	1	5%
Male	13	62%	Other Male	-	-
Female	8	38%	Other Female	-	-
Unknown	-	-	Unknown	-	-
Age			Transmission category		
<13	1	5%	Male adult or adolescent		
13-19	-	-	Male-to-male sexual contact	6	46%
20-24	-	-	Injection drug use (IDU)	-	-
25-29	-	-	Male-to-male sexual contact and IDU	-	-
30-39	6	29%	Heterosexual contact	-	-
40-49	12	57%	Blood product	1	8%
50-64	2	10%	Other/unknown	6	46%
65+	-	-	Subtotal	13	100%
Race/Ethnicity			Female adult or adolescent		
White, not Hispanic	19	90%	Injection drug use	-	-
Black, not Hispanic	2	10%	Heterosexual contact	2	29%
Hispanic	-	-	Blood product	-	-
Asian/Pacific Islander	-	-	Other/unknown	5	71%
American Indian/Alaska Native	-	-	Subtotal	7	100%
Unknown	-	-	Child (<13 yrs at diagnosis)		
			Perinatal	1	100%
			Other	-	-
			Subtotal	1	100%
Total	21				

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Champaign County, Ohio

Cumulative cases of AIDS diagnosed through 2003, by selected characteristics and transmission category

	Cumulative AIDS Diagnoses				
	No.	%		No.	%
Gender			Transmission category		
Male	16	76%	Male adult or adolescent		
Female	5	24%	Male-to-male sexual contact	11	69%
Unknown	-	-	Injection drug use (IDU)	-	-
			Male-to-male sexual contact and IDU	-	-
Age			Heterosexual contact	-	-
<13	2	10%	Blood product	1	6%
13-19	-	-	Other/unknown	4	25%
20-24	-	-	Subtotal	16	100%
25-29	6	29%	Female adult or adolescent		
30-39	6	29%	Injection drug use	-	-
40-49	6	29%	Heterosexual contact	2	67%
50-64	1	5%	Blood product	-	-
65+	-	-	Other/unknown	1	33%
			Subtotal	3	100%
Race/Ethnicity			Child (<13 yrs at diagnosis)		
White, not Hispanic	19	90%	Perinatal	2	100%
Black, not Hispanic	2	10%	Other	-	-
Hispanic	-	-	Subtotal	2	100%
Asian/Pacific Islander	-	-			
American Indian/Alaska Native	-	-			
Unknown	-	-			
Race/Gender					
White, not Hispanic Male	15	71%			
White, not Hispanic Female	4	19%			
Black, not Hispanic Male	1	5%			
Black, not Hispanic Female	1	5%			
Other Male	-	-			
Other Female	-	-			
Unknown	-	-			
Total	21				

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.