

## Clinton County, Ohio

### Reported persons living with HIV/AIDS as of Dec. 31, 2003, by selected characteristics and transmission category

Reported Persons Living with HIV/AIDS						
			No.	%		
			No.	%	No.	%
<b>Current disease status</b>					<b>Race/Gender</b>	
HIV (not AIDS)	6	50%	White, not Hispanic Male	9	75%	
AIDS	6	50%	White, not Hispanic Female	2	17%	
			Black, not Hispanic Male	-	-	
<b>Gender</b>						
Male	10	83%	Black, not Hispanic Female	-	-	
Female	2	17%	Other Male	-	-	
Unknown	-	-	Other Female	-	-	
			Unknown	1	8%	
<b>Age</b>			<b>Transmission category</b>			
<13	-	-	<b>Male adult or adolescent</b>			
13-19	-	-	Male-to-male sexual contact	5	50%	
20-24	-	-	Injection drug use (IDU)	-	-	
25-29	1	8%	Male-to-male sexual contact and IDU	-	-	
30-39	2	17%	Heterosexual contact	-	-	
40-49	6	50%	Blood product	-	-	
50-64	1	8%	Other/unknown	5	50%	
65+	2	17%	Subtotal	10	100%	
<b>Race/Ethnicity</b>			<b>Female adult or adolescent</b>			
White, not Hispanic	11	92%	Injection drug use	-	-	
Black, not Hispanic	-	-	Heterosexual contact	1	50%	
Hispanic	-	-	Blood product	-	-	
Asian/Pacific Islander	-	-	Other/unknown	1	50%	
American Indian/Alaska Native	-	-	Subtotal	2	100%	
Unknown	1	8%	<b>Child (&lt;13 yrs at diagnosis)</b>			
			Perinatal	-	-	
			Other	-	-	
			Subtotal	-	-	
<b>Total</b>	<b>12</b>					

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Clinton County, Ohio

### Cumulative cases of AIDS diagnosed through 2003, by selected characteristics and transmission category

Cumulative AIDS Diagnoses					
	No.	%		No.	%
<b>Gender</b>			<b>Transmission category</b>		
Male	17	94%	<b>Male adult or adolescent</b>		
Female	1	6%	Male-to-male sexual contact	11	65%
Unknown	-	-	Injection drug use (IDU)	-	-
			Male-to-male sexual contact and IDU	-	-
<b>Age</b>			Heterosexual contact	1	6%
<13	-	-	Blood product	1	6%
13-19	-	-	Other/unknown	4	24%
20-24	1	6%	Subtotal	17	100%
25-29	1	6%	<b>Female adult or adolescent</b>		
30-39	3	17%	Injection drug use	-	-
40-49	6	33%	Heterosexual contact	-	-
50-64	6	33%	Blood product	1	100%
65+	1	6%	Other/unknown	-	-
			Subtotal	1	100%
<b>Race/Ethnicity</b>			<b>Child (&lt;13 yrs at diagnosis)</b>		
White, not Hispanic	17	94%	Perinatal	-	-
Black, not Hispanic	1	6%	Other	-	-
Hispanic	-	-	Subtotal	-	-
Asian/Pacific Islander	-	-			
American Indian/Alaska Native	-	-			
Unknown	-	-			
<b>Race/Gender</b>					
White, not Hispanic Male	16	89%			
White, not Hispanic Female	1	6%			
Black, not Hispanic Male	1	6%			
Black, not Hispanic Female	-	-			
Other Male	-	-			
Other Female	-	-			
Unknown	-	-			
<b>Total</b>	<b>18</b>				

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.