

## Cuyahoga County, Ohio

### Reported persons living with HIV/AIDS as of Dec. 31, 2003, by current disease status and selected characteristics

	HIV/AIDS			Current Disease Status			
	Rate <sup>a</sup>	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
<b>Gender</b>							
Male	379.8	2,501	77%	1,094	74%	1,407	80%
Female	100.3	738	23%	385	26%	353	20%
Unknown	*	1	<1%	1	<1%	-	-
<b>Age (as of 12/2003)</b>							
<13	10.3	26	1%	14	1%	12	1%
13-19	20.2	26	1%	16	1%	10	1%
20-24	83.9	65	2%	45	3%	20	1%
25-29	151.3	138	4%	90	6%	48	3%
30-39	486.3	995	31%	515	35%	480	27%
40-49	631.9	1,349	42%	546	37%	803	46%
50-64	279.7	583	18%	232	16%	351	20%
65+	26.7	58	2%	22	1%	36	2%
<b>Race/Ethnicity</b>							
White, not Hispanic	116.2	1,081	33%	460	31%	621	35%
Black, not Hispanic	455.1	1,758	54%	801	54%	957	54%
Hispanic	622.4	293	9%	122	8%	171	10%
Asian/Pacific Islander	36.5	10	<1%	5	<1%	5	<1%
American Indian/Alaska Native	299.2	9	<1%	4	<1%	5	<1%
Unknown	*	89	3%	88	6%	1	<1%
<b>Race/Gender</b>							
White, not Hispanic Male	213.0	948	29%	389	26%	559	32%
White, not Hispanic Female	27.4	133	4%	71	5%	62	4%
Black, not Hispanic Male	730.0	1,277	39%	559	38%	718	41%
Black, not Hispanic Female	227.1	480	15%	241	16%	239	14%
Other Male	541.0	208	6%	79	5%	129	7%
Other Female	266.6	104	3%	52	4%	52	3%
Unknown	*	90	3%	89	6%	1	<1%
<b>Total</b>	<b>232.4</b>	<b>3,240</b>		<b>1,480</b>		<b>1,760</b>	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (\*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> The rate is the number of persons living with HIV/AIDS per 100,000 population calculated using 2000 census data.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Cuyahoga County, Ohio

### Reported persons living with HIV/AIDS as of Dec. 31, 2003, by current disease status and transmission category

Transmission category	HIV/AIDS		Current Disease Status			
			HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	1,388	56%	515	48%	873	62%
Injection drug use (IDU)	249	10%	80	7%	169	12%
Male-to-male sexual contact and IDU	148	6%	50	5%	98	7%
Heterosexual contact	101	4%	39	4%	62	4%
Blood product	13	1%	5	<1%	8	1%
Other/unknown	581	23%	393	36%	188	13%
Subtotal	2,480	100%	1,082	100%	1,398	100%
<b>Female adult or adolescent</b>						
Injection drug use	143	20%	49	13%	94	28%
Heterosexual contact	268	38%	118	32%	150	44%
Blood product	5	1%	2	1%	3	1%
Other/unknown	295	41%	203	55%	92	27%
Subtotal	711	100%	372	100%	339	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	36	75%	18	72%	18	78%
Other	12	25%	7	28%	5	22%
Subtotal	48	100%	25	100%	23	100%
<b>Total</b>	<b>3,240</b>		<b>1,480</b>		<b>1,760</b>	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (\*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Cuyahoga County, Ohio

### Reported cases of HIV infection, by year of diagnosis and selected characteristics

	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
<b>Gender</b>				
Male	473	76%	378	73%
Female	153	24%	139	27%
Unknown	-	-	-	-
<b>Age</b>				
<13	11	2%	6	1%
13-19	18	3%	7	1%
20-24	48	8%	48	9%
25-29	70	11%	67	13%
30-39	259	41%	191	37%
40-49	160	26%	145	28%
50-64	54	9%	52	10%
65+	6	1%	1	<1%
<b>Race/Ethnicity</b>				
White, not Hispanic	199	32%	153	30%
Black, not Hispanic	375	60%	300	58%
Hispanic	41	7%	41	8%
Asian/Pacific Islander	2	<1%	2	<1%
American Indian/Alaska Native	2	<1%	2	<1%
Unknown	7	1%	19	4%
<b>Race/Gender</b>				
White, not Hispanic Male	168	27%	137	26%
White, not Hispanic Female	31	5%	16	3%
Black, not Hispanic Male	263	42%	197	38%
Black, not Hispanic Female	112	18%	103	20%
Other Male	36	6%	29	6%
Other Female	9	1%	16	3%
Unknown	7	1%	19	4%
<b>Total</b>	<b>626</b>		<b>517</b>	

*Note:* HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Cuyahoga County, Ohio

### Reported cases of HIV infection, by year of diagnosis and transmission category

Transmission category	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
<b>Male adult or adolescent</b>				
Male-to-male sexual contact	248	53%	165	44%
Injection drug use (IDU)	41	9%	18	5%
Male-to-male sexual contact and IDU	14	3%	6	2%
Heterosexual contact	20	4%	13	3%
Blood product	-	-	-	-
Other/unknown	143	31%	174	46%
Subtotal	466	100%	376	100%
<b>Female adult or adolescent</b>				
Injection drug use	13	9%	9	7%
Heterosexual contact	51	34%	26	19%
Blood product	2	1%	1	1%
Other/unknown	83	56%	99	73%
Subtotal	149	100%	135	100%
<b>Child (&lt;13 yrs at diagnosis)</b>				
Perinatal	8	73%	5	83%
Other	3	27%	1	17%
Subtotal	11	-	6	-
<b>Total</b>	<b>626</b>		<b>517</b>	

*Note:* HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Cuyahoga County, Ohio

### Reported cases of AIDS, by year of diagnosis and selected characteristics

	Year of AIDS Diagnosis				Cumulative	
	1998 - 2000		2001 - 2003			
	No.	%	No.	%	No.	%
<b>Gender</b>						
Male	335	76%	308	73%	2,770	84%
Female	107	24%	112	27%	546	16%
Unknown	-	-	-	-	-	-
<b>Age</b>						
<13	2	<1%	4	1%	39	1%
13-19	2	<1%	3	1%	20	1%
20-24	15	3%	22	5%	132	4%
25-29	38	9%	34	8%	443	13%
30-39	202	46%	172	41%	1,522	46%
40-49	140	32%	133	32%	819	25%
50-64	38	9%	50	12%	303	9%
65+	5	1%	2	<1%	38	1%
<b>Race/Ethnicity</b>						
White, not Hispanic	128	29%	124	30%	1,281	39%
Black, not Hispanic	289	65%	256	61%	1,792	54%
Hispanic	21	5%	38	9%	231	7%
Asian/Pacific Islander	1	<1%	-	-	6	<1%
American Indian/Alaska Native	3	1%	1	<1%	5	<1%
Unknown	-	-	1	<1%	1	<1%
<b>Race/Gender</b>						
White, not Hispanic Male	112	25%	107	25%	1,193	36%
White, not Hispanic Female	16	4%	17	4%	88	3%
Black, not Hispanic Male	202	46%	176	42%	1,400	42%
Black, not Hispanic Female	87	20%	80	19%	392	12%
Other Male	21	5%	24	6%	176	5%
Other Female	4	1%	15	4%	66	2%
Unknown	-	-	1	<1%	1	<1%
<b>Total</b>	<b>442</b>		<b>420</b>		<b>3,316</b>	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Cuyahoga County, Ohio

### Reported cases of AIDS, by year of diagnosis and transmission category

Transmission category	Year of AIDS Diagnosis				Cumulative	
	1998 - 2000		2001 - 2003			
	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	194	58%	173	57%	1,845	67%
Injection drug use (IDU)	32	10%	25	8%	316	11%
Male-to-male sexual contact and IDU	18	5%	17	6%	221	8%
Heterosexual contact	18	5%	14	5%	98	4%
Blood product	2	1%	-	-	31	1%
Other/unknown	70	21%	77	25%	240	9%
Subtotal	334	100%	306	100%	2,751	100%
<b>Female adult or adolescent</b>						
Injection drug use	21	20%	20	19%	149	29%
Heterosexual contact	45	43%	37	34%	247	48%
Blood product	1	1%	1	1%	7	1%
Other/unknown	38	36%	50	46%	117	23%
Subtotal	105	100%	108	100%	520	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	2	67%	4	67%	37	82%
Other	1	33%	2	33%	8	18%
Subtotal	3	100%	6	100%	45	100%
<b>Total</b>	<b>442</b>		<b>420</b>		<b>3,316</b>	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.