

Erie County, Ohio

Reported persons living with HIV/AIDS as of Dec. 31, 2003, by current disease status and selected characteristics

	Current Disease Status						
	HIV/AIDS			HIV (not AIDS)		AIDS	
	Rate ^a	No.	%	No.	%	No.	%
Gender							
Male	113.5	44	76%	22	76%	22	76%
Female	34.3	14	24%	7	24%	7	24%
Unknown	*	-	-	-	-	-	-
Age (as of 12/2003)							
<13	*	2	3%	1	3%	1	3%
13-19	*	1	2%	1	3%	-	-
20-24	*	-	-	-	-	-	-
25-29	*	1	2%	1	3%	-	-
30-39	205.7	22	38%	11	38%	11	38%
40-49	189.0	24	41%	10	34%	14	48%
50-64	56.9	8	14%	5	17%	3	10%
65+	*	-	-	-	-	-	-
Race/Ethnicity							
White, not Hispanic	51.4	36	62%	17	59%	19	66%
Black, not Hispanic	258.7	19	33%	9	31%	10	34%
Hispanic	*	-	-	-	-	-	-
Asian/Pacific Islander	*	-	-	-	-	-	-
American Indian/Alaska Native	*	-	-	-	-	-	-
Unknown	*	3	5%	3	10%	-	-
Race/Gender							
White, not Hispanic Male	87.9	30	52%	15	52%	15	52%
White, not Hispanic Female	16.7	6	10%	2	7%	4	14%
Black, not Hispanic Male	310.2	11	19%	4	14%	7	24%
Black, not Hispanic Female	210.7	8	14%	5	17%	3	10%
Other Male	*	-	-	-	-	-	-
Other Female	*	-	-	-	-	-	-
Unknown	*	3	5%	3	10%	-	-
Total	72.9	58		29		29	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with HIV/AIDS per 100,000 population calculated using 2000 census data.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Erie County, Ohio

Reported persons living with HIV/AIDS as of Dec. 31, 2003, by current disease status and transmission category

Transmission category	HIV/AIDS		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	20	47%	10	45%	10	48%
Injection drug use (IDU)	1	2%	-	-	1	5%
Male-to-male sexual contact and IDU	4	9%	-	-	4	19%
Heterosexual contact	2	5%	2	9%	-	-
Blood product	-	-	-	-	-	-
Other/unknown	16	37%	10	45%	6	29%
Subtotal	43	100%	22	100%	21	100%
Female adult or adolescent						
Injection drug use	1	8%	-	-	1	14%
Heterosexual contact	6	46%	3	50%	3	43%
Blood product	-	-	-	-	-	-
Other/unknown	6	46%	3	50%	3	43%
Subtotal	13	100%	6	100%	7	100%
Child (<13 yrs at diagnosis)						
Perinatal	2	100%	1	100%	1	100%
Other	-	-	-	-	-	-
Subtotal	2	100%	1	100%	1	100%
Total	58		29		29	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Erie County, Ohio

Reported cases of HIV infection, by year of diagnosis and selected characteristics

	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
Gender				
Male	14	78%	10	83%
Female	4	22%	2	17%
Unknown	-	-	-	-
Age				
<13	-	-	-	-
13-19	-	-	1	8%
20-24	3	17%	-	-
25-29	1	6%	2	17%
30-39	9	50%	4	33%
40-49	4	22%	3	25%
50-64	-	-	2	17%
65+	1	6%	-	-
Race/Ethnicity				
White, not Hispanic	11	61%	10	83%
Black, not Hispanic	7	39%	1	8%
Hispanic	-	-	-	-
Asian/Pacific Islander	-	-	-	-
American Indian/Alaska Native	-	-	-	-
Unknown	-	-	1	8%
Race/Gender				
White, not Hispanic Male	9	50%	8	67%
White, not Hispanic Female	2	11%	2	17%
Black, not Hispanic Male	5	28%	1	8%
Black, not Hispanic Female	2	11%	-	-
Other Male	-	-	-	-
Other Female	-	-	-	-
Unknown	-	-	1	8%
Total	18		12	

Note: HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Erie County, Ohio

Reported cases of HIV infection, by year of diagnosis and transmission category

Transmission category	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
Male adult or adolescent				
Male-to-male sexual contact	5	36%	2	20%
Injection drug use (IDU)	1	7%	-	-
Male-to-male sexual contact and IDU	1	7%	-	-
Heterosexual contact	1	7%	-	-
Blood product	-	-	-	-
Other/unknown	6	43%	8	80%
Subtotal	14	100%	10	100%
Female adult or adolescent				
Injection drug use	1	25%	-	-
Heterosexual contact	1	25%	-	-
Blood product	-	-	-	-
Other/unknown	2	50%	2	100%
Subtotal	4	100%	2	100%
Child (<13 yrs at diagnosis)				
Perinatal	-	-	-	-
Other	-	-	-	-
Subtotal	-	-	-	-
Total	18		12	

Note: HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Erie County, Ohio

Cumulative cases of AIDS diagnosed through 2003, by selected characteristics and transmission category

Cumulative AIDS Diagnoses					
	No.	%		No.	%
Gender			Transmission category		
Male	53	83%	Male adult or adolescent		
Female	11	17%	Male-to-male sexual contact	32	63%
Unknown	-	-	Injection drug use (IDU)	3	6%
			Male-to-male sexual contact and IDU	6	12%
Age			Heterosexual contact	-	-
<13	2	3%	Blood product	-	-
13-19	-	-	Other/unknown	10	20%
20-24	2	3%	Subtotal	51	100%
25-29	15	23%	Female adult or adolescent		
30-39	28	44%	Injection drug use	2	18%
40-49	11	17%	Heterosexual contact	6	55%
50-64	5	8%	Blood product	-	-
65+	1	2%	Other/unknown	3	27%
			Subtotal	11	100%
Race/Ethnicity			Child (<13 yrs at diagnosis)		
White, not Hispanic	50	78%	Perinatal	1	50%
Black, not Hispanic	13	20%	Other	1	50%
Hispanic	1	2%	Subtotal	2	100%
Asian/Pacific Islander	-	-			
American Indian/Alaska Native	-	-			
Unknown	-	-			
Race/Gender					
White, not Hispanic Male	44	69%			
White, not Hispanic Female	6	9%			
Black, not Hispanic Male	8	13%			
Black, not Hispanic Female	5	8%			
Other Male	1	2%			
Other Female	-	-			
Unknown	-	-			
Total	64				

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.