

Fayette County, Ohio

Reported persons living with HIV/AIDS as of Dec. 31, 2003, by selected characteristics and transmission category

Reported Persons Living with HIV/AIDS						
			No.	%		
			No.	%	No.	%
Current disease status					Race/Gender	
HIV (not AIDS)	4	33%	White, not Hispanic Male	10	83%	
AIDS	8	67%	White, not Hispanic Female	1	8%	
			Black, not Hispanic Male	1	8%	
Gender			Black, not Hispanic Female	-	-	
Male	11	92%	Other Male	-	-	
Female	1	8%	Other Female	-	-	
Unknown	-	-	Unknown	-	-	
Age			Transmission category			
<13	-	-	Male adult or adolescent			
13-19	-	-	Male-to-male sexual contact	5	45%	
20-24	1	8%	Injection drug use (IDU)	2	18%	
25-29	-	-	Male-to-male sexual contact and IDU	-	-	
30-39	4	33%	Heterosexual contact	1	9%	
40-49	6	50%	Blood product	-	-	
50-64	1	8%	Other/unknown	3	27%	
65+	-	-	Subtotal	11	100%	
Race/Ethnicity			Female adult or adolescent			
White, not Hispanic	11	92%	Injection drug use	-	-	
Black, not Hispanic	1	8%	Heterosexual contact	-	-	
Hispanic	-	-	Blood product	-	-	
Asian/Pacific Islander	-	-	Other/unknown	1	100%	
American Indian/Alaska Native	-	-	Subtotal	1	100%	
Unknown	-	-	Child (<13 yrs at diagnosis)			
			Perinatal	-	-	
			Other	-	-	
			Subtotal	-	-	
Total	12					

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Fayette County, Ohio

Cumulative cases of AIDS diagnosed through 2003, by selected characteristics and transmission category

		Cumulative AIDS Diagnoses			
		No.	%		
		No.	%	No.	%
Gender				Transmission category	
Male	19	95%	Male adult or adolescent		
Female	1	5%	Male-to-male sexual contact	12	67%
Unknown	-	-	Injection drug use (IDU)	2	11%
			Male-to-male sexual contact and IDU	-	-
Age			Heterosexual contact	2	11%
<13	-	-	Blood product	1	6%
13-19	1	5%	Other/unknown	1	6%
20-24	-	-	Subtotal	18	100%
25-29	2	10%	Female adult or adolescent		
30-39	8	40%	Injection drug use	-	-
40-49	7	35%	Heterosexual contact	-	-
50-64	2	10%	Blood product	1	100%
65+	-	-	Other/unknown	-	-
			Subtotal	1	100%
Race/Ethnicity			Child (<13 yrs at diagnosis)		
White, not Hispanic	19	95%	Perinatal	-	-
Black, not Hispanic	1	5%	Other	1	100%
Hispanic	-	-	Subtotal	1	100%
Asian/Pacific Islander	-	-			
American Indian/Alaska Native	-	-			
Unknown	-	-			
Race/Gender					
White, not Hispanic Male	18	90%			
White, not Hispanic Female	1	5%			
Black, not Hispanic Male	1	5%			
Black, not Hispanic Female	-	-			
Other Male	-	-			
Other Female	-	-			
Unknown	-	-			
Total	20				

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.