

## Fulton County, Ohio

### Reported persons living with HIV/AIDS as of Dec. 31, 2003, by selected characteristics and transmission category

Reported Persons Living with HIV/AIDS					
	No.	%		No.	%
<b>Current disease status</b>			<b>Race/Gender</b>		
HIV (not AIDS)	9	56%	White, not Hispanic Male	11	69%
AIDS	7	44%	White, not Hispanic Female	3	19%
			Black, not Hispanic Male	-	-
<b>Gender</b>			Black, not Hispanic Female	-	-
Male	13	81%	Other Male	2	13%
Female	3	19%	Other Female	-	-
Unknown	-	-	Unknown	-	-
<b>Age</b>			<b>Transmission category</b>		
<13	-	-	<b>Male adult or adolescent</b>		
13-19	-	-	Male-to-male sexual contact	9	69%
20-24	-	-	Injection drug use (IDU)	1	8%
25-29	3	19%	Male-to-male sexual contact and IDU	-	-
30-39	6	38%	Heterosexual contact	-	-
40-49	5	31%	Blood product	-	-
50-64	2	13%	Other/unknown	3	23%
65+	-	-	Subtotal	13	100%
<b>Race/Ethnicity</b>			<b>Female adult or adolescent</b>		
White, not Hispanic	14	88%	Injection drug use	1	33%
Black, not Hispanic	-	-	Heterosexual contact	1	33%
Hispanic	2	13%	Blood product	-	-
Asian/Pacific Islander	-	-	Other/unknown	1	33%
American Indian/Alaska Native	-	-	Subtotal	3	100%
Unknown	-	-	<b>Child (&lt;13 yrs at diagnosis)</b>		
			Perinatal	-	-
			Other	-	-
			Subtotal	-	-
<b>Total</b>	<b>16</b>				

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Fulton County, Ohio

### Cumulative cases of AIDS diagnosed through 2003, by selected characteristics and transmission category

		Cumulative AIDS Diagnoses			
		No.	%		
		No.	%	No.	%
<b>Gender</b>				<b>Transmission category</b>	
Male	10	77%		<b>Male adult or adolescent</b>	
Female	3	23%		Male-to-male sexual contact	7 70%
Unknown	-	-		Injection drug use (IDU)	- -
				Male-to-male sexual contact and IDU	- -
<b>Age</b>				Heterosexual contact	1 10%
<13	-	-		Blood product	- -
13-19	-	-		Other/unknown	2 20%
20-24	1	8%		Subtotal	10 100%
25-29	4	31%		<b>Female adult or adolescent</b>	
30-39	4	31%		Injection drug use	- -
40-49	2	15%		Heterosexual contact	2 67%
50-64	2	15%		Blood product	1 33%
65+	-	-		Other/unknown	- -
				Subtotal	3 100%
<b>Race/Ethnicity</b>				<b>Child (&lt;13 yrs at diagnosis)</b>	
White, not Hispanic	12	92%		Perinatal	- -
Black, not Hispanic	-	-		Other	- -
Hispanic	1	8%		Subtotal	- -
Asian/Pacific Islander	-	-			
American Indian/Alaska Native	-	-			
Unknown	-	-			
<b>Race/Gender</b>					
White, not Hispanic Male	9	69%			
White, not Hispanic Female	3	23%			
Black, not Hispanic Male	-	-			
Black, not Hispanic Female	-	-			
Other Male	1	8%			
Other Female	-	-			
Unknown	-	-			
<b>Total</b>	<b>13</b>				

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.