

Gallia County, Ohio

Reported persons living with HIV/AIDS as of Dec. 31, 2003, by selected characteristics and transmission category

Reported Persons Living with HIV/AIDS					
	No.	%		No.	%
Current disease status			Race/Gender		
HIV (not AIDS)	5	38%	White, not Hispanic Male	7	54%
AIDS	8	62%	White, not Hispanic Female	3	23%
			Black, not Hispanic Male	-	-
Gender			Black, not Hispanic Female	1	8%
Male	9	69%	Other Male	1	8%
Female	4	31%	Other Female	-	-
Unknown	-	-	Unknown	1	8%
Age			Transmission category		
<13	-	-	Male adult or adolescent		
13-19	-	-	Male-to-male sexual contact	4	50%
20-24	-	-	Injection drug use (IDU)	1	13%
25-29	1	8%	Male-to-male sexual contact and IDU	1	13%
30-39	6	46%			-
40-49	4	31%	Heterosexual contact	-	-
50-64	2	15%	Blood product	-	-
65+	-	-	Other/unknown	2	25%
			Subtotal	8	100%
Race/Ethnicity			Female adult or adolescent		
White, not Hispanic	10	77%	Injection drug use	1	25%
Black, not Hispanic	1	8%	Heterosexual contact	2	50%
Hispanic	1	8%	Blood product	-	-
Asian/Pacific Islander	-	-	Other/unknown	1	25%
American Indian/Alaska Native	-	-	Subtotal	4	100%
Unknown	1	8%	Child (<13 yrs at diagnosis)		
			Perinatal	-	-
			Other	1	100%
			Subtotal	1	100%
Total	13				

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Gallia County, Ohio

Cumulative cases of AIDS diagnosed through 2003, by selected characteristics and transmission category

Cumulative AIDS Diagnoses					
	No.	%		No.	%
Gender			Transmission category		
Male	11	92%	Male adult or adolescent		
Female	1	8%	Male-to-male sexual contact	5	50%
Unknown	-	-	Injection drug use (IDU)	3	30%
			Male-to-male sexual contact and IDU	-	-
Age			Heterosexual contact	-	-
<13	-	-	Blood product	-	-
13-19	-	-	Other/unknown	2	20%
20-24	1	8%	Subtotal	10	100%
25-29	1	8%	Female adult or adolescent		
30-39	7	58%	Injection drug use	-	-
40-49	2	17%	Heterosexual contact	1	100%
50-64	1	8%	Blood product	-	-
65+	-	-	Other/unknown	-	-
			Subtotal	1	100%
Race/Ethnicity			Child (<13 yrs at diagnosis)		
White, not Hispanic	7	58%	Perinatal	-	-
Black, not Hispanic	3	25%	Other	1	100%
Hispanic	1	8%	Subtotal	1	100%
Asian/Pacific Islander	1	8%			
American Indian/Alaska Native	-	-			
Unknown	-	-			
Race/Gender					
White, not Hispanic Male	6	50%			
White, not Hispanic Female	1	8%			
Black, not Hispanic Male	3	25%			
Black, not Hispanic Female	-	-			
Other Male	2	17%			
Other Female	-	-			
Unknown	-	-			
Total	12				

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.