

Geauga County, Ohio

Reported persons living with HIV/AIDS as of Dec. 31, 2003, by selected characteristics and transmission category

Reported Persons Living with HIV/AIDS						
			No.	%		
			No.	%	No.	%
Current disease status					Race/Gender	
HIV (not AIDS)	4	18%	White, not Hispanic Male	19	86%	
AIDS	18	82%	White, not Hispanic Female	1	5%	
			Black, not Hispanic Male	1	5%	
Gender			Black, not Hispanic Female	-	-	
Male	21	95%	Other Male	1	5%	
Female	1	5%	Other Female	-	-	
Unknown	-	-	Unknown	-	-	
Age			Transmission category			
<13	-	-	Male adult or adolescent			
13-19	-	-	Male-to-male sexual contact	14	67%	
20-24	-	-	Injection drug use (IDU)	2	10%	
25-29	-	-	Male-to-male sexual contact and IDU	1	5%	
30-39	4	18%	Heterosexual contact	-	-	
40-49	15	68%	Blood product	-	-	
50-64	1	5%	Other/unknown	4	19%	
65+	2	9%	Subtotal	21	100%	
Race/Ethnicity			Female adult or adolescent			
White, not Hispanic	20	91%	Injection drug use	-	-	
Black, not Hispanic	1	5%	Heterosexual contact	-	-	
Hispanic	-	-	Blood product	1	100%	
Asian/Pacific Islander	1	5%	Other/unknown	-	-	
American Indian/Alaska Native	-	-	Subtotal	1	100%	
Unknown	-	-	Child (<13 yrs at diagnosis)			
			Perinatal	-	-	
			Other	-	-	
			Subtotal	-	-	
Total	22					

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Geauga County, Ohio

Cumulative cases of AIDS diagnosed through 2003, by selected characteristics and transmission category

Cumulative AIDS Diagnoses					
	No.	%		No.	%
Gender			Transmission category		
Male	31	94%	Male adult or adolescent		
Female	2	6%	Male-to-male sexual contact	24	77%
Unknown	-	-	Injection drug use (IDU)	1	3%
			Male-to-male sexual contact and IDU	2	6%
Age			Heterosexual contact	-	-
<13	-	-	Blood product	1	3%
13-19	-	-	Other/unknown	3	10%
20-24	2	6%	Subtotal	31	100%
25-29	3	9%	Female adult or adolescent		
30-39	14	42%	Injection drug use	-	-
40-49	10	30%	Heterosexual contact	1	50%
50-64	1	3%	Blood product	1	50%
65+	3	9%	Other/unknown	-	-
			Subtotal	2	100%
Race/Ethnicity			Child (<13 yrs at diagnosis)		
White, not Hispanic	30	91%	Perinatal	-	-
Black, not Hispanic	3	9%	Other	-	-
Hispanic	-	-	Subtotal	-	-
Asian/Pacific Islander	-	-			
American Indian/Alaska Native	-	-			
Unknown	-	-			
Race/Gender					
White, not Hispanic Male	28	85%			
White, not Hispanic Female	2	6%			
Black, not Hispanic Male	3	9%			
Black, not Hispanic Female	-	-			
Other Male	-	-			
Other Female	-	-			
Unknown	-	-			
Total	33				

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.