

Hamilton County, Ohio

Reported persons living with HIV/AIDS as of Dec. 31, 2003, by current disease status and selected characteristics

| | Current Disease Status | | | | | | |
|-------------------------------|------------------------|--------------|-----|----------------|-----|------------|-----|
| | HIV/AIDS | | | HIV (not AIDS) | | AIDS | |
| | Rate ^a | No. | % | No. | % | No. | % |
| Gender | | | | | | | |
| Male | 368.0 | 1,483 | 82% | 784 | 78% | 699 | 87% |
| Female | 74.4 | 329 | 18% | 222 | 22% | 107 | 13% |
| Unknown | * | - | - | - | - | - | - |
| Age (as of 12/2003) | | | | | | | |
| <13 | 7.1 | 11 | 1% | 8 | 1% | 3 | <1% |
| 13-19 | * | 4 | <1% | 3 | <1% | 1 | <1% |
| 20-24 | 94.8 | 54 | 3% | 49 | 5% | 5 | 1% |
| 25-29 | 202.7 | 118 | 7% | 87 | 9% | 31 | 4% |
| 30-39 | 455.3 | 568 | 31% | 337 | 33% | 231 | 29% |
| 40-49 | 576.2 | 742 | 41% | 379 | 38% | 363 | 45% |
| 50-64 | 237.0 | 286 | 16% | 132 | 13% | 154 | 19% |
| 65+ | 25.5 | 29 | 2% | 11 | 1% | 18 | 2% |
| Race/Ethnicity | | | | | | | |
| White, not Hispanic | 131.2 | 810 | 45% | 435 | 43% | 375 | 47% |
| Black, not Hispanic | 455.4 | 918 | 51% | 510 | 51% | 408 | 51% |
| Hispanic | 262.8 | 25 | 1% | 13 | 1% | 12 | 1% |
| Asian/Pacific Islander | 53.8 | 8 | <1% | 3 | <1% | 5 | 1% |
| American Indian/Alaska Native | * | 3 | <1% | 1 | <1% | 2 | <1% |
| Unknown | * | 48 | 3% | 44 | 4% | 4 | <1% |
| Race/Gender | | | | | | | |
| White, not Hispanic Male | 244.1 | 726 | 40% | 373 | 37% | 353 | 44% |
| White, not Hispanic Female | 26.2 | 84 | 5% | 62 | 6% | 22 | 3% |
| Black, not Hispanic Male | 747.0 | 688 | 38% | 361 | 36% | 327 | 41% |
| Black, not Hispanic Female | 210.1 | 230 | 13% | 149 | 15% | 81 | 10% |
| Other Male | 223.9 | 30 | 2% | 14 | 1% | 16 | 2% |
| Other Female | 46.8 | 6 | <1% | 3 | <1% | 3 | <1% |
| Unknown | * | 48 | 3% | 44 | 4% | 4 | <1% |
| Total | 214.4 | 1,812 | | 1,006 | | 806 | |

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with HIV/AIDS per 100,000 population calculated using 2000 census data.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Hamilton County, Ohio

Reported persons living with HIV/AIDS as of Dec. 31, 2003, by current disease status and transmission category

| Transmission category | HIV/AIDS | | Current Disease Status | | | |
|--|--------------|------|------------------------|------|------------|------|
| | | | HIV (not AIDS) | | AIDS | |
| | No. | % | No. | % | No. | % |
| Male adult or adolescent | | | | | | |
| Male-to-male sexual contact | 780 | 53% | 333 | 43% | 447 | 64% |
| Injection drug use (IDU) | 87 | 6% | 31 | 4% | 56 | 8% |
| Male-to-male sexual contact and IDU | 73 | 5% | 25 | 3% | 48 | 7% |
| Heterosexual contact | 36 | 2% | 14 | 2% | 22 | 3% |
| Blood product | 11 | 1% | 5 | 1% | 6 | 1% |
| Other/unknown | 486 | 33% | 371 | 48% | 115 | 17% |
| Subtotal | 1,473 | 100% | 779 | 100% | 694 | 100% |
| Female adult or adolescent | | | | | | |
| Injection drug use | 35 | 11% | 15 | 7% | 20 | 19% |
| Heterosexual contact | 108 | 33% | 64 | 29% | 44 | 42% |
| Blood product | 2 | 1% | 2 | 1% | - | - |
| Other/unknown | 178 | 55% | 136 | 63% | 42 | 40% |
| Subtotal | 323 | 100% | 217 | 100% | 106 | 100% |
| Child (<13 yrs at diagnosis) | | | | | | |
| Perinatal | 11 | 69% | 8 | 80% | 3 | 50% |
| Other | 5 | 31% | 2 | 20% | 3 | 50% |
| Subtotal | 16 | 100% | 10 | 100% | 6 | 100% |
| Total | 1,812 | | 1,006 | | 806 | |

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Hamilton County, Ohio

Reported cases of HIV infection, by year of diagnosis and selected characteristics

| | Year of HIV Diagnosis | | | |
|-------------------------------|-----------------------|-----|-------------|-----|
| | 1998 - 2000 | | 2001 - 2003 | |
| | No. | % | No. | % |
| Gender | | | | |
| Male | 269 | 79% | 275 | 78% |
| Female | 72 | 21% | 78 | 22% |
| Unknown | - | - | - | - |
| Age | | | | |
| <13 | 2 | 1% | - | - |
| 13-19 | 13 | 4% | 12 | 3% |
| 20-24 | 35 | 10% | 39 | 11% |
| 25-29 | 55 | 16% | 49 | 14% |
| 30-39 | 137 | 40% | 118 | 33% |
| 40-49 | 72 | 21% | 99 | 28% |
| 50-64 | 24 | 7% | 31 | 9% |
| 65+ | 3 | 1% | 5 | 1% |
| Race/Ethnicity | | | | |
| White, not Hispanic | 139 | 41% | 133 | 38% |
| Black, not Hispanic | 180 | 53% | 194 | 55% |
| Hispanic | 5 | 1% | 6 | 2% |
| Asian/Pacific Islander | 1 | <1% | - | - |
| American Indian/Alaska Native | - | - | - | - |
| Unknown | 16 | 5% | 20 | 6% |
| Race/Gender | | | | |
| White, not Hispanic Male | 120 | 35% | 114 | 32% |
| White, not Hispanic Female | 19 | 6% | 19 | 5% |
| Black, not Hispanic Male | 132 | 39% | 142 | 40% |
| Black, not Hispanic Female | 48 | 14% | 52 | 15% |
| Other Male | 4 | 1% | 4 | 1% |
| Other Female | 2 | 1% | 2 | 1% |
| Unknown | 16 | 5% | 20 | 6% |
| Total | 341 | | 353 | |

Note: HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Hamilton County, Ohio

Reported cases of HIV infection, by year of diagnosis and transmission category

| Transmission category | Year of HIV Diagnosis | | | |
|--|-----------------------|------|-------------|------|
| | 1998 - 2000 | | 2001 - 2003 | |
| | No. | % | No. | % |
| Male adult or adolescent | | | | |
| Male-to-male sexual contact | 119 | 44% | 98 | 36% |
| Injection drug use (IDU) | 11 | 4% | 12 | 4% |
| Male-to-male sexual contact and IDU | 5 | 2% | 3 | 1% |
| Heterosexual contact | 2 | 1% | 12 | 4% |
| Blood product | 1 | <1% | 2 | 1% |
| Other/unknown | 131 | 49% | 148 | 54% |
| Subtotal | 269 | 100% | 275 | 100% |
| Female adult or adolescent | | | | |
| Injection drug use | 3 | 4% | 3 | 4% |
| Heterosexual contact | 21 | 30% | 17 | 22% |
| Blood product | - | - | - | - |
| Other/unknown | 46 | 66% | 58 | 74% |
| Subtotal | 70 | 100% | 78 | 100% |
| Child (<13 yrs at diagnosis) | | | | |
| Perinatal | - | - | - | - |
| Other | 2 | 100% | - | - |
| Subtotal | 2 | - | - | - |
| Total | 341 | | 353 | |

Note: HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Hamilton County, Ohio

Reported cases of AIDS, by year of diagnosis and selected characteristics

| | Year of AIDS Diagnosis | | | | Cumulative | |
|-------------------------------|------------------------|-----|-------------|-----|--------------|-----|
| | 1998 - 2000 | | 2001 - 2003 | | | |
| | No. | % | No. | % | No. | % |
| Gender | | | | | | |
| Male | 151 | 86% | 135 | 84% | 1,532 | 91% |
| Female | 25 | 14% | 26 | 16% | 160 | 9% |
| Unknown | - | - | - | - | - | - |
| Age | | | | | | |
| <13 | 1 | 1% | - | - | 13 | 1% |
| 13-19 | - | - | 2 | 1% | 8 | <1% |
| 20-24 | 5 | 3% | 7 | 4% | 54 | 3% |
| 25-29 | 29 | 16% | 14 | 9% | 273 | 16% |
| 30-39 | 81 | 46% | 67 | 42% | 796 | 47% |
| 40-49 | 45 | 26% | 50 | 31% | 404 | 24% |
| 50-64 | 13 | 7% | 17 | 11% | 126 | 7% |
| 65+ | 2 | 1% | 4 | 2% | 18 | 1% |
| Race/Ethnicity | | | | | | |
| White, not Hispanic | 72 | 41% | 57 | 35% | 931 | 55% |
| Black, not Hispanic | 101 | 57% | 97 | 60% | 737 | 44% |
| Hispanic | - | - | 3 | 2% | 12 | 1% |
| Asian/Pacific Islander | 2 | 1% | 1 | 1% | 5 | <1% |
| American Indian/Alaska Native | - | - | - | - | 2 | <1% |
| Unknown | 1 | 1% | 3 | 2% | 5 | <1% |
| Race/Gender | | | | | | |
| White, not Hispanic Male | 68 | 39% | 49 | 30% | 886 | 52% |
| White, not Hispanic Female | 4 | 2% | 8 | 5% | 45 | 3% |
| Black, not Hispanic Male | 81 | 46% | 80 | 50% | 626 | 37% |
| Black, not Hispanic Female | 20 | 11% | 17 | 11% | 111 | 7% |
| Other Male | 1 | 1% | 4 | 2% | 16 | 1% |
| Other Female | 1 | 1% | - | - | 3 | <1% |
| Unknown | 1 | 1% | 3 | 2% | 5 | <1% |
| Total | 176 | | 161 | | 1,692 | |

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Hamilton County, Ohio

Reported cases of AIDS, by year of diagnosis and transmission category

| Transmission category | Year of AIDS Diagnosis | | | | Cumulative | |
|--|------------------------|------|-------------|------|--------------|------|
| | 1998 - 2000 | | 2001 - 2003 | | | |
| | No. | % | No. | % | No. | % |
| Male adult or adolescent | | | | | | |
| Male-to-male sexual contact | 92 | 61% | 63 | 47% | 1,105 | 73% |
| Injection drug use (IDU) | 9 | 6% | 11 | 8% | 83 | 5% |
| Male-to-male sexual contact and IDU | 6 | 4% | 6 | 4% | 106 | 7% |
| Heterosexual contact | 1 | 1% | 4 | 3% | 29 | 2% |
| Blood product | 1 | 1% | - | - | 26 | 2% |
| Other/unknown | 42 | 28% | 50 | 37% | 172 | 11% |
| Subtotal | 151 | 100% | 134 | 100% | 1,521 | 100% |
| Female adult or adolescent | | | | | | |
| Injection drug use | 3 | 13% | 2 | 8% | 35 | 22% |
| Heterosexual contact | 7 | 29% | 8 | 31% | 67 | 43% |
| Blood product | - | - | - | - | 2 | 1% |
| Other/unknown | 14 | 58% | 16 | 62% | 52 | 33% |
| Subtotal | 24 | 100% | 26 | 100% | 156 | 100% |
| Child (<13 yrs at diagnosis) | | | | | | |
| Perinatal | - | - | - | - | 10 | 67% |
| Other | 1 | 100% | 1 | 100% | 5 | 33% |
| Subtotal | 1 | 100% | 1 | 100% | 15 | 100% |
| Total | 176 | | 161 | | 1,692 | |

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.