

## Highland County, Ohio

### Reported persons living with HIV/AIDS as of Dec. 31, 2003, by selected characteristics and transmission category

Reported Persons Living with HIV/AIDS					
	No.	%		No.	%
<b>Current disease status</b>			<b>Race/Gender</b>		
HIV (not AIDS)	7	41%	White, not Hispanic Male	14	82%
AIDS	10	59%	White, not Hispanic Female	1	6%
			Black, not Hispanic Male	1	6%
			Black, not Hispanic Female	1	6%
<b>Gender</b>			Other Male	-	-
Male	15	88%	Other Female	-	-
Female	2	12%	Unknown	-	-
Unknown	-	-			
<b>Age</b>			<b>Transmission category</b>		
<13	-	-	<b>Male adult or adolescent</b>		
13-19	-	-	Male-to-male sexual contact	4	29%
20-24	-	-	Injection drug use (IDU)	-	-
25-29	1	6%	Male-to-male sexual contact and IDU	1	7%
30-39	5	29%	Heterosexual contact	-	-
40-49	9	53%	Blood product	-	-
50-64	2	12%	Other/unknown	9	64%
65+	-	-	Subtotal	14	100%
<b>Race/Ethnicity</b>			<b>Female adult or adolescent</b>		
White, not Hispanic	15	88%	Injection drug use	-	-
Black, not Hispanic	2	12%	Heterosexual contact	2	100%
Hispanic	-	-	Blood product	-	-
Asian/Pacific Islander	-	-	Other/unknown	-	-
American Indian/Alaska Native	-	-	Subtotal	2	100%
Unknown	-	-			
			<b>Child (&lt;13 yrs at diagnosis)</b>		
			Perinatal	-	-
			Other	1	100%
			Subtotal	1	100%
<b>Total</b>	<b>17</b>				

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Highland County, Ohio

### Cumulative cases of AIDS diagnosed through 2003, by selected characteristics and transmission category

		Cumulative AIDS Diagnoses			
		No.	%		
		No.	%	No.	%
<b>Gender</b>				<b>Transmission category</b>	
Male	13	87%	<b>Male adult or adolescent</b>		
Female	2	13%	Male-to-male sexual contact	8	62%
Unknown	-	-	Injection drug use (IDU)	-	-
			Male-to-male sexual contact and IDU	-	-
<b>Age</b>			Heterosexual contact	-	-
<13	-	-	Blood product	-	-
13-19	-	-	Other/unknown	5	38%
20-24	1	7%	Subtotal	13	100%
25-29	1	7%	<b>Female adult or adolescent</b>		
30-39	8	53%	Injection drug use	-	-
40-49	4	27%	Heterosexual contact	1	50%
50-64	1	7%	Blood product	-	-
65+	-	-	Other/unknown	1	50%
			Subtotal	2	100%
<b>Race/Ethnicity</b>			<b>Child (&lt;13 yrs at diagnosis)</b>		
White, not Hispanic	15	100%	Perinatal	-	-
Black, not Hispanic	-	-	Other	-	-
Hispanic	-	-	Subtotal	-	-
Asian/Pacific Islander	-	-			
American Indian/Alaska Native	-	-			
Unknown	-	-			
<b>Race/Gender</b>					
White, not Hispanic Male	13	87%			
White, not Hispanic Female	2	13%			
Black, not Hispanic Male	-	-			
Black, not Hispanic Female	-	-			
Other Male	-	-			
Other Female	-	-			
Unknown	-	-			
<b>Total</b>	<b>15</b>				

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.