

Hocking County, Ohio

Reported persons living with HIV/AIDS as of Dec. 31, 2003, by selected characteristics and transmission category

Reported Persons Living with HIV/AIDS						
			No.	%		
			No.	%	No.	%
Current disease status					Race/Gender	
HIV (not AIDS)	9	53%	White, not Hispanic Male	12	71%	
AIDS	8	47%	White, not Hispanic Female	5	29%	
			Black, not Hispanic Male	-	-	
Gender						
Male	12	71%	Black, not Hispanic Female	-	-	
Female	5	29%	Other Male	-	-	
Unknown	-	-	Other Female	-	-	
			Unknown	-	-	
Age					Transmission category	
<13	1	6%	Male adult or adolescent			
13-19	1	6%	Male-to-male sexual contact	3	25%	
20-24	-	-	Injection drug use (IDU)	2	17%	
25-29	-	-	Male-to-male sexual contact and IDU	-	-	
30-39	4	24%	Heterosexual contact	1	8%	
40-49	8	47%	Blood product	-	-	
50-64	3	18%	Other/unknown	6	50%	
65+	-	-	Subtotal	12	100%	
Race/Ethnicity					Female adult or adolescent	
White, not Hispanic	17	100%	Injection drug use	-	-	
Black, not Hispanic	-	-	Heterosexual contact	2	67%	
Hispanic	-	-	Blood product	-	-	
Asian/Pacific Islander	-	-	Other/unknown	1	33%	
American Indian/Alaska Native	-	-	Subtotal	3	100%	
Unknown	-	-	Child (<13 yrs at diagnosis)			
			Perinatal	2	100%	
			Other	-	-	
			Subtotal	2	100%	
Total	17					

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Hocking County, Ohio

Cumulative cases of AIDS diagnosed through 2003, by selected characteristics and transmission category

Cumulative AIDS Diagnoses					
	No.	%		No.	%
Gender			Transmission category		
Male	13	87%	Male adult or adolescent		
Female	2	13%	Male-to-male sexual contact	7	54%
Unknown	-	-	Injection drug use (IDU)	1	8%
			Male-to-male sexual contact and IDU	-	-
Age			Heterosexual contact	1	8%
<13	-	-	Blood product	-	-
13-19	-	-	Other/unknown	4	31%
20-24	-	-	Subtotal	13	100%
25-29	2	13%	Female adult or adolescent		
30-39	4	27%	Injection drug use	-	-
40-49	5	33%	Heterosexual contact	1	50%
50-64	4	27%	Blood product	-	-
65+	-	-	Other/unknown	1	50%
			Subtotal	2	100%
Race/Ethnicity			Child (<13 yrs at diagnosis)		
White, not Hispanic	15	100%	Perinatal	-	-
Black, not Hispanic	-	-	Other	-	-
Hispanic	-	-	Subtotal	-	-
Asian/Pacific Islander	-	-			
American Indian/Alaska Native	-	-			
Unknown	-	-			
Race/Gender					
White, not Hispanic Male	13	87%			
White, not Hispanic Female	2	13%			
Black, not Hispanic Male	-	-			
Black, not Hispanic Female	-	-			
Other Male	-	-			
Other Female	-	-			
Unknown	-	-			
Total	15				

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.