

Jackson County, Ohio

Reported persons living with HIV/AIDS as of Dec. 31, 2003, by selected characteristics and transmission category

Reported Persons Living with HIV/AIDS					
	No.	%		No.	%
Current disease status			Race/Gender		
HIV (not AIDS)	7	50%	White, not Hispanic Male	12	86%
AIDS	7	50%	White, not Hispanic Female	1	7%
			Black, not Hispanic Male	-	-
Gender			Black, not Hispanic Female	1	7%
Male	12	86%	Other Male	-	-
Female	2	14%	Other Female	-	-
Unknown	-	-	Unknown	-	-
Age			Transmission category		
<13	-	-	Male adult or adolescent		
13-19	-	-	Male-to-male sexual contact	6	50%
20-24	1	7%	Injection drug use (IDU)	-	-
25-29	-	-	Male-to-male sexual contact and IDU	2	17%
30-39	7	50%	Heterosexual contact	-	-
40-49	3	21%	Blood product	-	-
50-64	3	21%	Other/unknown	4	33%
65+	-	-	Subtotal	12	100%
Race/Ethnicity			Female adult or adolescent		
White, not Hispanic	13	93%	Injection drug use	-	-
Black, not Hispanic	1	7%	Heterosexual contact	1	50%
Hispanic	-	-	Blood product	-	-
Asian/Pacific Islander	-	-	Other/unknown	1	50%
American Indian/Alaska Native	-	-	Subtotal	2	100%
Unknown	-	-	Child (<13 yrs at diagnosis)		
			Perinatal	-	-
			Other	-	-
			Subtotal	-	-
Total	14				

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Jackson County, Ohio

Cumulative cases of AIDS diagnosed through 2003, by selected characteristics and transmission category

	Cumulative AIDS Diagnoses				
	No.	%		No.	%
Gender			Transmission category		
Male	8	80%	Male adult or adolescent		
Female	2	20%	Male-to-male sexual contact	8	100%
Unknown	-	-	Injection drug use (IDU)	-	-
			Male-to-male sexual contact and IDU	-	-
Age			Heterosexual contact	-	-
<13	-	-	Blood product	-	-
13-19	-	-	Other/unknown	-	-
20-24	1	10%	Subtotal	8	100%
25-29	1	10%	Female adult or adolescent		
30-39	5	50%	Injection drug use	-	-
40-49	3	30%	Heterosexual contact	1	50%
50-64	-	-	Blood product	1	50%
65+	-	-	Other/unknown	-	-
			Subtotal	2	100%
Race/Ethnicity			Child (<13 yrs at diagnosis)		
White, not Hispanic	10	100%	Perinatal	-	-
Black, not Hispanic	-	-	Other	-	-
Hispanic	-	-	Subtotal	-	-
Asian/Pacific Islander	-	-			
American Indian/Alaska Native	-	-			
Unknown	-	-			
Race/Gender					
White, not Hispanic Male	8	80%			
White, not Hispanic Female	2	20%			
Black, not Hispanic Male	-	-			
Black, not Hispanic Female	-	-			
Other Male	-	-			
Other Female	-	-			
Unknown	-	-			
Total	10				

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.