

Knox County, Ohio

Reported persons living with HIV/AIDS as of Dec. 31, 2003, by selected characteristics and transmission category

Reported Persons Living with HIV/AIDS					
	No.	%		No.	%
Current disease status			Race/Gender		
HIV (not AIDS)	12	63%	White, not Hispanic Male	13	68%
AIDS	7	37%	White, not Hispanic Female	1	5%
			Black, not Hispanic Male	3	16%
Gender			Black, not Hispanic Female	1	5%
Male	16	84%	Other Male	-	-
Female	3	16%	Other Female	-	-
Unknown	-	-	Unknown	1	5%
Age			Transmission category		
<13	1	5%	Male adult or adolescent		
13-19	-	-	Male-to-male sexual contact	7	47%
20-24	-	-	Injection drug use (IDU)	1	7%
25-29	2	11%	Male-to-male sexual contact and IDU	1	7%
30-39	6	32%			-
40-49	4	21%	Heterosexual contact	1	7%
50-64	6	32%	Blood product	-	-
65+	-	-	Other/unknown	5	33%
			Subtotal	15	100%
Race/Ethnicity			Female adult or adolescent		
White, not Hispanic	14	74%	Injection drug use	-	-
Black, not Hispanic	4	21%	Heterosexual contact	2	67%
Hispanic	-	-	Blood product	-	-
Asian/Pacific Islander	-	-	Other/unknown	1	33%
American Indian/Alaska Native	-	-	Subtotal	3	100%
Unknown	1	5%	Child (<13 yrs at diagnosis)		
			Perinatal	1	100%
			Other	-	-
			Subtotal	1	100%
Total	19				

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Knox County, Ohio

Cumulative cases of AIDS diagnosed through 2003, by selected characteristics and transmission category

		Cumulative AIDS Diagnoses			
		No.	%		
				No.	%
Gender				Transmission category	
Male	15	75%		Male adult or adolescent	
Female	5	25%		Male-to-male sexual contact	8 57%
Unknown	-	-		Injection drug use (IDU)	- -
				Male-to-male sexual contact and IDU	1 7%
Age				Heterosexual contact	1 7%
<13	1	5%		Blood product	1 7%
13-19	-	-		Other/unknown	3 21%
20-24	2	10%		Subtotal	14 100%
25-29	3	15%		Female adult or adolescent	
30-39	8	40%		Injection drug use	1 20%
40-49	4	20%		Heterosexual contact	2 40%
50-64	1	5%		Blood product	1 20%
65+	1	5%		Other/unknown	1 20%
				Subtotal	5 100%
Race/Ethnicity				Child (<13 yrs at diagnosis)	
White, not Hispanic	16	80%		Perinatal	1 100%
Black, not Hispanic	4	20%		Other	- -
Hispanic	-	-		Subtotal	1 100%
Asian/Pacific Islander	-	-			
American Indian/Alaska Native	-	-			
Unknown	-	-			
Race/Gender					
White, not Hispanic Male	13	65%			
White, not Hispanic Female	3	15%			
Black, not Hispanic Male	2	10%			
Black, not Hispanic Female	2	10%			
Other Male	-	-			
Other Female	-	-			
Unknown	-	-			
Total	20				

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.