

Logan County, Ohio

Reported persons living with HIV/AIDS as of Dec. 31, 2003, by selected characteristics and transmission category

Reported Persons Living with HIV/AIDS						
			No.	%		
			No.	%	No.	%
Current disease status					Race/Gender	
HIV (not AIDS)	9	43%	White, not Hispanic Male	14	67%	
AIDS	12	57%	White, not Hispanic Female	6	29%	
			Black, not Hispanic Male	-	-	
Gender			Black, not Hispanic Female	1	5%	
Male	14	67%	Other Male	-	-	
Female	7	33%	Other Female	-	-	
Unknown	-	-	Unknown	-	-	
Age			Transmission category			
<13	-	-	Male adult or adolescent			
13-19	-	-	Male-to-male sexual contact	7	50%	
20-24	-	-	Injection drug use (IDU)	1	7%	
25-29	-	-	Male-to-male sexual contact and IDU	1	7%	
30-39	8	38%			-	
40-49	6	29%	Heterosexual contact	1	7%	
50-64	7	33%	Blood product	-	-	
65+	-	-	Other/unknown	4	29%	
			Subtotal	14	100%	
Race/Ethnicity			Female adult or adolescent			
White, not Hispanic	20	95%	Injection drug use	1	14%	
Black, not Hispanic	1	5%	Heterosexual contact	1	14%	
Hispanic	-	-	Blood product	-	-	
Asian/Pacific Islander	-	-	Other/unknown	5	71%	
American Indian/Alaska Native	-	-	Subtotal	7	100%	
Unknown	-	-	Child (<13 yrs at diagnosis)			
			Perinatal	-	-	
			Other	-	-	
			Subtotal	-	-	
Total	21					

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Logan County, Ohio

Cumulative cases of AIDS diagnosed through 2003, by selected characteristics and transmission category

		Cumulative AIDS Diagnoses			
		No.	%		
		No.	%	No.	%
Gender				Transmission category	
Male	17	81%	Male adult or adolescent		
Female	4	19%	Male-to-male sexual contact	9	53%
Unknown	-	-	Injection drug use (IDU)	1	6%
			Male-to-male sexual contact and IDU	2	12%
Age			Heterosexual contact	1	6%
<13	-	-	Blood product	1	6%
13-19	-	-	Other/unknown	3	18%
20-24	-	-	Subtotal	17	100%
25-29	4	19%	Female adult or adolescent		
30-39	9	43%	Injection drug use	1	25%
40-49	5	24%	Heterosexual contact	1	25%
50-64	3	14%	Blood product	-	-
65+	-	-	Other/unknown	2	50%
			Subtotal	4	100%
Race/Ethnicity			Child (<13 yrs at diagnosis)		
White, not Hispanic	21	100%	Perinatal	-	-
Black, not Hispanic	-	-	Other	-	-
Hispanic	-	-	Subtotal	-	-
Asian/Pacific Islander	-	-			
American Indian/Alaska Native	-	-			
Unknown	-	-			
Race/Gender					
White, not Hispanic Male	17	81%			
White, not Hispanic Female	4	19%			
Black, not Hispanic Male	-	-			
Black, not Hispanic Female	-	-			
Other Male	-	-			
Other Female	-	-			
Unknown	-	-			
Total	21				

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.