

Medina County, Ohio

Reported persons living with HIV/AIDS as of Dec. 31, 2003, by selected characteristics and transmission category

Reported Persons Living with HIV/AIDS						
			No.	%		
			No.	%	No.	%
Current disease status					Race/Gender	
HIV (not AIDS)	8	33%	White, not Hispanic Male	16	67%	
AIDS	16	67%	White, not Hispanic Female	3	13%	
			Black, not Hispanic Male	4	17%	
Gender			Black, not Hispanic Female	1	4%	
Male	20	83%	Other Male	-	-	
Female	4	17%	Other Female	-	-	
Unknown	-	-	Unknown	-	-	
Age			Transmission category			
<13	-	-	Male adult or adolescent			
13-19	-	-	Male-to-male sexual contact	9	45%	
20-24	-	-	Injection drug use (IDU)	-	-	
25-29	1	4%	Male-to-male sexual contact and IDU	1	5%	
30-39	7	29%			-	
40-49	11	46%	Heterosexual contact	2	10%	
50-64	4	17%	Blood product	2	10%	
65+	1	4%	Other/unknown	6	30%	
			Subtotal	20	100%	
Race/Ethnicity			Female adult or adolescent			
White, not Hispanic	19	79%	Injection drug use	-	-	
Black, not Hispanic	5	21%	Heterosexual contact	3	75%	
Hispanic	-	-	Blood product	-	-	
Asian/Pacific Islander	-	-	Other/unknown	1	25%	
American Indian/Alaska Native	-	-	Subtotal	4	100%	
Unknown	-	-	Child (<13 yrs at diagnosis)			
			Perinatal	-	-	
			Other	-	-	
			Subtotal	-	-	
Total	24					

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Medina County, Ohio

Cumulative cases of AIDS diagnosed through 2003, by selected characteristics and transmission category

Cumulative AIDS Diagnoses					
	No.	%		No.	%
Gender			Transmission category		
Male	29	91%	Male adult or adolescent		
Female	3	9%	Male-to-male sexual contact	21	72%
Unknown	-	-	Injection drug use (IDU)	1	3%
			Male-to-male sexual contact and IDU	1	3%
Age			Heterosexual contact	2	7%
<13	-	-	Blood product	2	7%
13-19	-	-	Other/unknown	2	7%
20-24	2	6%	Subtotal	29	100%
25-29	5	16%	Female adult or adolescent		
30-39	18	56%	Injection drug use	-	-
40-49	3	9%	Heterosexual contact	3	100%
50-64	3	9%	Blood product	-	-
65+	1	3%	Other/unknown	-	-
			Subtotal	3	100%
Race/Ethnicity			Child (<13 yrs at diagnosis)		
White, not Hispanic	29	91%	Perinatal	-	-
Black, not Hispanic	1	3%	Other	-	-
Hispanic	2	6%	Subtotal	-	-
Asian/Pacific Islander	-	-			
American Indian/Alaska Native	-	-			
Unknown	-	-			
Race/Gender					
White, not Hispanic Male	27	84%			
White, not Hispanic Female	2	6%			
Black, not Hispanic Male	1	3%			
Black, not Hispanic Female	-	-			
Other Male	1	3%			
Other Female	1	3%			
Unknown	-	-			
Total	32				

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.