

Mercer County, Ohio

Cumulative cases of AIDS diagnosed through 2003, by selected characteristics and transmission category

		Cumulative AIDS Diagnoses			
		No.	%		
				No.	%
Gender		Transmission category			
Male	14	100%	Male adult or adolescent		
Female	-	-	Male-to-male sexual contact	8	57%
Unknown	-	-	Injection drug use (IDU)	-	-
			Male-to-male sexual contact and IDU	1	7%
Age				Heterosexual contact	-
<13	-	-	Blood product	3	21%
13-19	-	-	Other/unknown	2	14%
20-24	-	-	Subtotal	14	100%
25-29	2	14%	Female adult or adolescent		
30-39	6	43%	Injection drug use	-	-
40-49	2	14%	Heterosexual contact	-	-
50-64	4	29%	Blood product	-	-
65+	-	-	Other/unknown	-	-
			Subtotal	-	-
Race/Ethnicity		Child (<13 yrs at diagnosis)			
White, not Hispanic	14	100%	Perinatal	-	-
Black, not Hispanic	-	-	Other	-	-
Hispanic	-	-	Subtotal	-	-
Asian/Pacific Islander	-	-			
American Indian/Alaska Native	-	-			
Unknown	-	-			
Race/Gender					
White, not Hispanic Male	14	100%			
White, not Hispanic Female	-	-			
Black, not Hispanic Male	-	-			
Black, not Hispanic Female	-	-			
Other Male	-	-			
Other Female	-	-			
Unknown	-	-			
Total	14				

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.