

## Morrow County, Ohio

### Cumulative cases of AIDS diagnosed through 2003, by selected characteristics and transmission category

Cumulative AIDS Diagnoses					
	No.	%		No.	%
<b>Gender</b>			<b>Transmission category</b>		
Male	10	91%	<b>Male adult or adolescent</b>		
Female	1	9%	Male-to-male sexual contact	6	60%
Unknown	-	-	Injection drug use (IDU)	1	10%
			Male-to-male sexual contact and IDU	-	-
<b>Age</b>			Heterosexual contact	2	20%
<13	-	-	Blood product	-	-
13-19	-	-	Other/unknown	1	10%
20-24	-	-	Subtotal	10	100%
25-29	1	9%	<b>Female adult or adolescent</b>		
30-39	4	36%	Injection drug use	1	100%
40-49	5	45%	Heterosexual contact	-	-
50-64	1	9%	Blood product	-	-
65+	-	-	Other/unknown	-	-
			Subtotal	1	100%
<b>Race/Ethnicity</b>			<b>Child (&lt;13 yrs at diagnosis)</b>		
White, not Hispanic	11	100%	Perinatal	-	-
Black, not Hispanic	-	-	Other	-	-
Hispanic	-	-	Subtotal	-	-
Asian/Pacific Islander	-	-			
American Indian/Alaska Native	-	-			
Unknown	-	-			
<b>Race/Gender</b>					
White, not Hispanic Male	10	91%			
White, not Hispanic Female	1	9%			
Black, not Hispanic Male	-	-			
Black, not Hispanic Female	-	-			
Other Male	-	-			
Other Female	-	-			
Unknown	-	-			
<b>Total</b>	<b>11</b>				

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.