

Muskingum County, Ohio

Reported persons living with HIV/AIDS as of Dec. 31, 2003, by current disease status and selected characteristics

	Current Disease Status						
	HIV/AIDS			HIV (not AIDS)		AIDS	
	Rate ^a	No.	%	No.	%	No.	%
Gender							
Male	108.6	44	73%	20	61%	24	89%
Female	36.3	16	27%	13	39%	3	11%
Unknown	*	-	-	-	-	-	-
Age (as of 12/2003)							
<13	*	-	-	-	-	-	-
13-19	*	-	-	-	-	-	-
20-24	*	3	5%	2	6%	1	4%
25-29	*	1	2%	1	3%	-	-
30-39	186.7	22	37%	10	30%	12	44%
40-49	168.0	21	35%	15	45%	6	22%
50-64	90.7	12	20%	5	15%	7	26%
65+	*	1	2%	-	-	1	4%
Race/Ethnicity							
White, not Hispanic	57.7	46	77%	25	76%	21	78%
Black, not Hispanic	281.1	11	18%	5	15%	6	22%
Hispanic	*	-	-	-	-	-	-
Asian/Pacific Islander	*	-	-	-	-	-	-
American Indian/Alaska Native	*	1	2%	1	3%	-	-
Unknown	*	2	3%	2	6%	-	-
Race/Gender							
White, not Hispanic Male	86.7	33	55%	15	45%	18	67%
White, not Hispanic Female	31.2	13	22%	10	30%	3	11%
Black, not Hispanic Male	404.7	8	13%	2	6%	6	22%
Black, not Hispanic Female	*	3	5%	3	9%	-	-
Other Male	*	1	2%	1	3%	-	-
Other Female	*	-	-	-	-	-	-
Unknown	*	2	3%	2	6%	-	-
Total	70.9	60		33		27	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with HIV/AIDS per 100,000 population calculated using 2000 census data.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Muskingum County, Ohio

Reported persons living with HIV/AIDS as of Dec. 31, 2003, by current disease status and transmission category

Transmission category	HIV/AIDS		Current Disease Status			
			HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	18	42%	4	20%	14	61%
Injection drug use (IDU)	3	7%	-	-	3	13%
Male-to-male sexual contact and IDU	-	-	-	-	-	-
Heterosexual contact	1	2%	1	5%	-	-
Blood product	2	5%	-	-	2	9%
Other/unknown	19	44%	15	75%	4	17%
Subtotal	43	100%	20	100%	23	100%
Female adult or adolescent						
Injection drug use	-	-	-	-	-	-
Heterosexual contact	7	44%	5	38%	2	67%
Blood product	-	-	-	-	-	-
Other/unknown	9	56%	8	62%	1	33%
Subtotal	16	100%	13	100%	3	100%
Child (<13 yrs at diagnosis)						
Perinatal	-	-	-	-	-	-
Other	1	100%	-	-	1	100%
Subtotal	1	100%	-	-	1	100%
Total	60		33		27	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Muskingum County, Ohio

Reported cases of HIV infection, by year of diagnosis and selected characteristics

	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
Gender				
Male	10	71%	11	69%
Female	4	29%	5	31%
Unknown	-	-	-	-
Age				
<13	-	-	-	-
13-19	-	-	-	-
20-24	-	-	1	6%
25-29	4	29%	1	6%
30-39	6	43%	6	38%
40-49	4	29%	5	31%
50-64	-	-	2	13%
65+	-	-	1	6%
Race/Ethnicity				
White, not Hispanic	11	79%	12	75%
Black, not Hispanic	2	14%	4	25%
Hispanic	-	-	-	-
Asian/Pacific Islander	-	-	-	-
American Indian/Alaska Native	-	-	-	-
Unknown	1	7%	-	-
Race/Gender				
White, not Hispanic Male	7	50%	8	50%
White, not Hispanic Female	4	29%	4	25%
Black, not Hispanic Male	2	14%	3	19%
Black, not Hispanic Female	-	-	1	6%
Other Male	-	-	-	-
Other Female	-	-	-	-
Unknown	1	7%	-	-
Total	14		16	

Note: HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Muskingum County, Ohio

Reported cases of HIV infection, by year of diagnosis and transmission category

Transmission category	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
Male adult or adolescent				
Male-to-male sexual contact	6	60%	2	18%
Injection drug use (IDU)	-	-	1	9%
Male-to-male sexual contact and IDU	-	-	-	-
Heterosexual contact	2	20%	-	-
Blood product	-	-	-	-
Other/unknown	2	20%	8	73%
Subtotal	10	100%	11	100%
Female adult or adolescent				
Injection drug use	1	25%	-	-
Heterosexual contact	1	25%	1	20%
Blood product	-	-	-	-
Other/unknown	2	50%	4	80%
Subtotal	4	100%	5	100%
Child (<13 yrs at diagnosis)				
Perinatal	-	-	-	-
Other	-	-	-	-
Subtotal	-	-	-	-
Total	14		16	

Note: HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Muskingum County, Ohio

Cumulative cases of AIDS diagnosed through 2003, by selected characteristics and transmission category

	Cumulative AIDS Diagnoses				
	No.	%		No.	%
Gender			Transmission category		
Male	40	91%	Male adult or adolescent		
Female	4	9%	Male-to-male sexual contact	25	64%
Unknown	-	-	Injection drug use (IDU)	4	10%
			Male-to-male sexual contact and IDU	1	3%
Age			Heterosexual contact	-	-
<13	-	-	Blood product	2	5%
13-19	1	2%	Other/unknown	7	18%
20-24	-	-	Subtotal	39	100%
25-29	6	14%	Female adult or adolescent		
30-39	17	39%	Injection drug use	1	25%
40-49	14	32%	Heterosexual contact	3	75%
50-64	5	11%	Blood product	-	-
65+	1	2%	Other/unknown	-	-
			Subtotal	4	100%
Race/Ethnicity			Child (<13 yrs at diagnosis)		
White, not Hispanic	40	91%	Perinatal	-	-
Black, not Hispanic	4	9%	Other	1	100%
Hispanic	-	-	Subtotal	1	100%
Asian/Pacific Islander	-	-			
American Indian/Alaska Native	-	-			
Unknown	-	-			
Race/Gender					
White, not Hispanic Male	36	82%			
White, not Hispanic Female	4	9%			
Black, not Hispanic Male	4	9%			
Black, not Hispanic Female	-	-			
Other Male	-	-			
Other Female	-	-			
Unknown	-	-			
Total	44				

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.