

Ohio

Reported persons living with HIV/AIDS as of Dec. 31, 2003, by current disease status and selected characteristics

	Current Disease Status						
	HIV/AIDS			HIV (not AIDS)		AIDS	
	Rate ^a	No.	%	No.	%	No.	%
Gender							
Male	210.3	11,591	80%	6,002	77%	5,589	83%
Female	49.4	2,884	20%	1,774	23%	1,110	17%
Unknown	*	2	<1%	2	<1%	-	-
Age (as of 12/2003)							
<13	4.9	102	1%	62	1%	40	1%
13-19	7.3	84	1%	54	1%	30	<1%
20-24	46.4	338	2%	279	4%	59	1%
25-29	111.6	821	6%	598	8%	223	3%
30-39	282.3	4,709	33%	2,712	35%	1,997	30%
40-49	326.1	5,728	40%	2,848	37%	2,880	43%
50-64	141.5	2,463	17%	1,125	14%	1,338	20%
65+	15.4	232	2%	100	1%	132	2%
Race/Ethnicity							
White, not Hispanic	75.5	7,268	50%	3,695	48%	3,573	53%
Black, not Hispanic	450.1	6,031	42%	3,308	43%	2,723	41%
Hispanic	316.9	688	5%	336	4%	352	5%
Asian/Pacific Islander	37.1	54	<1%	25	<1%	29	<1%
American Indian/Alaska Native	108.1	31	<1%	19	<1%	12	<1%
Unknown	*	405	3%	395	5%	10	<1%
Race/Gender							
White, not Hispanic Male	134.3	6,283	43%	3,086	40%	3,197	48%
White, not Hispanic Female	19.9	985	7%	609	8%	376	6%
Black, not Hispanic Male	698.6	4,436	31%	2,347	30%	2,089	31%
Black, not Hispanic Female	226.2	1,594	11%	960	12%	634	9%
Other Male	279.5	552	4%	258	3%	294	4%
Other Female	113.9	221	2%	122	2%	99	1%
Unknown	*	406	3%	396	5%	10	<1%
Total	127.5	14,477		7,778		6,699	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with HIV/AIDS per 100,000 population calculated using 2000 census data.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Ohio

Reported persons living with HIV/AIDS as of Dec. 31, 2003, by current disease status and transmission category

Transmission category	HIV/AIDS		Current Disease Status			
			HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	5,731	50%	2,406	40%	3,325	60%
Injection drug use (IDU)	909	8%	398	7%	511	9%
Male-to-male sexual contact and IDU	579	5%	213	4%	366	7%
Heterosexual contact	421	4%	191	3%	230	4%
Blood product	103	1%	40	1%	63	1%
Other/unknown	3,766	33%	2,713	46%	1,053	19%
Subtotal	11,509	100%	5,961	100%	5,548	100%
Female adult or adolescent						
Injection drug use	388	14%	170	10%	218	20%
Heterosexual contact	1,038	37%	563	33%	475	44%
Blood product	25	1%	11	1%	14	1%
Other/unknown	1,346	48%	980	57%	366	34%
Subtotal	2,797	100%	1,724	100%	1,073	100%
Child (<13 yrs at diagnosis)						
Perinatal	132	78%	73	80%	59	76%
Other	37	22%	18	20%	19	24%
Subtotal	169	100%	91	100%	78	100%
Total	14,477		7,776		6,699	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Ohio

Reported cases of HIV infection, by year of diagnosis and selected characteristics

	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
Gender				
Male	2,385	79%	2,313	77%
Female	624	21%	688	23%
Unknown	1	<1%	-	-
Age				
<13	26	1%	20	1%
13-19	75	2%	71	2%
20-24	250	8%	263	9%
25-29	414	14%	368	12%
30-39	1,251	42%	1,119	37%
40-49	734	24%	840	28%
50-64	235	8%	297	10%
65+	25	1%	23	1%
Race/Ethnicity				
White, not Hispanic	1,467	49%	1,418	47%
Black, not Hispanic	1,329	44%	1,286	43%
Hispanic	118	4%	130	4%
Asian/Pacific Islander	14	<1%	15	<1%
American Indian/Alaska Native	8	<1%	7	<1%
Unknown	74	2%	145	5%
Race/Gender				
White, not Hispanic Male	1,254	42%	1,208	40%
White, not Hispanic Female	213	7%	210	7%
Black, not Hispanic Male	972	32%	876	29%
Black, not Hispanic Female	357	12%	410	14%
Other Male	100	3%	113	4%
Other Female	40	1%	39	1%
Unknown	74	2%	145	5%
Total	3010		3001	

Note: Total HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Ohio

Reported cases of HIV infection, by year of diagnosis and transmission category

Transmission category	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
Male adult or adolescent				
Male-to-male sexual contact	1,130	48%	746	32%
Injection drug use (IDU)	151	6%	83	4%
Male-to-male sexual contact and IDU	70	3%	32	1%
Heterosexual contact	90	4%	76	3%
Blood product	4	<1%	6	<1%
Other/unknown	927	39%	1,363	59%
Subtotal	2,372	100%	2,306	100%
Female adult or adolescent				
Injection drug use	47	8%	29	4%
Heterosexual contact	216	35%	137	20%
Blood product	8	1%	3	<1%
Other/unknown	340	56%	506	75%
Subtotal	611	100%	675	100%
Child (<13 yrs at diagnosis)				
Perinatal	21	81%	17	85%
Other	5	19%	3	15%
Subtotal	26	-	20	-
Total	3,010		3,001	

Note: Total HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Ohio

Reported cases of AIDS, by year of diagnosis and selected characteristics

	Year of AIDS Diagnosis				Cumulative	
	1998 - 2000		2001 - 2003			
	No.	%	No.	%	No.	%
Gender						
Male	1,385	82%	1,358	80%	11,475	87%
Female	313	18%	349	20%	1,703	13%
Unknown	-	-	-	-	-	-
Age						
<13	8	<1%	9	1%	130	1%
13-19	9	1%	12	1%	92	1%
20-24	45	3%	62	4%	517	4%
25-29	176	10%	136	8%	1,882	14%
30-39	791	47%	711	42%	5,967	45%
40-49	499	29%	560	33%	3,324	25%
50-64	151	9%	200	12%	1,127	9%
65+	19	1%	17	1%	139	1%
Race/Ethnicity						
White, not Hispanic	843	50%	837	49%	7,892	60%
Black, not Hispanic	775	46%	773	45%	4,759	36%
Hispanic	61	4%	84	5%	467	4%
Asian/Pacific Islander	9	1%	5	<1%	35	<1%
American Indian/Alaska Native	7	<1%	1	<1%	14	<1%
Unknown	3	<1%	7	<1%	11	<1%
Race/Gender						
White, not Hispanic Male	742	44%	721	42%	7,261	55%
White, not Hispanic Female	101	6%	116	7%	631	5%
Black, not Hispanic Male	583	34%	565	33%	3,809	29%
Black, not Hispanic Female	192	11%	208	12%	950	7%
Other Male	57	3%	66	4%	395	3%
Other Female	20	1%	24	1%	121	1%
Unknown	3	<1%	7	<1%	11	<1%
Total	1,698		1,707		13,178	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Ohio

Reported cases of AIDS, by year of diagnosis and transmission category

Transmission category	Year of AIDS Diagnosis				Cumulative	
	1998 - 2000		2001 - 2003			
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	796	58%	609	45%	7,696	68%
Injection drug use (IDU)	112	8%	83	6%	943	8%
Male-to-male sexual contact and IDU	64	5%	56	4%	751	7%
Heterosexual contact	69	5%	50	4%	368	3%
Blood product	8	1%	3	<1%	254	2%
Other/unknown	333	24%	550	41%	1,380	12%
Subtotal	1,382	100%	1,351	100%	11,392	100%
Female adult or adolescent						
Injection drug use	55	18%	38	11%	374	23%
Heterosexual contact	125	41%	105	31%	748	46%
Blood product	5	2%	2	1%	53	3%
Other/unknown	122	40%	196	57%	455	28%
Subtotal	307	100%	341	100%	1,630	100%
Child (<13 yrs at diagnosis)						
Perinatal	7	78%	10	67%	115	74%
Other	2	22%	5	33%	41	26%
Subtotal	9	100%	15	100%	156	100%
Total	1,698		1,707		13,178	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.