

## Ottawa County, Ohio

### Reported persons living with HIV/AIDS as of Dec. 31, 2003, by selected characteristics and transmission category

Reported Persons Living with HIV/AIDS					
	No.	%		No.	%
<b>Current disease status</b>			<b>Race/Gender</b>		
HIV (not AIDS)	10	56%	White, not Hispanic Male	10	56%
AIDS	8	44%	White, not Hispanic Female	3	17%
			Black, not Hispanic Male	3	17%
<b>Gender</b>			Black, not Hispanic Female	-	-
Male	15	83%	Other Male	2	11%
Female	3	17%	Other Female	-	-
Unknown	-	-	Unknown	-	-
<b>Age</b>			<b>Transmission category</b>		
<13	-	-	<b>Male adult or adolescent</b>		
13-19	-	-	Male-to-male sexual contact	9	60%
20-24	-	-	Injection drug use (IDU)	2	13%
25-29	1	6%	Male-to-male sexual contact and IDU	-	-
30-39	5	28%	Heterosexual contact	1	7%
40-49	6	33%	Blood product	-	-
50-64	6	33%	Other/unknown	3	20%
65+	-	-	Subtotal	15	100%
<b>Race/Ethnicity</b>			<b>Female adult or adolescent</b>		
White, not Hispanic	13	72%	Injection drug use	1	33%
Black, not Hispanic	3	17%	Heterosexual contact	1	33%
Hispanic	2	11%	Blood product	-	-
Asian/Pacific Islander	-	-	Other/unknown	1	33%
American Indian/Alaska Native	-	-	Subtotal	3	100%
Unknown	-	-	<b>Child (&lt;13 yrs at diagnosis)</b>		
			Perinatal	-	-
			Other	-	-
			Subtotal	-	-
<b>Total</b>	<b>18</b>				

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ottawa County, Ohio

### Cumulative cases of AIDS diagnosed through 2003, by selected characteristics and transmission category

		Cumulative AIDS Diagnoses			
		No.	%		
		No.	%	No.	%
<b>Gender</b>				<b>Transmission category</b>	
Male	11	85%		<b>Male adult or adolescent</b>	
Female	2	15%		Male-to-male sexual contact	7 64%
Unknown	-	-		Injection drug use (IDU)	- -
				Male-to-male sexual contact and IDU	- -
<b>Age</b>				Heterosexual contact	1 9%
<13	-	-		Blood product	- -
13-19	-	-		Other/unknown	3 27%
20-24	1	8%		Subtotal	11 100%
25-29	1	8%		<b>Female adult or adolescent</b>	
30-39	1	8%		Injection drug use	- -
40-49	6	46%		Heterosexual contact	- -
50-64	4	31%		Blood product	1 50%
65+	-	-		Other/unknown	1 50%
				Subtotal	2 100%
<b>Race/Ethnicity</b>				<b>Child (&lt;13 yrs at diagnosis)</b>	
White, not Hispanic	11	85%		Perinatal	- -
Black, not Hispanic	2	15%		Other	- -
Hispanic	-	-		Subtotal	- -
Asian/Pacific Islander	-	-			
American Indian/Alaska Native	-	-			
Unknown	-	-			
<b>Race/Gender</b>					
White, not Hispanic Male	9	69%			
White, not Hispanic Female	2	15%			
Black, not Hispanic Male	2	15%			
Black, not Hispanic Female	-	-			
Other Male	-	-			
Other Female	-	-			
Unknown	-	-			
<b>Total</b>	<b>13</b>				

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.