

## Perry County, Ohio

### Reported persons living with HIV/AIDS as of Dec. 31, 2003, by selected characteristics and transmission category

Reported Persons Living with HIV/AIDS					
	No.	%		No.	%
<b>Current disease status</b>			<b>Race/Gender</b>		
HIV (not AIDS)	7	58%	White, not Hispanic Male	10	83%
AIDS	5	42%	White, not Hispanic Female	2	17%
			Black, not Hispanic Male	-	-
<b>Gender</b>			Black, not Hispanic Female	-	-
Male	10	83%	Other Male	-	-
Female	2	17%	Other Female	-	-
Unknown	-	-	Unknown	-	-
<b>Age</b>			<b>Transmission category</b>		
<13	1	8%	<b>Male adult or adolescent</b>		
13-19	-	-	Male-to-male sexual contact	2	22%
20-24	-	-	Injection drug use (IDU)	-	-
25-29	3	25%	Male-to-male sexual contact and IDU	-	-
30-39	5	42%	Heterosexual contact	2	22%
40-49	2	17%	Blood product	1	11%
50-64	1	8%	Other/unknown	4	44%
65+	-	-	Subtotal	9	100%
<b>Race/Ethnicity</b>			<b>Female adult or adolescent</b>		
White, not Hispanic	12	100%	Injection drug use	-	-
Black, not Hispanic	-	-	Heterosexual contact	1	50%
Hispanic	-	-	Blood product	-	-
Asian/Pacific Islander	-	-	Other/unknown	1	50%
American Indian/Alaska Native	-	-	Subtotal	2	100%
Unknown	-	-	<b>Child (&lt;13 yrs at diagnosis)</b>		
			Perinatal	1	100%
			Other	-	-
			Subtotal	1	100%
<b>Total</b>	<b>12</b>				

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Perry County, Ohio

### Cumulative cases of AIDS diagnosed through 2003, by selected characteristics and transmission category

Cumulative AIDS Diagnoses					
	No.	%		No.	%
<b>Gender</b>			<b>Transmission category</b>		
Male	11	92%	<b>Male adult or adolescent</b>		
Female	1	8%	Male-to-male sexual contact	6	55%
Unknown	-	-	Injection drug use (IDU)	-	-
			Male-to-male sexual contact and IDU	-	-
<b>Age</b>			Heterosexual contact	1	9%
<13	-	-	Blood product	3	27%
13-19	-	-	Other/unknown	1	9%
20-24	1	8%	Subtotal	11	100%
25-29	5	42%	<b>Female adult or adolescent</b>		
30-39	4	33%	Injection drug use	1	100%
40-49	1	8%	Heterosexual contact	-	-
50-64	1	8%	Blood product	-	-
65+	-	-	Other/unknown	-	-
			Subtotal	1	100%
<b>Race/Ethnicity</b>			<b>Child (&lt;13 yrs at diagnosis)</b>		
White, not Hispanic	12	100%	Perinatal	-	-
Black, not Hispanic	-	-	Other	-	-
Hispanic	-	-	Subtotal	-	-
Asian/Pacific Islander	-	-			
American Indian/Alaska Native	-	-			
Unknown	-	-			
<b>Race/Gender</b>					
White, not Hispanic Male	11	92%			
White, not Hispanic Female	1	8%			
Black, not Hispanic Male	-	-			
Black, not Hispanic Female	-	-			
Other Male	-	-			
Other Female	-	-			
Unknown	-	-			
<b>Total</b>	<b>12</b>				

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.