

Pickaway County, Ohio

Reported persons living with HIV/AIDS as of Dec. 31, 2003, by selected characteristics and transmission category

Reported Persons Living with HIV/AIDS					
	No.	%		No.	%
Current disease status			Race/Gender		
HIV (not AIDS)	13	65%	White, not Hispanic Male	11	55%
AIDS	7	35%	White, not Hispanic Female	1	5%
			Black, not Hispanic Male	5	25%
Gender			Black, not Hispanic Female	-	-
Male	17	85%	Other Male	1	5%
Female	3	15%	Other Female	2	10%
Unknown	-	-	Unknown	-	-
Age			Transmission category		
<13	-	-	Male adult or adolescent		
13-19	1	5%	Male-to-male sexual contact	7	41%
20-24	1	5%	Injection drug use (IDU)	2	12%
25-29	-	-	Male-to-male sexual contact and IDU	-	-
30-39	8	40%	Heterosexual contact	-	-
40-49	5	25%	Blood product	1	6%
50-64	4	20%	Other/unknown	7	41%
65+	1	5%	Subtotal	17	100%
Race/Ethnicity			Female adult or adolescent		
White, not Hispanic	12	60%	Injection drug use	-	-
Black, not Hispanic	5	25%	Heterosexual contact	1	50%
Hispanic	3	15%	Blood product	-	-
Asian/Pacific Islander	-	-	Other/unknown	1	50%
American Indian/Alaska Native	-	-	Subtotal	2	100%
Unknown	-	-	Child (<13 yrs at diagnosis)		
			Perinatal	1	100%
			Other	-	-
			Subtotal	1	100%
Total	20				

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Pickaway County, Ohio

Cumulative cases of AIDS diagnosed through 2003, by selected characteristics and transmission category

		Cumulative AIDS Diagnoses			
		No.	%		
				No.	%
Gender				Transmission category	
Male	20	87%	Male adult or adolescent		
Female	3	13%	Male-to-male sexual contact	11	55%
Unknown	-	-	Injection drug use (IDU)	4	20%
			Male-to-male sexual contact and IDU	-	-
Age			Heterosexual contact	2	10%
<13	-	-	Blood product	1	5%
13-19	-	-	Other/unknown	2	10%
20-24	1	4%	Subtotal	20	100%
25-29	5	22%	Female adult or adolescent		
30-39	11	48%	Injection drug use	-	-
40-49	4	17%	Heterosexual contact	2	67%
50-64	2	9%	Blood product	-	-
65+	-	-	Other/unknown	1	33%
			Subtotal	3	100%
Race/Ethnicity			Child (<13 yrs at diagnosis)		
White, not Hispanic	17	74%	Perinatal	-	-
Black, not Hispanic	3	13%	Other	-	-
Hispanic	3	13%	Subtotal	-	-
Asian/Pacific Islander	-	-			
American Indian/Alaska Native	-	-			
Unknown	-	-			
Race/Gender					
White, not Hispanic Male	15	65%			
White, not Hispanic Female	2	9%			
Black, not Hispanic Male	3	13%			
Black, not Hispanic Female	-	-			
Other Male	2	9%			
Other Female	1	4%			
Unknown	-	-			
Total	23				

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.