

Pike County, Ohio

Reported persons living with HIV/AIDS as of Dec. 31, 2003, by selected characteristics and transmission category

Reported Persons Living with HIV/AIDS						
			No.	%		
			No.	%	No.	%
Current disease status					Race/Gender	
HIV (not AIDS)	7	50%	White, not Hispanic Male	8	57%	
AIDS	7	50%	White, not Hispanic Female	3	21%	
			Black, not Hispanic Male	1	7%	
Gender			Black, not Hispanic Female	-	-	
Male	10	71%	Other Male	1	7%	
Female	4	29%	Other Female	-	-	
Unknown	-	-	Unknown	1	7%	
Age			Transmission category			
<13	-	-	Male adult or adolescent			
13-19	-	-	Male-to-male sexual contact	6	60%	
20-24	1	7%	Injection drug use (IDU)	1	10%	
25-29	1	7%	Male-to-male sexual contact and IDU	-	-	
30-39	5	36%	Heterosexual contact	-	-	
40-49	5	36%	Blood product	-	-	
50-64	2	14%	Other/unknown	3	30%	
65+	-	-	Subtotal	10	100%	
Race/Ethnicity			Female adult or adolescent			
White, not Hispanic	11	79%	Injection drug use	-	-	
Black, not Hispanic	1	7%	Heterosexual contact	3	75%	
Hispanic	-	-	Blood product	-	-	
Asian/Pacific Islander	-	-	Other/unknown	1	25%	
American Indian/Alaska Native	1	7%	Subtotal	4	100%	
Unknown	1	7%	Child (<13 yrs at diagnosis)			
			Perinatal	-	-	
			Other	-	-	
			Subtotal	-	-	
Total	14					

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Pike County, Ohio

Cumulative cases of AIDS diagnosed through 2003, by selected characteristics and transmission category

	Cumulative AIDS Diagnoses				
	No.	%		No.	%
Gender			Transmission category		
Male	10	91%	Male adult or adolescent		
Female	1	9%	Male-to-male sexual contact	8	80%
Unknown	-	-	Injection drug use (IDU)	-	-
			Male-to-male sexual contact and IDU	-	-
Age			Heterosexual contact	-	-
<13	-	-	Blood product	-	-
13-19	1	9%	Other/unknown	2	20%
20-24	-	-	Subtotal	10	100%
25-29	4	36%	Female adult or adolescent		
30-39	2	18%	Injection drug use	1	100%
40-49	2	18%	Heterosexual contact	-	-
50-64	1	9%	Blood product	-	-
65+	1	9%	Other/unknown	-	-
			Subtotal	1	100%
Race/Ethnicity			Child (<13 yrs at diagnosis)		
White, not Hispanic	10	91%	Perinatal	-	-
Black, not Hispanic	1	9%	Other	-	-
Hispanic	-	-	Subtotal	-	-
Asian/Pacific Islander	-	-			
American Indian/Alaska Native	-	-			
Unknown	-	-			
Race/Gender					
White, not Hispanic Male	9	82%			
White, not Hispanic Female	1	9%			
Black, not Hispanic Male	1	9%			
Black, not Hispanic Female	-	-			
Other Male	-	-			
Other Female	-	-			
Unknown	-	-			
Total	11				

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.