

## Portage County, Ohio

### Reported persons living with HIV/AIDS as of Dec. 31, 2003, by current disease status and selected characteristics

	Current Disease Status						
	HIV/AIDS			HIV (not AIDS)		AIDS	
	Rate <sup>a</sup>	No.	%	No.	%	No.	%
<b>Gender</b>							
Male	66.0	49	86%	21	78%	28	93%
Female	10.3	8	14%	6	22%	2	7%
Unknown	*	-	-	-	-	-	-
<b>Age (as of 12/2003)</b>							
<13	*	-	-	-	-	-	-
13-19	*	-	-	-	-	-	-
20-24	*	2	4%	1	4%	1	3%
25-29	*	1	2%	1	4%	-	-
30-39	72.9	16	28%	9	33%	7	23%
40-49	77.3	18	32%	9	33%	9	30%
50-64	78.7	18	32%	6	22%	12	40%
65+	*	2	4%	1	4%	1	3%
<b>Race/Ethnicity</b>							
White, not Hispanic	32.6	47	82%	21	78%	26	87%
Black, not Hispanic	190.0	10	18%	6	22%	4	13%
Hispanic	*	-	-	-	-	-	-
Asian/Pacific Islander	*	-	-	-	-	-	-
American Indian/Alaska Native	*	-	-	-	-	-	-
Unknown	*	-	-	-	-	-	-
<b>Race/Gender</b>							
White, not Hispanic Male	59.7	42	74%	18	67%	24	80%
White, not Hispanic Female	6.8	5	9%	3	11%	2	7%
Black, not Hispanic Male	277.2	7	12%	3	11%	4	13%
Black, not Hispanic Female	*	3	5%	3	11%	-	-
Other Male	*	-	-	-	-	-	-
Other Female	*	-	-	-	-	-	-
Unknown	*	-	-	-	-	-	-
<b>Total</b>	<b>37.5</b>	<b>57</b>		<b>27</b>		<b>30</b>	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (\*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> The rate is the number of persons living with HIV/AIDS per 100,000 population calculated using 2000 census data.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Portage County, Ohio

### Reported persons living with HIV/AIDS as of Dec. 31, 2003, by current disease status and transmission category

Transmission category	HIV/AIDS		Current Disease Status			
			HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	29	59%	9	43%	20	71%
Injection drug use (IDU)	1	2%	-	-	1	4%
Male-to-male sexual contact and IDU	-	-	-	-	-	-
Heterosexual contact	-	-	-	-	-	-
Blood product	-	-	-	-	-	-
Other/unknown	19	39%	12	57%	7	25%
Subtotal	49	100%	21	100%	28	100%
<b>Female adult or adolescent</b>						
Injection drug use	-	-	-	-	-	-
Heterosexual contact	2	25%	1	17%	1	50%
Blood product	1	13%	1	17%	-	-
Other/unknown	5	63%	4	67%	1	50%
Subtotal	8	100%	6	100%	2	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	-	-	-	-	-	-
Other	-	-	-	-	-	-
Subtotal	-	-	-	-	-	-
<b>Total</b>	<b>57</b>		<b>27</b>		<b>30</b>	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (\*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Portage County, Ohio

### Reported cases of HIV infection, by year of diagnosis and selected characteristics

	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
<b>Gender</b>				
Male	9	90%	12	80%
Female	1	10%	3	20%
Unknown	-	-	-	-
<b>Age</b>				
<13	-	-	-	-
13-19	-	-	-	-
20-24	1	10%	3	20%
25-29	-	-	-	-
30-39	5	50%	6	40%
40-49	3	30%	3	20%
50-64	1	10%	2	13%
65+	-	-	1	7%
<b>Race/Ethnicity</b>				
White, not Hispanic	9	90%	11	73%
Black, not Hispanic	1	10%	4	27%
Hispanic	-	-	-	-
Asian/Pacific Islander	-	-	-	-
American Indian/Alaska Native	-	-	-	-
Unknown	-	-	-	-
<b>Race/Gender</b>				
White, not Hispanic Male	8	80%	9	60%
White, not Hispanic Female	1	10%	2	13%
Black, not Hispanic Male	1	10%	3	20%
Black, not Hispanic Female	-	-	1	7%
Other Male	-	-	-	-
Other Female	-	-	-	-
Unknown	-	-	-	-
<b>Total</b>	<b>10</b>		<b>15</b>	

*Note:* HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Portage County, Ohio

### Reported cases of HIV infection, by year of diagnosis and transmission category

Transmission category	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
<b>Male adult or adolescent</b>				
Male-to-male sexual contact	4	44%	3	25%
Injection drug use (IDU)	-	-	-	-
Male-to-male sexual contact and IDU	-	-	1	8%
Heterosexual contact	-	-	-	-
Blood product	-	-	-	-
Other/unknown	5	56%	8	67%
Subtotal	9	100%	12	100%
<b>Female adult or adolescent</b>				
Injection drug use	-	-	-	-
Heterosexual contact	1	100%	-	-
Blood product	-	-	-	-
Other/unknown	-	-	3	100%
Subtotal	1	100%	3	100%
<b>Child (&lt;13 yrs at diagnosis)</b>				
Perinatal	-	-	-	-
Other	-	-	-	-
Subtotal	-	-	-	-
<b>Total</b>	<b>10</b>		<b>15</b>	

*Note:* HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Portage County, Ohio

### Cumulative cases of AIDS diagnosed through 2003, by selected characteristics and transmission category

Cumulative AIDS Diagnoses					
	No.	%		No.	%
<b>Gender</b>			<b>Transmission category</b>		
Male	52	90%	<b>Male adult or adolescent</b>		
Female	6	10%	Male-to-male sexual contact	37	71%
Unknown	-	-	Injection drug use (IDU)	1	2%
			Male-to-male sexual contact and IDU	4	8%
<b>Age</b>			Heterosexual contact	1	2%
<13	-	-	Blood product	1	2%
13-19	-	-	Other/unknown	8	15%
20-24	4	7%	Subtotal	52	100%
25-29	7	12%	<b>Female adult or adolescent</b>		
30-39	27	47%	Injection drug use	2	33%
40-49	15	26%	Heterosexual contact	3	50%
50-64	5	9%	Blood product	-	-
65+	-	-	Other/unknown	1	17%
			Subtotal	6	100%
<b>Race/Ethnicity</b>			<b>Child (&lt;13 yrs at diagnosis)</b>		
White, not Hispanic	50	86%	Perinatal	-	-
Black, not Hispanic	8	14%	Other	-	-
Hispanic	-	-	Subtotal	-	-
Asian/Pacific Islander	-	-			
American Indian/Alaska Native	-	-			
Unknown	-	-			
<b>Race/Gender</b>					
White, not Hispanic Male	46	79%			
White, not Hispanic Female	4	7%			
Black, not Hispanic Male	6	10%			
Black, not Hispanic Female	2	3%			
Other Male	-	-			
Other Female	-	-			
Unknown	-	-			
<b>Total</b>	<b>58</b>				

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.