

Preble County, Ohio

Reported persons living with HIV/AIDS as of Dec. 31, 2003, by selected characteristics and transmission category

Reported Persons Living with HIV/AIDS					
	No.	%		No.	%
Current disease status			Race/Gender		
HIV (not AIDS)	3	19%	White, not Hispanic Male	11	69%
AIDS	13	81%	White, not Hispanic Female	3	19%
			Black, not Hispanic Male	1	6%
Gender			Black, not Hispanic Female	-	-
Male	12	75%	Other Male	-	-
Female	4	25%	Other Female	1	6%
Unknown	-	-	Unknown	-	-
Age			Transmission category		
<13	-	-	Male adult or adolescent		
13-19	-	-	Male-to-male sexual contact	5	42%
20-24	-	-	Injection drug use (IDU)	1	8%
25-29	-	-	Male-to-male sexual contact and IDU	1	8%
30-39	4	25%			-
40-49	9	56%	Heterosexual contact	1	8%
50-64	2	13%	Blood product	1	8%
65+	1	6%	Other/unknown	3	25%
			Subtotal	12	100%
Race/Ethnicity			Female adult or adolescent		
White, not Hispanic	14	88%	Injection drug use	2	50%
Black, not Hispanic	1	6%	Heterosexual contact	-	-
Hispanic	-	-	Blood product	1	25%
Asian/Pacific Islander	1	6%	Other/unknown	1	25%
American Indian/Alaska Native	-	-	Subtotal	4	100%
Unknown	-	-	Child (<13 yrs at diagnosis)		
			Perinatal	-	-
			Other	-	-
			Subtotal	-	-
Total	16				

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Preble County, Ohio

Cumulative cases of AIDS diagnosed through 2003, by selected characteristics and transmission category

	Cumulative AIDS Diagnoses			
	No.	%	No.	%
Gender			Transmission category	
Male	14	82%	Male adult or adolescent	
Female	3	18%	Male-to-male sexual contact	6 43%
Unknown	-	-	Injection drug use (IDU)	- -
			Male-to-male sexual contact and IDU	2 14%
Age			Heterosexual contact	- -
<13	-	-	Blood product	4 29%
13-19	1	6%	Other/unknown	2 14%
20-24	1	6%	Subtotal	14 100%
25-29	2	12%	Female adult or adolescent	
30-39	4	24%	Injection drug use	1 33%
40-49	8	47%	Heterosexual contact	- -
50-64	-	-	Blood product	- -
65+	1	6%	Other/unknown	2 67%
			Subtotal	3 100%
Race/Ethnicity			Child (<13 yrs at diagnosis)	
White, not Hispanic	16	94%	Perinatal	- -
Black, not Hispanic	-	-	Other	- -
Hispanic	-	-	Subtotal	- -
Asian/Pacific Islander	1	6%		
American Indian/Alaska Native	-	-		
Unknown	-	-		
Race/Gender				
White, not Hispanic Male	14	82%		
White, not Hispanic Female	2	12%		
Black, not Hispanic Male	-	-		
Black, not Hispanic Female	-	-		
Other Male	-	-		
Other Female	1	6%		
Unknown	-	-		
Total	17			

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.