

Seneca County, Ohio

Reported persons living with HIV/AIDS as of Dec. 31, 2003, by selected characteristics and transmission category

Reported Persons Living with HIV/AIDS					
	No.	%		No.	%
Current disease status			Race/Gender		
HIV (not AIDS)	4	25%	White, not Hispanic Male	13	81%
AIDS	12	75%	White, not Hispanic Female	1	6%
			Black, not Hispanic Male	-	-
Gender			Black, not Hispanic Female	-	-
Male	15	94%	Other Male	2	13%
Female	1	6%	Other Female	-	-
Unknown	-	-	Unknown	-	-
Age			Transmission category		
<13	-	-	Male adult or adolescent		
13-19	-	-	Male-to-male sexual contact	10	67%
20-24	-	-	Injection drug use (IDU)	2	13%
25-29	1	6%	Male-to-male sexual contact and IDU	2	13%
30-39	6	38%			-
40-49	7	44%	Heterosexual contact	-	-
50-64	1	6%	Blood product	-	-
65+	1	6%	Other/unknown	1	7%
			Subtotal	15	100%
Race/Ethnicity			Female adult or adolescent		
White, not Hispanic	14	88%	Injection drug use	-	-
Black, not Hispanic	-	-	Heterosexual contact	1	100%
Hispanic	1	6%	Blood product	-	-
Asian/Pacific Islander	-	-	Other/unknown	-	-
American Indian/Alaska Native	1	6%	Subtotal	1	100%
Unknown	-	-	Child (<13 yrs at diagnosis)		
			Perinatal	-	-
			Other	-	-
			Subtotal	-	-
Total	16				

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Seneca County, Ohio

Cumulative cases of AIDS diagnosed through 2003, by selected characteristics and transmission category

		Cumulative AIDS Diagnoses				
		No.	%		No.	%
Gender				Transmission category		
Male	25	100%		Male adult or adolescent		
Female	-	-		Male-to-male sexual contact	17	68%
Unknown	-	-		Injection drug use (IDU)	4	16%
				Male-to-male sexual contact and IDU	2	8%
Age				Heterosexual contact	-	-
<13	-	-		Blood product	1	4%
13-19	-	-		Other/unknown	1	4%
20-24	1	4%		Subtotal	25	100%
25-29	6	24%		Female adult or adolescent		
30-39	11	44%		Injection drug use	-	-
40-49	6	24%		Heterosexual contact	-	-
50-64	-	-		Blood product	-	-
65+	1	4%		Other/unknown	-	-
				Subtotal	-	-
Race/Ethnicity				Child (<13 yrs at diagnosis)		
White, not Hispanic	23	92%		Perinatal	-	-
Black, not Hispanic	-	-		Other	-	-
Hispanic	1	4%		Subtotal	-	-
Asian/Pacific Islander	-	-				
American Indian/Alaska Native	1	4%				
Unknown	-	-				
Race/Gender						
White, not Hispanic Male	23	92%				
White, not Hispanic Female	-	-				
Black, not Hispanic Male	-	-				
Black, not Hispanic Female	-	-				
Other Male	2	8%				
Other Female	-	-				
Unknown	-	-				
Total	25					

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.