

## Shelby County, Ohio

### Reported persons living with HIV/AIDS as of Dec. 31, 2003, by selected characteristics and transmission category

Reported Persons Living with HIV/AIDS						
			No.	%		
			No.	%	No.	%
<b>Current disease status</b>					<b>Race/Gender</b>	
HIV (not AIDS)	7	44%	White, not Hispanic Male	11	69%	
AIDS	9	56%	White, not Hispanic Female	4	25%	
			Black, not Hispanic Male	-	-	
<b>Gender</b>			Black, not Hispanic Female	1	6%	
Male	11	69%	Other Male	-	-	
Female	5	31%	Other Female	-	-	
Unknown	-	-	Unknown	-	-	
<b>Age</b>			<b>Transmission category</b>			
<13	-	-	<b>Male adult or adolescent</b>			
13-19	-	-	Male-to-male sexual contact	6	55%	
20-24	1	6%	Injection drug use (IDU)	-	-	
25-29	2	13%	Male-to-male sexual contact and IDU	-	-	
30-39	7	44%	Heterosexual contact	1	9%	
40-49	4	25%	Blood product	1	9%	
50-64	2	13%	Other/unknown	3	27%	
65+	-	-	Subtotal	11	100%	
<b>Race/Ethnicity</b>			<b>Female adult or adolescent</b>			
White, not Hispanic	15	94%	Injection drug use	-	-	
Black, not Hispanic	1	6%	Heterosexual contact	1	20%	
Hispanic	-	-	Blood product	-	-	
Asian/Pacific Islander	-	-	Other/unknown	4	80%	
American Indian/Alaska Native	-	-	Subtotal	5	100%	
Unknown	-	-	<b>Child (&lt;13 yrs at diagnosis)</b>			
			Perinatal	-	-	
			Other	-	-	
			Subtotal	-	-	
<b>Total</b>	<b>16</b>					

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Shelby County, Ohio

### Cumulative cases of AIDS diagnosed through 2003, by selected characteristics and transmission category

		Cumulative AIDS Diagnoses			
		No.	%		
		No.	%	No.	%
<b>Gender</b>				<b>Transmission category</b>	
Male	22	92%	<b>Male adult or adolescent</b>		
Female	2	8%	Male-to-male sexual contact	15	68%
Unknown	-	-	Injection drug use (IDU)	-	-
			Male-to-male sexual contact and IDU	-	-
<b>Age</b>			Heterosexual contact	1	5%
<13	-	-	Blood product	3	14%
13-19	1	4%	Other/unknown	3	14%
20-24	2	8%	Subtotal	22	100%
25-29	3	13%	<b>Female adult or adolescent</b>		
30-39	12	50%	Injection drug use	-	-
40-49	2	8%	Heterosexual contact	2	100%
50-64	3	13%	Blood product	-	-
65+	1	4%	Other/unknown	-	-
			Subtotal	2	100%
<b>Race/Ethnicity</b>			<b>Child (&lt;13 yrs at diagnosis)</b>		
White, not Hispanic	21	88%	Perinatal	-	-
Black, not Hispanic	3	13%	Other	-	-
Hispanic	-	-	Subtotal	-	-
Asian/Pacific Islander	-	-			
American Indian/Alaska Native	-	-			
Unknown	-	-			
<b>Race/Gender</b>					
White, not Hispanic Male	20	83%			
White, not Hispanic Female	1	4%			
Black, not Hispanic Male	2	8%			
Black, not Hispanic Female	1	4%			
Other Male	-	-			
Other Female	-	-			
Unknown	-	-			
<b>Total</b>	<b>24</b>				

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.