

Tuscarawas County, Ohio

Reported persons living with HIV/AIDS as of Dec. 31, 2003, by selected characteristics and transmission category

Reported Persons Living with HIV/AIDS					
	No.	%		No.	%
Current disease status			Race/Gender		
HIV (not AIDS)	9	56%	White, not Hispanic Male	13	81%
AIDS	7	44%	White, not Hispanic Female	2	13%
			Black, not Hispanic Male	1	6%
Gender			Black, not Hispanic Female	-	-
Male	14	88%	Other Male	-	-
Female	2	13%	Other Female	-	-
Unknown	-	-	Unknown	-	-
Age			Transmission category		
<13	-	-	Male adult or adolescent		
13-19	-	-	Male-to-male sexual contact	7	50%
20-24	-	-	Injection drug use (IDU)	-	-
25-29	-	-	Male-to-male sexual contact and IDU	2	14%
30-39	6	38%	Heterosexual contact	-	-
40-49	7	44%	Blood product	-	-
50-64	3	19%	Other/unknown	5	36%
65+	-	-	Subtotal	14	100%
Race/Ethnicity			Female adult or adolescent		
White, not Hispanic	15	94%	Injection drug use	-	-
Black, not Hispanic	1	6%	Heterosexual contact	1	50%
Hispanic	-	-	Blood product	-	-
Asian/Pacific Islander	-	-	Other/unknown	1	50%
American Indian/Alaska Native	-	-	Subtotal	2	100%
Unknown	-	-	Child (<13 yrs at diagnosis)		
			Perinatal	-	-
			Other	-	-
			Subtotal	-	-
Total	16				

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

Tuscarawas County, Ohio

Cumulative cases of AIDS diagnosed through 2003, by selected characteristics and transmission category

		Cumulative AIDS Diagnoses			
		No.	%		
		No.	%	No.	%
Gender				Transmission category	
Male	13	76%		Male adult or adolescent	
Female	4	24%		Male-to-male sexual contact	10 77%
Unknown	-	-		Injection drug use (IDU)	- -
				Male-to-male sexual contact and IDU	1 8%
Age				Heterosexual contact	- -
<13	-	-		Blood product	1 8%
13-19	-	-		Other/unknown	1 8%
20-24	-	-		Subtotal	13 100%
25-29	3	18%		Female adult or adolescent	
30-39	6	35%		Injection drug use	- -
40-49	4	24%		Heterosexual contact	3 75%
50-64	3	18%		Blood product	- -
65+	1	6%		Other/unknown	1 25%
				Subtotal	4 100%
Race/Ethnicity				Child (<13 yrs at diagnosis)	
White, not Hispanic	15	88%		Perinatal	- -
Black, not Hispanic	2	12%		Other	- -
Hispanic	-	-		Subtotal	- -
Asian/Pacific Islander	-	-			
American Indian/Alaska Native	-	-			
Unknown	-	-			
Race/Gender					
White, not Hispanic Male	12	71%			
White, not Hispanic Female	3	18%			
Black, not Hispanic Male	1	6%			
Black, not Hispanic Female	1	6%			
Other Male	-	-			
Other Female	-	-			
Unknown	-	-			
Total	17				

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.