

## Union County, Ohio

### Reported persons living with HIV/AIDS as of Dec. 31, 2003, by selected characteristics and transmission category

Reported Persons Living with HIV/AIDS						
			No.	%		
			No.	%	No.	%
<b>Current disease status</b>					<b>Race/Gender</b>	
HIV (not AIDS)	6	43%	White, not Hispanic Male	10	71%	
AIDS	8	57%	White, not Hispanic Female	2	14%	
			Black, not Hispanic Male	-	-	
<b>Gender</b>			Black, not Hispanic Female	-	-	
Male	11	79%	Other Male	-	-	
Female	3	21%	Other Female	1	7%	
Unknown	-	-	Unknown	1	7%	
<b>Age</b>			<b>Transmission category</b>			
<13	-	-	<b>Male adult or adolescent</b>			
13-19	-	-	Male-to-male sexual contact	6	55%	
20-24	-	-	Injection drug use (IDU)	-	-	
25-29	2	14%	Male-to-male sexual contact and IDU	1	9%	
30-39	3	21%	Heterosexual contact	-	-	
40-49	7	50%	Blood product	-	-	
50-64	2	14%	Other/unknown	4	36%	
65+	-	-	Subtotal	11	100%	
<b>Race/Ethnicity</b>			<b>Female adult or adolescent</b>			
White, not Hispanic	12	86%	Injection drug use	-	-	
Black, not Hispanic	-	-	Heterosexual contact	2	67%	
Hispanic	1	7%	Blood product	-	-	
Asian/Pacific Islander	-	-	Other/unknown	1	33%	
American Indian/Alaska Native	-	-	Subtotal	3	100%	
Unknown	1	7%	<b>Child (&lt;13 yrs at diagnosis)</b>			
			Perinatal	-	-	
			Other	-	-	
			Subtotal	-	-	
<b>Total</b>	<b>14</b>					

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Union County, Ohio

### Cumulative cases of AIDS diagnosed through 2003, by selected characteristics and transmission category

	Cumulative AIDS Diagnoses			
	No.	%	No.	%
<b>Gender</b>			<b>Transmission category</b>	
Male	14	82%	<b>Male adult or adolescent</b>	
Female	3	18%	Male-to-male sexual contact	9 69%
Unknown	-	-	Injection drug use (IDU)	2 15%
			Male-to-male sexual contact and IDU	- -
<b>Age</b>			Heterosexual contact	- -
<13	-	-	Blood product	- -
13-19	1	6%	Other/unknown	2 15%
20-24	1	6%	Subtotal	13 100%
25-29	2	12%	<b>Female adult or adolescent</b>	
30-39	9	53%	Injection drug use	1 33%
40-49	1	6%	Heterosexual contact	2 67%
50-64	3	18%	Blood product	- -
65+	-	-	Other/unknown	- -
			Subtotal	3 100%
<b>Race/Ethnicity</b>			<b>Child (&lt;13 yrs at diagnosis)</b>	
White, not Hispanic	16	94%	Perinatal	- -
Black, not Hispanic	1	6%	Other	1 100%
Hispanic	-	-	Subtotal	1 100%
Asian/Pacific Islander	-	-		
American Indian/Alaska Native	-	-		
Unknown	-	-		
<b>Race/Gender</b>				
White, not Hispanic Male	14	82%		
White, not Hispanic Female	2	12%		
Black, not Hispanic Male	-	-		
Black, not Hispanic Female	1	6%		
Other Male	-	-		
Other Female	-	-		
Unknown	-	-		
<b>Total</b>	<b>17</b>			

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.