

HIV/AIDS Data by HIV Prevention Regional Advisory Groups (RAGs)

The following tables present HIV/AIDS data for the nine HIV Prevention RAGs. For a listing and a map depiction of the counties associated with each of these regions, please refer to section 3 starting on page 235. Additional data sources (i.e. Census, STDs) are presented by RAGs in section 3.

There are six tables of data for each of the regional advisory groups. The first table is reported persons living with HIV/AIDS as of Dec. 31, 2003, by current disease status and selected characteristics. The second table is reported persons living with HIV/AIDS by current disease status and transmission category. The third table is reported cases of HIV infection (including persons diagnosed with HIV and later progressed to AIDS and cases concurrently diagnosed with HIV and AIDS) by year of diagnosis and selected characteristics for time periods 1998 to 2000 and 2001 to 2003. The fourth table is reported cases of HIV infection by year of diagnosis and transmission category. The fifth table is reported cases of AIDS by year of diagnosis and selected characteristics for time periods 1998 to 2000 and 2001 to 2003 as well as cumulative AIDS diagnoses. The sixth table is AIDS diagnosis by year of diagnosis and transmission category.

When viewing these tables it is important to take into account the limitations of these data. These data represent confidential reports of HIV infection and AIDS diagnoses and not all persons with HIV infection. The HIV diagnosis data represent the earliest date of diagnosis reported to the HIV/AIDS Surveillance Program. They do not always reflect the earliest date the individual became aware of his or her HIV-infection status. The possibility exists that individuals might have been previously tested anonymously and not confidentially reported or might have been previously diagnosed out-of-state and not reported to Ohio until a later test was performed or diagnosis made.

HIV Prevention Akron Regional Advisory Group

Table 17. Reported persons living with HIV/AIDS as of Dec. 31, 2003, HIV Prevention Akron Regional Advisory Group, by current disease status and selected characteristics

	HIV/AIDS			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Gender							
Male	154.0	517	81%	232	77%	285	84%
Female	34.2	123	19%	68	23%	55	16%
Unknown	*	-	-	-	-	-	-
Age (as of 12/2003)							
<13	*	4	1%	3	1%	1	<1%
13-19	*	3	<1%	2	1%	1	<1%
20-24	26.3	12	2%	11	4%	1	<1%
25-29	91.3	40	6%	24	8%	16	5%
30-39	179.3	185	29%	93	31%	92	27%
40-49	227.0	249	39%	111	37%	138	41%
50-64	127.4	136	21%	53	18%	83	24%
65+	11.8	11	2%	3	1%	8	2%
Race/Ethnicity							
White, not Hispanic	62.0	371	58%	154	51%	217	64%
Black, not Hispanic	304.8	241	38%	124	41%	117	34%
Hispanic	204.3	12	2%	7	2%	5	1%
Asian/Pacific Islander	*	4	1%	3	1%	1	<1%
American Indian/Alaska Native	*	1	<1%	1	<1%	-	-
Unknown	*	11	2%	11	4%	-	-
Race/Gender							
White, not Hispanic Male	112.2	326	51%	134	45%	192	56%
White, not Hispanic Female	14.6	45	7%	20	7%	25	7%
Black, not Hispanic Male	455.1	167	26%	79	26%	88	26%
Black, not Hispanic Female	174.6	74	12%	45	15%	29	9%
Hispanic Male	335.5	10	2%	6	2%	4	1%
Hispanic Female	*	2	<1%	1	<1%	1	<1%
Asian/Pacific Islander Male	*	3	<1%	2	1%	1	<1%
Asian/Pacific Islander Female	*	1	<1%	1	<1%	-	-
American Indian/Alaska Native Male	*	1	<1%	1	<1%	-	-
American Indian/Alaska Native Female	*	-	-	-	-	-	-
Unknown	*	11	2%	11	4%	-	-
Total	92.1	640		300		340	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with HIV/AIDS per 100,000 population calculated using 2000 census data.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Akron Regional Advisory Group

Table 18. Reported persons living with HIV/AIDS as of Dec. 31, 2003, HIV Prevention Akron Regional Advisory Group, by current disease status and transmission category

Transmission category	HIV/AIDS		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	248	48%	68	30%	180	64%
Injection drug use (IDU)	29	6%	9	4%	20	7%
Male-to-male sexual contact and IDU	25	5%	7	3%	18	6%
Heterosexual contact	10	2%	4	2%	6	2%
Blood product	1	<1%	-	-	1	<1%
Other/unknown	200	39%	142	62%	58	20%
Subtotal	513	100%	230	100%	283	100%
Female adult or adolescent						
Injection drug use	13	11%	3	5%	10	18%
Heterosexual contact	40	33%	14	22%	26	47%
Blood product	1	1%	1	2%	-	-
Other/unknown	66	55%	47	72%	19	35%
Subtotal	120	100%	65	100%	55	100%
Child (<13 yrs at diagnosis)						
Perinatal	7	100%	5	100%	2	100%
Other	-	-	-	-	-	-
Subtotal	7	100%	5	100%	2	100%
Total	640		300		340	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Akron Regional Advisory Group

Table 19. Reported cases of HIV infection, HIV Prevention Akron Regional Advisory Group, by year of diagnosis and selected characteristics

	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
Gender				
Male	118	82%	116	77%
Female	26	18%	34	23%
Unknown	-	-	-	-
Age				
<13	2	1%	2	1%
13-19	3	2%	1	1%
20-24	11	8%	19	13%
25-29	13	9%	12	8%
30-39	58	40%	60	40%
40-49	43	30%	33	22%
50-64	13	9%	21	14%
65+	1	1%	2	1%
Race/Ethnicity				
White, not Hispanic	77	53%	83	55%
Black, not Hispanic	61	42%	55	37%
Hispanic	4	3%	4	3%
Asian/Pacific Islander	1	1%	3	2%
American Indian/Alaska Native	-	-	1	1%
Unknown	1	1%	4	3%
Race/Gender				
White, not Hispanic Male	68	47%	70	47%
White, not Hispanic Female	9	6%	13	9%
Black, not Hispanic Male	44	31%	36	24%
Black, not Hispanic Female	17	12%	19	13%
Hispanic Male	4	3%	3	2%
Hispanic Female	-	-	1	1%
Asian/Pacific Islander Male	1	1%	2	1%
Asian/Pacific Islander Female	-	-	1	1%
American Indian/Alaska Native Male	-	-	1	1%
American Indian/Alaska Native Female	-	-	-	-
Unknown	1	1%	4	3%
Total	144		150	

Note: HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Akron Regional Advisory Group

Table 20. Reported cases of HIV infection, HIV Prevention Akron Regional Advisory Group, by year of diagnosis and transmission category

Transmission category	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
Male adult or adolescent				
Male-to-male sexual contact	58	50%	32	28%
Injection drug use (IDU)	5	4%	4	3%
Male-to-male sexual contact and IDU	2	2%	1	1%
Heterosexual contact	1	1%	4	3%
Blood product	-	-	-	-
Other/unknown	51	44%	74	64%
Subtotal	117	100%	115	100%
Female adult or adolescent				
Injection drug use	-	-	2	6%
Heterosexual contact	10	40%	5	15%
Blood product	-	-	-	-
Other/unknown	15	60%	26	79%
Subtotal	25	100%	33	100%
Child (<13 yrs at diagnosis)				
Perinatal	2	100%	2	100%
Other	-	-	-	-
Subtotal	2	-	2	-
Total	144		150	

Note: HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Akron Regional Advisory Group

Table 21. Reported cases of AIDS, HIV Prevention Akron Regional Advisory Group, by year of diagnosis and selected characteristics

	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
Gender						
Male	87	84%	49	75%	570	88%
Female	17	16%	16	25%	75	12%
Unknown	-	-	-	-	-	-
Age						
<13	1	1%	-	-	1	<1%
13-19	1	1%	-	-	2	<1%
20-24	2	2%	3	5%	27	4%
25-29	13	13%	7	11%	91	14%
30-39	45	43%	26	40%	279	43%
40-49	34	33%	21	32%	182	28%
50-64	7	7%	7	11%	59	9%
65+	1	1%	1	2%	4	1%
Race/Ethnicity						
White, not Hispanic	63	61%	40	62%	414	64%
Black, not Hispanic	38	37%	24	37%	223	35%
Hispanic	2	2%	1	2%	7	1%
Asian/Pacific Islander	1	1%	-	-	1	<1%
American Indian/Alaska Native	-	-	-	-	-	-
Unknown	-	-	-	-	-	-
Race/Gender						
White, not Hispanic Male	54	52%	32	49%	381	59%
White, not Hispanic Female	9	9%	8	12%	33	5%
Black, not Hispanic Male	30	29%	16	25%	182	28%
Black, not Hispanic Female	8	8%	8	12%	41	6%
Hispanic Male	2	2%	1	2%	6	1%
Hispanic Female	-	-	-	-	1	<1%
Asian/Pacific Islander Male	1	1%	-	-	1	<1%
Asian/Pacific Islander Female	-	-	-	-	-	-
American Indian/Alaska Native Male	-	-	-	-	-	-
American Indian/Alaska Native Female	-	-	-	-	-	-
Unknown	-	-	-	-	-	-
Total	104		65		645	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Akron Regional Advisory Group

Table 22. Reported cases of AIDS, HIV Prevention Akron Regional Advisory Group, by year of diagnosis and transmission category

Transmission category	Year of AIDS Diagnosis				Cumulative	
	1998 - 2000		2001 - 2003			
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	62	72%	14	29%	405	71%
Injection drug use (IDU)	4	5%	2	4%	32	6%
Male-to-male sexual contact and IDU	2	2%	3	6%	38	7%
Heterosexual contact	-	-	1	2%	11	2%
Blood product	-	-	-	-	13	2%
Other/unknown	18	21%	29	59%	70	12%
Subtotal	86	100%	49	100%	569	100%
Female adult or adolescent						
Injection drug use	1	6%	1	6%	11	15%
Heterosexual contact	7	41%	7	44%	36	48%
Blood product	-	-	-	-	4	5%
Other/unknown	9	53%	8	50%	24	32%
Subtotal	17	100%	16	100%	75	100%
Child (<13 yrs at diagnosis)						
Perinatal	1	100%	-	-	1	100%
Other	-	-	-	-	-	-
Subtotal	1	100%	-	-	1	100%
Total	104		65		645	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Canton Regional Advisory Group

Table 23. Reported persons living with HIV/AIDS as of Dec. 31, 2003, HIV Prevention Canton Regional Advisory Group, by current disease status and selected characteristics

	HIV/AIDS			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Gender							
Male	106.2	208	77%	119	74%	89	82%
Female	28.9	61	23%	41	26%	20	18%
Unknown	*	-	-	-	-	-	-
Age (as of 12/2003)							
<13	*	3	1%	1	1%	2	2%
13-19	*	-	-	-	-	-	-
20-24	*	4	1%	4	3%	-	-
25-29	51.2	12	4%	9	6%	3	3%
30-39	164.7	93	35%	66	41%	27	25%
40-49	152.8	98	36%	49	31%	49	45%
50-64	83.5	56	21%	29	18%	27	25%
65+	*	3	1%	2	1%	1	1%
Race/Ethnicity							
White, not Hispanic	51.6	191	71%	107	67%	84	77%
Black, not Hispanic	229.4	67	25%	45	28%	22	20%
Hispanic	137.0	5	2%	2	1%	3	3%
Asian/Pacific Islander	*	-	-	-	-	-	-
American Indian/Alaska Native	*	-	-	-	-	-	-
Unknown	*	6	2%	6	4%	-	-
Race/Gender							
White, not Hispanic Male	87.4	156	58%	86	54%	70	64%
White, not Hispanic Female	18.2	35	13%	21	13%	14	13%
Black, not Hispanic Male	309.6	43	16%	27	17%	16	15%
Black, not Hispanic Female	156.6	24	9%	18	11%	6	6%
Hispanic Male	272.5	5	2%	2	1%	3	3%
Hispanic Female	*	-	-	-	-	-	-
Asian/Pacific Islander Male	*	-	-	-	-	-	-
Asian/Pacific Islander Female	*	-	-	-	-	-	-
American Indian/Alaska Native Male	*	-	-	-	-	-	-
American Indian/Alaska Native Female	*	-	-	-	-	-	-
Unknown	*	6	2%	6	4%	-	-
Total	66.1	269		160		109	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with HIV/AIDS per 100,000 population calculated using 2000 census data.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Canton Regional Advisory Group

Table 24. Reported persons living with HIV/AIDS as of Dec. 31, 2003, HIV Prevention Canton Regional Advisory Group, by current disease status and transmission category

Transmission category	Current Disease Status					
	HIV/AIDS		HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	113	54%	49	41%	64	72%
Injection drug use (IDU)	8	4%	4	3%	4	4%
Male-to-male sexual contact and IDU	6	3%	2	2%	4	4%
Heterosexual contact	7	3%	5	4%	2	2%
Blood product	3	1%	-	-	3	3%
Other/unknown	71	34%	59	50%	12	13%
Subtotal	208	100%	119	100%	89	100%
Female adult or adolescent						
Injection drug use	3	5%	3	8%	-	-
Heterosexual contact	21	36%	11	28%	10	56%
Blood product	1	2%	1	3%	-	-
Other/unknown	33	57%	25	63%	8	44%
Subtotal	58	100%	40	100%	18	100%
Child (<13 yrs at diagnosis)						
Perinatal	3	100%	1	100%	2	100%
Other	-	-	-	-	-	-
Subtotal	3	100%	1	100%	2	100%
Total	269		160		109	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Canton Regional Advisory Group

Table 25. Reported cases of HIV infection, HIV Prevention Canton Regional Advisory Group, by year of diagnosis and selected characteristics

	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
Gender				
Male	40	66%	35	78%
Female	21	34%	10	22%
Unknown	-	-	-	-
Age				
<13	2	3%	-	-
13-19	1	2%	1	2%
20-24	5	8%	2	4%
25-29	9	15%	4	9%
30-39	24	39%	21	47%
40-49	15	25%	8	18%
50-64	5	8%	9	20%
65+	-	-	-	-
Race/Ethnicity				
White, not Hispanic	47	77%	27	60%
Black, not Hispanic	12	20%	13	29%
Hispanic	1	2%	1	2%
Asian/Pacific Islander	-	-	-	-
American Indian/Alaska Native	-	-	-	-
Unknown	1	2%	4	9%
Race/Gender				
White, not Hispanic Male	31	51%	23	51%
White, not Hispanic Female	16	26%	4	9%
Black, not Hispanic Male	8	13%	8	18%
Black, not Hispanic Female	4	7%	5	11%
Hispanic Male	1	2%	1	2%
Hispanic Female	-	-	-	-
Asian/Pacific Islander Male	-	-	-	-
Asian/Pacific Islander Female	-	-	-	-
American Indian/Alaska Native Male	-	-	-	-
American Indian/Alaska Native Female	-	-	-	-
Unknown	1	2%	4	9%
Total	61		45	

Note: HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Canton Regional Advisory Group

Table 26. Reported cases of HIV infection, HIV Prevention Canton Regional Advisory Group, by year of diagnosis and transmission category

Transmission category	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
Male adult or adolescent				
Male-to-male sexual contact	26	65%	10	29%
Injection drug use (IDU)	1	3%	1	3%
Male-to-male sexual contact and IDU	-	-	-	-
Heterosexual contact	1	3%	1	3%
Blood product	-	-	-	-
Other/unknown	12	30%	23	66%
Subtotal	40	100%	35	100%
Female adult or adolescent				
Injection drug use	2	11%	-	-
Heterosexual contact	9	47%	2	20%
Blood product	-	-	-	-
Other/unknown	8	42%	8	80%
Subtotal	19	100%	10	100%
Child (<13 yrs at diagnosis)				
Perinatal	2	100%	-	-
Other	-	-	-	-
Subtotal	2	-	-	-
Total	61		45	

Note: HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Canton Regional Advisory Group

Table 27. Reported cases of AIDS, HIV Prevention Canton Regional Advisory Group, by year of diagnosis and selected characteristics

	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
Gender						
Male	27	82%	17	77%	210	86%
Female	6	18%	5	23%	33	14%
Unknown	-	-	-	-	-	-
Age						
<13	1	3%	-	-	6	2%
13-19	-	-	-	-	-	-
20-24	1	3%	-	-	13	5%
25-29	1	3%	-	-	29	12%
30-39	15	45%	12	55%	110	45%
40-49	9	27%	7	32%	59	24%
50-64	6	18%	2	9%	23	9%
65+	-	-	1	5%	3	1%
Race/Ethnicity						
White, not Hispanic	25	76%	16	73%	190	78%
Black, not Hispanic	7	21%	5	23%	48	20%
Hispanic	1	3%	1	5%	5	2%
Asian/Pacific Islander	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-
Unknown	-	-	-	-	-	-
Race/Gender						
White, not Hispanic Male	21	64%	12	55%	168	69%
White, not Hispanic Female	4	12%	4	18%	22	9%
Black, not Hispanic Male	5	15%	4	18%	37	15%
Black, not Hispanic Female	2	6%	1	5%	11	5%
Hispanic Male	1	3%	1	5%	5	2%
Hispanic Female	-	-	-	-	-	-
Asian/Pacific Islander Male	-	-	-	-	-	-
Asian/Pacific Islander Female	-	-	-	-	-	-
American Indian/Alaska Native Male	-	-	-	-	-	-
American Indian/Alaska Native Female	-	-	-	-	-	-
Unknown	-	-	-	-	-	-
Total	33		22		243	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Canton Regional Advisory Group

Table 28. Reported cases of AIDS, HIV Prevention Canton Regional Advisory Group, by year of diagnosis and transmission category

Transmission category	Year of AIDS Diagnosis				Cumulative	
	1998 - 2000		2001 - 2003			
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	21	78%	8	47%	152	73%
Injection drug use (IDU)	-	-	-	-	10	5%
Male-to-male sexual contact and IDU	-	-	-	-	7	3%
Heterosexual contact	-	-	-	-	5	2%
Blood product	-	-	-	-	9	4%
Other/unknown	6	22%	9	53%	25	12%
Subtotal	27	100%	17	100%	208	100%
Female adult or adolescent						
Injection drug use	1	20%	-	-	8	28%
Heterosexual contact	2	40%	2	40%	10	34%
Blood product	-	-	-	-	1	3%
Other/unknown	2	40%	3	60%	10	34%
Subtotal	5	100%	5	100%	29	100%
Child (<13 yrs at diagnosis)						
Perinatal	1	100%	-	-	6	100%
Other	-	-	-	-	-	-
Subtotal	1	100%	-	-	6	100%
Total	33		22		243	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Cincinnati Regional Advisory Group

Table 29. Reported persons living with HIV/AIDS as of Dec. 31, 2003, HIV Prevention Cincinnati Regional Advisory Group, by current disease status and selected characteristics

	HIV/AIDS			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Gender							
Male	258.0	1,684	82%	880	78%	804	86%
Female	52.9	372	18%	246	22%	126	14%
Unknown	*	-	-	-	-	-	-
Age (as of 12/2003)							
<13	4.7	12	1%	9	1%	3	<1%
13-19	4.2	6	<1%	3	<1%	3	<1%
20-24	62.7	59	3%	52	5%	7	1%
25-29	148.5	135	7%	99	9%	36	4%
30-39	319.1	652	32%	383	34%	269	29%
40-49	394.7	828	40%	415	37%	413	44%
50-64	168.4	329	16%	150	13%	179	19%
65+	21.1	35	2%	15	1%	20	2%
Race/Ethnicity							
White, not Hispanic	91.1	996	48%	527	47%	469	50%
Black, not Hispanic	434.8	964	47%	532	47%	432	46%
Hispanic	195.8	31	2%	15	1%	16	2%
Asian/Pacific Islander	41.4	9	<1%	3	<1%	6	1%
American Indian/Alaska Native	*	4	<1%	2	<1%	2	<1%
Unknown	*	52	3%	47	4%	5	1%
Race/Gender							
White, not Hispanic Male	166.6	883	43%	451	40%	432	46%
White, not Hispanic Female	20.0	113	5%	76	7%	37	4%
Black, not Hispanic Male	708.3	721	35%	374	33%	347	37%
Black, not Hispanic Female	202.7	243	12%	158	14%	85	9%
Hispanic Male	294.6	25	1%	11	1%	14	2%
Hispanic Female	81.7	6	<1%	4	<1%	2	<1%
Asian/Pacific Islander Male	74.0	8	<1%	3	<1%	5	1%
Asian/Pacific Islander Female	*	1	<1%	-	-	1	<1%
American Indian/Alaska Native Male	*	4	<1%	2	<1%	2	<1%
American Indian/Alaska Native Female	*	-	-	-	-	-	-
Unknown	*	52	3%	47	4%	5	1%
Total	151.6	2,056		1,126		930	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with HIV/AIDS per 100,000 population calculated using 2000 census data.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Cincinnati Regional Advisory Group

Table 30. Reported persons living with HIV/AIDS as of Dec. 31, 2003, HIV Prevention Cincinnati Regional Advisory Group, by current disease status and transmission category

Transmission category	HIV/AIDS		Current Disease Status			
			HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	895	54%	381	44%	514	64%
Injection drug use (IDU)	101	6%	38	4%	63	8%
Male-to-male sexual contact and IDU	78	5%	27	3%	51	6%
Heterosexual contact	40	2%	15	2%	25	3%
Blood product	14	1%	7	1%	7	1%
Other/unknown	544	33%	406	46%	138	17%
Subtotal	1,672	100%	874	100%	798	100%
Female adult or adolescent						
Injection drug use	40	11%	18	7%	22	18%
Heterosexual contact	122	33%	70	29%	52	42%
Blood product	3	1%	2	1%	1	1%
Other/unknown	200	55%	151	63%	49	40%
Subtotal	365	100%	241	100%	124	100%
Child (<13 yrs at diagnosis)						
Perinatal	13	68%	9	82%	4	50%
Other	6	32%	2	18%	4	50%
Subtotal	19	100%	11	100%	8	100%
Total	2,056		1,126		930	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Cincinnati Regional Advisory Group

Table 31. Reported cases of HIV infection, HIV Prevention Cincinnati Regional Advisory Group, by year of diagnosis and selected characteristics

	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
Gender				
Male	315	80%	310	77%
Female	79	20%	91	23%
Unknown	-	-	-	-
Age				
<13	3	1%	-	-
13-19	14	4%	13	3%
20-24	39	10%	45	11%
25-29	63	16%	52	13%
30-39	162	41%	132	33%
40-49	78	20%	117	29%
50-64	31	8%	35	9%
65+	4	1%	7	2%
Race/Ethnicity				
White, not Hispanic	179	45%	165	41%
Black, not Hispanic	189	48%	205	51%
Hispanic	7	2%	9	2%
Asian/Pacific Islander	1	<1%	-	-
American Indian/Alaska Native	-	-	1	<1%
Unknown	18	5%	21	5%
Race/Gender				
White, not Hispanic Male	155	39%	137	34%
White, not Hispanic Female	24	6%	28	7%
Black, not Hispanic Male	139	35%	150	37%
Black, not Hispanic Female	50	13%	55	14%
Hispanic Male	5	1%	6	1%
Hispanic Female	2	1%	3	1%
Asian/Pacific Islander Male	1	<1%	-	-
Asian/Pacific Islander Female	-	-	-	-
American Indian/Alaska Native Male	-	-	1	<1%
American Indian/Alaska Native Female	-	-	-	-
Unknown	18	5%	21	5%
Total	394		401	

Note: HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Cincinnati Regional Advisory Group

Table 32. Reported cases of HIV infection, HIV Prevention Cincinnati Regional Advisory Group, by year of diagnosis and transmission category

Transmission category	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
Male adult or adolescent				
Male-to-male sexual contact	138	44%	112	36%
Injection drug use (IDU)	13	4%	14	5%
Male-to-male sexual contact and IDU	5	2%	4	1%
Heterosexual contact	3	1%	13	4%
Blood product	1	<1%	2	1%
Other/unknown	154	49%	165	53%
Subtotal	314	100%	310	100%
Female adult or adolescent				
Injection drug use	3	4%	3	3%
Heterosexual contact	24	31%	19	21%
Blood product	-	-	-	-
Other/unknown	50	65%	69	76%
Subtotal	77	100%	91	100%
Child (<13 yrs at diagnosis)				
Perinatal	1	33%	-	-
Other	2	67%	-	-
Subtotal	3	-	-	-
Total	394		401	

Note: HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Cincinnati Regional Advisory Group

Table 33. Reported cases of AIDS, HIV Prevention Cincinnati Regional Advisory Group, by year of diagnosis and selected characteristics

	Year of AIDS Diagnosis				Cumulative	
	1998 - 2000		2001 - 2003			
	No.	%	No.	%	No.	%
Gender						
Male	181	87%	158	84%	1,746	91%
Female	26	13%	31	16%	181	9%
Unknown	-	-	-	-	-	-
Age						
<13	1	<1%	-	-	14	1%
13-19	1	<1%	3	2%	12	1%
20-24	6	3%	9	5%	65	3%
25-29	32	15%	16	8%	300	16%
30-39	97	47%	74	39%	907	47%
40-49	51	25%	62	33%	455	24%
50-64	17	8%	20	11%	153	8%
65+	2	1%	5	3%	21	1%
Race/Ethnicity						
White, not Hispanic	95	46%	75	40%	1,122	58%
Black, not Hispanic	106	51%	106	56%	773	40%
Hispanic	2	1%	4	2%	16	1%
Asian/Pacific Islander	2	1%	1	1%	8	<1%
American Indian/Alaska Native	-	-	-	-	2	<1%
Unknown	2	1%	3	2%	6	<1%
Race/Gender						
White, not Hispanic Male	90	43%	63	33%	1,064	55%
White, not Hispanic Female	5	2%	12	6%	58	3%
Black, not Hispanic Male	86	42%	88	47%	654	34%
Black, not Hispanic Female	20	10%	18	10%	119	6%
Hispanic Male	2	1%	4	2%	14	1%
Hispanic Female	-	-	-	-	2	<1%
Asian/Pacific Islander Male	1	<1%	1	1%	7	<1%
Asian/Pacific Islander Female	1	<1%	-	-	1	<1%
American Indian/Alaska Native Male	-	-	-	-	2	<1%
American Indian/Alaska Native Female	-	-	-	-	-	-
Unknown	2	1%	3	2%	6	<1%
Total	207		189		1,927	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Cincinnati Regional Advisory Group

Table 34. Reported cases of AIDS, HIV Prevention Cincinnati Regional Advisory Group, by year of diagnosis and transmission category

Transmission category	Year of AIDS Diagnosis				Cumulative	
	1998 - 2000		2001 - 2003			
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	110	61%	73	46%	1,254	72%
Injection drug use (IDU)	10	6%	13	8%	94	5%
Male-to-male sexual contact and IDU	7	4%	6	4%	115	7%
Heterosexual contact	1	1%	5	3%	34	2%
Blood product	1	1%	-	-	37	2%
Other/unknown	52	29%	60	38%	200	12%
Subtotal	181	100%	157	100%	1,734	100%
Female adult or adolescent						
Injection drug use	3	12%	2	6%	39	22%
Heterosexual contact	7	28%	8	26%	74	42%
Blood product	-	-	-	-	2	1%
Other/unknown	15	60%	21	68%	61	35%
Subtotal	25	100%	31	100%	176	100%
Child (<13 yrs at diagnosis)						
Perinatal	-	-	-	-	11	65%
Other	1	100%	1	100%	6	35%
Subtotal	1	100%	1	100%	17	100%
Total	207		189		1,927	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Cleveland Regional Advisory Group

Table 35. Reported persons living with HIV/AIDS as of Dec. 31, 2003, HIV Prevention Cleveland Regional Advisory Group, by current disease status and selected characteristic

	Current Disease Status						
	HIV/AIDS			HIV (not AIDS)		AIDS	
	Rate ^a	No.	%	No.	%	No.	%
Gender							
Male	379.8	2,501	77%	1,094	74%	1,407	80%
Female	100.3	738	23%	385	26%	353	20%
Unknown	*	1	<1%	1	<1%	-	-
Age (as of 12/2003)							
<13	10.3	26	1%	14	1%	12	1%
13-19	20.2	26	1%	16	1%	10	1%
20-24	83.9	65	2%	45	3%	20	1%
25-29	151.3	138	4%	90	6%	48	3%
30-39	486.3	995	31%	515	35%	480	27%
40-49	631.9	1,349	42%	546	37%	803	46%
50-64	279.7	583	18%	232	16%	351	20%
65+	26.7	58	2%	22	1%	36	2%
Race/Ethnicity							
White, not Hispanic	116.2	1,081	33%	460	31%	621	35%
Black, not Hispanic	455.1	1,758	54%	801	54%	957	54%
Hispanic	622.4	293	9%	122	8%	171	10%
Asian/Pacific Islander	36.5	10	<1%	5	<1%	5	<1%
American Indian/Alaska Native	299.2	9	<1%	4	<1%	5	<1%
Unknown	*	89	3%	88	6%	1	<1%
Race/Gender							
White, not Hispanic Male	213.0	948	29%	389	26%	559	32%
White, not Hispanic Female	27.4	133	4%	71	5%	62	4%
Black, not Hispanic Male	730.0	1,277	39%	559	38%	718	41%
Black, not Hispanic Female	227.1	480	15%	241	16%	239	14%
Hispanic Male	822.1	193	6%	74	5%	119	7%
Hispanic Female	423.7	100	3%	48	3%	52	3%
Asian/Pacific Islander Male	59.1	8	<1%	3	<1%	5	<1%
Asian/Pacific Islander Female	*	2	<1%	2	<1%	-	-
American Indian/Alaska Native Male	491.6	7	<1%	2	<1%	5	<1%
American Indian/Alaska Native Female	*	2	<1%	2	<1%	-	-
Unknown	*	90	3%	89	6%	1	<1%
Total	232.4	3,240		1,480		1,760	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with HIV/AIDS per 100,000 population calculated using 2000 census data.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Cleveland Regional Advisory Group

Table 36. Reported persons living with HIV/AIDS as of Dec. 31, 2003, HIV Prevention Cleveland Regional Advisory Group, by current disease status and transmission category

Transmission category	HIV/AIDS		Current Disease Status			
			HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	1,388	56%	515	48%	873	62%
Injection drug use (IDU)	249	10%	80	7%	169	12%
Male-to-male sexual contact and IDU	148	6%	50	5%	98	7%
Heterosexual contact	101	4%	39	4%	62	4%
Blood product	13	1%	5	<1%	8	1%
Other/unknown	581	23%	393	36%	188	13%
Subtotal	2,480	100%	1,082	100%	1,398	100%
Female adult or adolescent						
Injection drug use	143	20%	49	13%	94	28%
Heterosexual contact	268	38%	118	32%	150	44%
Blood product	5	1%	2	1%	3	1%
Other/unknown	295	41%	203	55%	92	27%
Subtotal	711	100%	372	100%	339	100%
Child (<13 yrs at diagnosis)						
Perinatal	36	75%	18	72%	18	78%
Other	12	25%	7	28%	5	22%
Subtotal	48	100%	25	100%	23	100%
Total	3,240		1,480		1,760	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Cleveland Regional Advisory Group

Table 37. Reported cases of HIV infection, HIV Prevention Cleveland Regional Advisory Group, by year of diagnosis and selected characteristics

	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
Gender				
Male	473	76%	378	73%
Female	153	24%	139	27%
Unknown	-	-	-	-
Age				
<13	11	2%	6	1%
13-19	18	3%	7	1%
20-24	48	8%	48	9%
25-29	70	11%	67	13%
30-39	259	41%	191	37%
40-49	160	26%	145	28%
50-64	54	9%	52	10%
65+	6	1%	1	<1%
Race/Ethnicity				
White, not Hispanic	199	32%	153	30%
Black, not Hispanic	375	60%	300	58%
Hispanic	41	7%	41	8%
Asian/Pacific Islander	2	<1%	2	<1%
American Indian/Alaska Native	2	<1%	2	<1%
Unknown	7	1%	19	4%
Race/Gender				
White, not Hispanic Male	168	27%	137	26%
White, not Hispanic Female	31	5%	16	3%
Black, not Hispanic Male	263	42%	197	38%
Black, not Hispanic Female	112	18%	103	20%
Hispanic Male	32	5%	28	5%
Hispanic Female	9	1%	13	3%
Asian/Pacific Islander Male	2	<1%	-	-
Asian/Pacific Islander Female	-	-	2	<1%
American Indian/Alaska Native Male	2	<1%	1	<1%
American Indian/Alaska Native Female	-	-	1	<1%
Unknown	7	1%	19	4%
Total	626		517	

Note: HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Cleveland Regional Advisory Group

Table 38. Reported cases of HIV infection, HIV Prevention Cleveland Regional Advisory Group, by year of diagnosis and transmission category

Transmission category	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
Male adult or adolescent				
Male-to-male sexual contact	248	53%	165	44%
Injection drug use (IDU)	41	9%	18	5%
Male-to-male sexual contact and IDU	14	3%	6	2%
Heterosexual contact	20	4%	13	3%
Blood product	-	-	-	-
Other/unknown	143	31%	174	46%
Subtotal	466	100%	376	100%
Female adult or adolescent				
Injection drug use	13	9%	9	7%
Heterosexual contact	51	34%	26	19%
Blood product	2	1%	1	1%
Other/unknown	83	56%	99	73%
Subtotal	149	100%	135	100%
Child (<13 yrs at diagnosis)				
Perinatal	8	73%	5	83%
Other	3	27%	1	17%
Subtotal	11	-	6	-
Total	626		517	

Note: HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Cleveland Regional Advisory Group

Table 39. Reported cases of AIDS, HIV Prevention Cleveland Regional Advisory Group, by year of diagnosis and selected characteristics

	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
Gender						
Male	335	76%	308	73%	2,770	84%
Female	107	24%	112	27%	546	16%
Unknown	-	-	-	-	-	-
Age						
<13	2	<1%	4	1%	39	1%
13-19	2	<1%	3	1%	20	1%
20-24	15	3%	22	5%	132	4%
25-29	38	9%	34	8%	443	13%
30-39	202	46%	172	41%	1,522	46%
40-49	140	32%	133	32%	819	25%
50-64	38	9%	50	12%	303	9%
65+	5	1%	2	<1%	38	1%
Race/Ethnicity						
White, not Hispanic	128	29%	124	30%	1,281	39%
Black, not Hispanic	289	65%	256	61%	1,792	54%
Hispanic	21	5%	38	9%	231	7%
Asian/Pacific Islander	1	<1%	-	-	6	<1%
American Indian/Alaska Native	3	1%	1	<1%	5	<1%
Unknown	-	-	1	<1%	1	<1%
Race/Gender						
White, not Hispanic Male	112	25%	107	25%	1,193	36%
White, not Hispanic Female	16	4%	17	4%	88	3%
Black, not Hispanic Male	202	46%	176	42%	1,400	42%
Black, not Hispanic Female	87	20%	80	19%	392	12%
Hispanic Male	17	4%	23	5%	166	5%
Hispanic Female	4	1%	15	4%	65	2%
Asian/Pacific Islander Male	1	<1%	-	-	6	<1%
Asian/Pacific Islander Female	-	-	-	-	-	-
American Indian/Alaska Native Male	3	1%	1	<1%	4	<1%
American Indian/Alaska Native Female	-	-	-	-	1	<1%
Unknown	-	-	1	<1%	1	<1%
Total	442		420		3,316	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Cleveland Regional Advisory Group

Table 40. Reported cases of AIDS, HIV Prevention Cleveland Regional Advisory Group, by year and transmission category

Transmission category	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	194	58%	173	57%	1,845	67%
Injection drug use (IDU)	32	10%	25	8%	316	11%
Male-to-male sexual contact and IDU	18	5%	17	6%	221	8%
Heterosexual contact	18	5%	14	5%	98	4%
Blood product	2	1%	-	-	31	1%
Other/unknown	70	21%	77	25%	240	9%
Subtotal	334	100%	306	100%	2,751	100%
Female adult or adolescent						
Injection drug use	21	20%	20	19%	149	29%
Heterosexual contact	45	43%	37	34%	247	48%
Blood product	1	1%	1	1%	7	1%
Other/unknown	38	36%	50	46%	117	23%
Subtotal	105	100%	108	100%	520	100%
Child (<13 yrs at diagnosis)						
Perinatal	2	67%	4	67%	37	82%
Other	1	33%	2	33%	8	18%
Subtotal	3	100%	6	100%	45	100%
Total	442		420		3,316	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Columbus Regional Advisory Group

Table 41. Reported persons living with HIV/AIDS as of Dec. 31, 2003, HIV Prevention Columbus Regional Advisory Group, by current disease status and selected characteristics

	Current Disease Status						
	HIV/AIDS			HIV (not AIDS)		AIDS	
	Rate ^a	No.	%	No.	%	No.	%
Gender							
Male	407.9	2,118	83%	1,251	82%	867	85%
Female	79.1	435	17%	282	18%	153	15%
Unknown	*	-	-	-	-	-	-
Age (as of 12/2003)							
<13	5.5	11	<1%	7	<1%	4	<1%
13-19	9.8	10	<1%	7	<1%	3	<1%
20-24	68.7	63	2%	54	4%	9	1%
25-29	160.1	151	6%	118	8%	33	3%
30-39	498.8	883	35%	562	37%	321	31%
40-49	621.2	984	39%	553	36%	431	42%
50-64	288.0	407	16%	209	14%	198	19%
65+	42.2	44	2%	23	2%	21	2%
Race/Ethnicity							
White, not Hispanic	182.6	1,473	58%	824	54%	649	64%
Black, not Hispanic	435.9	868	34%	539	35%	329	32%
Hispanic	300.7	73	3%	44	3%	29	3%
Asian/Pacific Islander	39.4	14	1%	5	<1%	9	1%
American Indian/Alaska Native	224.9	8	<1%	6	<1%	2	<1%
Unknown	*	117	5%	115	8%	2	<1%
Race/Gender							
White, not Hispanic Male	342.7	1,346	53%	743	48%	603	59%
White, not Hispanic Female	30.7	127	5%	81	5%	46	5%
Black, not Hispanic Male	643.7	600	24%	367	24%	233	23%
Black, not Hispanic Female	253.0	268	10%	172	11%	96	9%
Hispanic Male	410.5	56	2%	35	2%	21	2%
Hispanic Female	159.8	17	1%	9	1%	8	1%
Asian/Pacific Islander Male	67.0	12	<1%	4	<1%	8	1%
Asian/Pacific Islander Female	*	2	<1%	1	<1%	1	<1%
American Indian/Alaska Native Male	288.9	5	<1%	5	<1%	-	-
American Indian/Alaska Native Female	*	3	<1%	1	<1%	2	<1%
Unknown	*	117	5%	115	8%	2	<1%
Total	238.8	2,553		1,533		1,020	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with HIV/AIDS per 100,000 population calculated using 2000 census data.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Columbus Regional Advisory Group

Table 42. Reported persons living with HIV/AIDS as of Dec. 31, 2003, HIV Prevention Columbus Regional Advisory Group, by current disease status and transmission category

Transmission category	HIV/AIDS		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	947	45%	459	37%	488	57%
Injection drug use (IDU)	86	4%	32	3%	54	6%
Male-to-male sexual contact and IDU	58	3%	26	2%	32	4%
Heterosexual contact	49	2%	21	2%	28	3%
Blood product	12	1%	7	1%	5	1%
Other/unknown	955	45%	699	56%	256	30%
Subtotal	2,107	100%	1,244	100%	863	100%
Female adult or adolescent						
Injection drug use	44	10%	21	8%	23	15%
Heterosexual contact	144	34%	81	29%	63	42%
Blood product	5	1%	4	1%	1	1%
Other/unknown	234	55%	171	62%	63	42%
Subtotal	427	100%	277	100%	150	100%
Child (<13 yrs at diagnosis)						
Perinatal	14	74%	8	67%	6	86%
Other	5	26%	4	33%	1	14%
Subtotal	19	100%	12	100%	7	100%
Total	2,553		1,533		1,020	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Columbus Regional Advisory Group

Table 43. Reported cases of HIV infection, HIV Prevention Columbus Regional Advisory Group, by year of diagnosis and selected characteristics

	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
Gender				
Male	426	83%	544	81%
Female	87	17%	124	19%
Unknown	-	-	-	-
Age				
<13	1	<1%	5	1%
13-19	10	2%	11	2%
20-24	37	7%	57	9%
25-29	78	15%	73	11%
30-39	232	45%	261	39%
40-49	115	22%	194	29%
50-64	37	7%	64	10%
65+	3	1%	3	<1%
Race/Ethnicity				
White, not Hispanic	299	58%	337	50%
Black, not Hispanic	173	34%	242	36%
Hispanic	9	2%	25	4%
Asian/Pacific Islander	5	1%	4	1%
American Indian/Alaska Native	2	<1%	1	<1%
Unknown	25	5%	59	9%
Race/Gender				
White, not Hispanic Male	280	55%	307	46%
White, not Hispanic Female	19	4%	30	4%
Black, not Hispanic Male	116	23%	159	24%
Black, not Hispanic Female	57	11%	83	12%
Hispanic Male	5	1%	22	3%
Hispanic Female	4	1%	3	<1%
Asian/Pacific Islander Male	4	1%	3	<1%
Asian/Pacific Islander Female	1	<1%	1	<1%
American Indian/Alaska Native Male	1	<1%	1	<1%
American Indian/Alaska Native Female	1	<1%	-	-
Unknown	25	5%	59	9%
Total	513		668	

Note: HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Columbus Regional Advisory Group

Table 44. Reported cases of HIV infection, HIV Prevention Columbus Regional Advisory Group, by year of diagnosis and transmission category

Transmission category	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
Male adult or adolescent				
Male-to-male sexual contact	184	43%	125	23%
Injection drug use (IDU)	8	2%	6	1%
Male-to-male sexual contact and IDU	10	2%	3	1%
Heterosexual contact	12	3%	6	1%
Blood product	-	-	-	-
Other/unknown	211	50%	402	74%
Subtotal	425	100%	542	100%
Female adult or adolescent				
Injection drug use	7	8%	2	2%
Heterosexual contact	30	34%	18	15%
Blood product	2	2%	1	1%
Other/unknown	48	55%	100	83%
Subtotal	87	100%	121	100%
Child (<13 yrs at diagnosis)				
Perinatal	1	100%	4	80%
Other	-	-	1	20%
Subtotal	1	-	5	-
Total	513		668	

Note: HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Columbus Regional Advisory Group

Table 45. Reported cases of AIDS, HIV Prevention Columbus Regional Advisory Group, by year of diagnosis and selected characteristics

	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
Gender						
Male	197	79%	281	83%	2,042	90%
Female	53	21%	57	17%	237	10%
Unknown	-	-	-	-	-	-
Age						
<13	1	<1%	1	<1%	11	<1%
13-19	1	<1%	1	<1%	8	<1%
20-24	4	2%	11	3%	88	4%
25-29	22	9%	21	6%	339	15%
30-39	137	55%	138	41%	1,066	47%
40-49	62	25%	118	35%	567	25%
50-64	21	8%	46	14%	187	8%
65+	2	1%	2	1%	13	1%
Race/Ethnicity						
White, not Hispanic	144	58%	193	57%	1,624	71%
Black, not Hispanic	96	38%	130	38%	601	26%
Hispanic	4	2%	11	3%	36	2%
Asian/Pacific Islander	2	1%	2	1%	11	<1%
American Indian/Alaska Native	3	1%	-	-	4	<1%
Unknown	1	<1%	2	1%	3	<1%
Race/Gender						
White, not Hispanic Male	131	52%	173	51%	1,535	67%
White, not Hispanic Female	13	5%	20	6%	89	4%
Black, not Hispanic Male	61	24%	95	28%	463	20%
Black, not Hispanic Female	35	14%	35	10%	138	6%
Hispanic Male	2	1%	9	3%	29	1%
Hispanic Female	2	1%	2	1%	7	<1%
Asian/Pacific Islander Male	1	<1%	2	1%	10	<1%
Asian/Pacific Islander Female	1	<1%	-	-	1	<1%
American Indian/Alaska Native Male	1	<1%	-	-	2	<1%
American Indian/Alaska Native Female	2	1%	-	-	2	<1%
Unknown	1	<1%	2	1%	3	<1%
Total	250		338		2,279	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Columbus Regional Advisory Group

Table 46. Reported cases of AIDS, HIV Prevention Columbus Regional Advisory Group, by year of diagnosis and transmission category

Transmission category	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	105	54%	105	37%	1,487	73%
Injection drug use (IDU)	10	5%	9	3%	111	5%
Male-to-male sexual contact and IDU	7	4%	9	3%	77	4%
Heterosexual contact	7	4%	8	3%	45	2%
Blood product	-	-	-	-	19	1%
Other/unknown	67	34%	150	53%	297	15%
Subtotal	196	100%	281	100%	2,036	100%
Female adult or adolescent						
Injection drug use	10	19%	4	7%	41	18%
Heterosexual contact	24	45%	14	25%	106	46%
Blood product	2	4%	-	-	9	4%
Other/unknown	17	32%	38	68%	75	32%
Subtotal	53	100%	56	100%	231	100%
Child (<13 yrs at diagnosis)						
Perinatal	1	100%	1	100%	10	83%
Other	-	-	-	-	2	17%
Subtotal	1	100%	1	100%	12	100%
Total	250		338		2,279	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Dayton Regional Advisory Group

Table 47. Reported persons living with HIV/AIDS as of Dec. 31, 2003, HIV Prevention Dayton Regional Advisory Group, by current disease status and selected characteristics

	Current Disease Status						
	HIV/AIDS			HIV (not AIDS)		AIDS	
	Rate ^a	No.	%	No.	%	No.	%
Gender							
Male	304.4	817	81%	385	77%	432	85%
Female	66.4	193	19%	115	23%	78	15%
Unknown	*	-	-	-	-	-	-
Age (as of 12/2003)							
<13	*	4	<1%	2	<1%	2	<1%
13-19	22.3	12	1%	6	1%	6	1%
20-24	83.7	32	3%	25	5%	7	1%
25-29	179.0	67	7%	42	8%	25	5%
30-39	377.4	307	30%	158	32%	149	29%
40-49	454.8	381	38%	180	36%	201	39%
50-64	219.9	193	19%	81	16%	112	22%
65+	18.3	14	1%	6	1%	8	2%
Race/Ethnicity							
White, not Hispanic	114.8	492	49%	235	47%	257	50%
Black, not Hispanic	432.3	491	49%	245	49%	246	48%
Hispanic	183.2	13	1%	7	1%	6	1%
Asian/Pacific Islander	*	4	<1%	3	1%	1	<1%
American Indian/Alaska Native	*	-	-	-	-	-	-
Unknown	*	10	1%	10	2%	-	-
Race/Gender							
White, not Hispanic Male	210.6	437	43%	204	41%	233	46%
White, not Hispanic Female	24.9	55	5%	31	6%	24	5%
Black, not Hispanic Male	683.3	360	36%	168	34%	192	38%
Black, not Hispanic Female	215.1	131	13%	77	15%	54	11%
Hispanic Male	247.6	9	1%	3	1%	6	1%
Hispanic Female	*	4	<1%	4	1%	-	-
Asian/Pacific Islander Male	*	4	<1%	3	1%	1	<1%
Asian/Pacific Islander Female	*	-	-	-	-	-	-
American Indian/Alaska Native Male	*	-	-	-	-	-	-
American Indian/Alaska Native Female	*	-	-	-	-	-	-
Unknown	*	10	1%	10	2%	-	-
Total	180.7	1,010		500		510	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with HIV/AIDS per 100,000 population calculated using 2000 census data.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Dayton Regional Advisory Group

Table 48. Reported persons living with HIV/AIDS as of Dec. 31, 2003, HIV Prevention Dayton Regional Advisory Group, by current disease status and transmission category

Transmission category	HIV/AIDS		Current Disease Status			
	No.	%	HIV (not AIDS)		AIDS	
			No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	510	63%	228	59%	282	66%
Injection drug use (IDU)	36	4%	13	3%	23	5%
Male-to-male sexual contact and IDU	49	6%	13	3%	36	8%
Heterosexual contact	31	4%	8	2%	23	5%
Blood product	9	1%	3	1%	6	1%
Other/unknown	177	22%	119	31%	58	14%
Subtotal	812	100%	384	100%	428	100%
Female adult or adolescent						
Injection drug use	28	15%	12	11%	16	22%
Heterosexual contact	78	42%	49	43%	29	39%
Blood product	3	2%	-	-	3	4%
Other/unknown	78	42%	52	46%	26	35%
Subtotal	187	100%	113	100%	74	100%
Child (<13 yrs at diagnosis)						
Perinatal	8	73%	2	67%	6	75%
Other	3	27%	1	33%	2	25%
Subtotal	11	100%	3	100%	8	100%
Total	1,010		500		510	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Dayton Regional Advisory Group

Table 49. Reported cases of HIV infection, HIV Prevention Dayton Regional Advisory Group, by year of diagnosis and selected characteristics

	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
Gender				
Male	169	79%	145	71%
Female	46	21%	59	29%
Unknown	-	-	-	-
Age				
<13	-	-	1	<1%
13-19	5	2%	15	7%
20-24	20	9%	17	8%
25-29	31	14%	28	14%
30-39	85	40%	75	37%
40-49	50	23%	55	27%
50-64	20	9%	12	6%
65+	4	2%	1	<1%
Race/Ethnicity				
White, not Hispanic	93	43%	83	41%
Black, not Hispanic	116	54%	113	55%
Hispanic	2	1%	2	1%
Asian/Pacific Islander	1	<1%	3	1%
American Indian/Alaska Native	-	-	-	-
Unknown	3	1%	3	1%
Race/Gender				
White, not Hispanic Male	81	38%	69	34%
White, not Hispanic Female	12	6%	14	7%
Black, not Hispanic Male	85	40%	70	34%
Black, not Hispanic Female	31	14%	43	21%
Hispanic Male	1	<1%	1	<1%
Hispanic Female	1	<1%	1	<1%
Asian/Pacific Islander Male	1	<1%	3	1%
Asian/Pacific Islander Female	-	-	-	-
American Indian/Alaska Native Male	-	-	-	-
American Indian/Alaska Native Female	-	-	-	-
Unknown	3	1%	3	1%
Total	215		204	

Note: HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Dayton Regional Advisory Group

Table 50. Reported cases of HIV infection, HIV Prevention Dayton Regional Advisory Group, by year of diagnosis and transmission category

Transmission category	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
Male adult or adolescent				
Male-to-male sexual contact	114	67%	62	43%
Injection drug use (IDU)	6	4%	3	2%
Male-to-male sexual contact and IDU	4	2%	4	3%
Heterosexual contact	9	5%	6	4%
Blood product	-	-	-	-
Other/unknown	36	21%	70	48%
Subtotal	169	100%	145	100%
Female adult or adolescent				
Injection drug use	-	-	4	7%
Heterosexual contact	16	35%	15	26%
Blood product	1	2%	1	2%
Other/unknown	29	63%	38	66%
Subtotal	46	100%	58	100%
Child (<13 yrs at diagnosis)				
Perinatal	-	-	1	100%
Other	-	-	-	-
Subtotal	-	-	1	-
Total	215		204	

Note: HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Dayton Regional Advisory Group

Table 51. Reported cases of AIDS, HIV Prevention Dayton Regional Advisory Group, by year of diagnosis and selected characteristics

	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
Gender						
Male	106	85%	108	76%	784	87%
Female	18	15%	34	24%	119	13%
Unknown	-	-	-	-	-	-
Age						
<13	-	-	-	-	10	1%
13-19	2	2%	3	2%	10	1%
20-24	4	3%	7	5%	28	3%
25-29	11	9%	15	11%	113	13%
30-39	62	50%	60	42%	411	46%
40-49	29	23%	42	30%	245	27%
50-64	13	10%	13	9%	75	8%
65+	3	2%	2	1%	11	1%
Race/Ethnicity						
White, not Hispanic	60	48%	64	45%	534	59%
Black, not Hispanic	63	51%	77	54%	361	40%
Hispanic	1	1%	-	-	7	1%
Asian/Pacific Islander	-	-	1	1%	1	<1%
American Indian/Alaska Native	-	-	-	-	-	-
Unknown	-	-	-	-	-	-
Race/Gender						
White, not Hispanic Male	54	44%	57	40%	490	54%
White, not Hispanic Female	6	5%	7	5%	44	5%
Black, not Hispanic Male	51	41%	50	35%	286	32%
Black, not Hispanic Female	12	10%	27	19%	75	8%
Hispanic Male	1	1%	-	-	7	1%
Hispanic Female	-	-	-	-	-	-
Asian/Pacific Islander Male	-	-	1	1%	1	<1%
Asian/Pacific Islander Female	-	-	-	-	-	-
American Indian/Alaska Native Male	-	-	-	-	-	-
American Indian/Alaska Native Female	-	-	-	-	-	-
Unknown	-	-	-	-	-	-
Total	124		142		903	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Dayton Regional Advisory Group

Table 52. Reported cases of AIDS, HIV Prevention Dayton Regional Advisory Group, by year of diagnosis and transmission category

Transmission category	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	74	70%	58	54%	542	70%
Injection drug use (IDU)	5	5%	2	2%	41	5%
Male-to-male sexual contact and IDU	4	4%	5	5%	63	8%
Heterosexual contact	10	9%	2	2%	33	4%
Blood product	1	1%	-	-	17	2%
Other/unknown	12	11%	40	37%	80	10%
Subtotal	106	100%	107	100%	776	100%
Female adult or adolescent						
Injection drug use	4	22%	4	12%	28	25%
Heterosexual contact	5	28%	10	30%	42	37%
Blood product	1	6%	1	3%	7	6%
Other/unknown	8	44%	18	55%	36	32%
Subtotal	18	100%	33	100%	113	100%
Child (<13 yrs at diagnosis)						
Perinatal	-	-	1	50%	9	64%
Other	-	-	1	50%	5	36%
Subtotal	-	-	2	100%	14	100%
Total	124		142		903	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Toledo Regional Advisory Group

Table 53. Reported persons living with HIV/AIDS as of Dec. 31, 2003, HIV Prevention Toledo Regional Advisory Group, by current disease status and selected characteristics

	HIV/AIDS			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Gender							
Male	266.0	582	76%	303	72%	279	80%
Female	79.6	188	24%	119	28%	69	20%
Unknown	*	-	-	-	-	-	-
Age (as of 12/2003)							
<13	8.1	7	1%	3	1%	4	1%
13-19	10.9	5	1%	4	1%	1	<1%
20-24	66.3	21	3%	17	4%	4	1%
25-29	185.3	60	8%	48	11%	12	3%
30-39	395.2	255	33%	152	36%	103	30%
40-49	439.6	301	39%	146	35%	155	45%
50-64	165.2	109	14%	48	11%	61	18%
65+	20.2	12	2%	4	1%	8	2%
Race/Ethnicity							
White, not Hispanic	113.4	394	51%	223	53%	171	49%
Black, not Hispanic	403.6	321	42%	163	39%	158	45%
Hispanic	183.8	38	5%	20	5%	18	5%
Asian/Pacific Islander	81.1	5	1%	4	1%	1	<1%
American Indian/Alaska Native	*	-	-	-	-	-	-
Unknown	*	12	2%	12	3%	-	-
Race/Gender							
White, not Hispanic Male	195.6	328	43%	179	42%	149	43%
White, not Hispanic Female	36.7	66	9%	44	10%	22	6%
Black, not Hispanic Male	578.5	214	28%	98	23%	116	33%
Black, not Hispanic Female	251.5	107	14%	65	15%	42	12%
Hispanic Male	260.0	27	4%	14	3%	13	4%
Hispanic Female	107.0	11	1%	6	1%	5	1%
Asian/Pacific Islander Male	163.8	5	1%	4	1%	1	<1%
Asian/Pacific Islander Female	*	-	-	-	-	-	-
American Indian/Alaska Native Male	*	-	-	-	-	-	-
American Indian/Alaska Native Female	*	-	-	-	-	-	-
Unknown	*	12	2%	12	3%	-	-
Total	169.2	770		422		348	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with HIV/AIDS per 100,000 population calculated using 2000 census data.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Toledo Regional Advisory Group

Table 54. Reported persons living with HIV/AIDS as of Dec. 31, 2003, HIV Prevention Toledo Regional Advisory Group, by current disease status and transmission category

Transmission category	HIV/AIDS		Current Disease Status			
			HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	342	59%	173	57%	169	61%
Injection drug use (IDU)	42	7%	24	8%	18	7%
Male-to-male sexual contact and IDU	35	6%	9	3%	26	9%
Heterosexual contact	33	6%	17	6%	16	6%
Blood product	7	1%	3	1%	4	1%
Other/unknown	119	21%	76	25%	43	16%
Subtotal	578	100%	302	100%	276	100%
Female adult or adolescent						
Injection drug use	25	14%	11	10%	14	21%
Heterosexual contact	77	43%	51	44%	26	39%
Blood product	1	1%	-	-	1	2%
Other/unknown	78	43%	53	46%	25	38%
Subtotal	181	100%	115	100%	66	100%
Child (<13 yrs at diagnosis)						
Perinatal	10	91%	5	100%	5	83%
Other	1	9%	-	-	1	17%
Subtotal	11	100%	5	100%	6	100%
Total	770		422		348	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Toledo Regional Advisory Group

Table 55. Reported cases of HIV infection, HIV Prevention Toledo Regional Advisory Group, by year of diagnosis and selected characteristics

	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
Gender				
Male	95	67%	126	75%
Female	47	33%	42	25%
Unknown	-	-	-	-
Age				
<13	3	2%	2	1%
13-19	3	2%	1	1%
20-24	16	11%	15	9%
25-29	27	19%	23	14%
30-39	63	44%	62	37%
40-49	20	14%	51	30%
50-64	9	6%	14	8%
65+	1	1%	-	-
Race/Ethnicity				
White, not Hispanic	72	51%	71	42%
Black, not Hispanic	56	39%	83	49%
Hispanic	11	8%	9	5%
Asian/Pacific Islander	1	1%	2	1%
American Indian/Alaska Native	-	-	-	-
Unknown	2	1%	3	2%
Race/Gender				
White, not Hispanic Male	55	39%	61	36%
White, not Hispanic Female	17	12%	10	6%
Black, not Hispanic Male	30	21%	53	32%
Black, not Hispanic Female	26	18%	30	18%
Hispanic Male	7	5%	9	5%
Hispanic Female	4	3%	-	-
Asian/Pacific Islander Male	1	1%	2	1%
Asian/Pacific Islander Female	-	-	-	-
American Indian/Alaska Native Male	-	-	-	-
American Indian/Alaska Native Female	-	-	-	-
Unknown	2	1%	3	2%
Total	142		168	

Note: HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Toledo Regional Advisory Group

Table 56. Reported cases of HIV infection, HIV Prevention Toledo Regional Advisory Group, by year of diagnosis and transmission category

Transmission category	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
Male adult or adolescent				
Male-to-male sexual contact	49	52%	57	46%
Injection drug use (IDU)	5	5%	7	6%
Male-to-male sexual contact and IDU	3	3%	3	2%
Heterosexual contact	8	8%	8	6%
Blood product	-	-	-	-
Other/unknown	30	32%	50	40%
Subtotal	95	100%	125	100%
Female adult or adolescent				
Injection drug use	5	11%	1	2%
Heterosexual contact	14	32%	15	37%
Blood product	-	-	-	-
Other/unknown	25	57%	25	61%
Subtotal	44	100%	41	100%
Child (<13 yrs at diagnosis)				
Perinatal	3	100%	2	100%
Other	-	-	-	-
Subtotal	3	-	2	-
Total	142		168	

Note: HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Toledo Regional Advisory Group

Table 57. Reported cases of AIDS, HIV Prevention Toledo Regional Advisory Group, by year of diagnosis and selected characteristics

	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
Gender						
Male	62	72%	72	77%	516	82%
Female	24	28%	22	23%	115	18%
Unknown	-	-	-	-	-	-
Age						
<13	1	1%	1	1%	11	2%
13-19	1	1%	-	-	6	1%
20-24	4	5%	3	3%	43	7%
25-29	9	10%	9	10%	89	14%
30-39	42	49%	42	45%	280	44%
40-49	22	26%	30	32%	146	23%
50-64	6	7%	9	10%	50	8%
65+	1	1%	-	-	6	1%
Race/Ethnicity						
White, not Hispanic	44	51%	38	40%	369	58%
Black, not Hispanic	34	40%	52	55%	239	38%
Hispanic	8	9%	4	4%	22	3%
Asian/Pacific Islander	-	-	-	-	1	<1%
American Indian/Alaska Native	-	-	-	-	-	-
Unknown	-	-	-	-	-	-
Race/Gender						
White, not Hispanic Male	31	36%	33	35%	324	51%
White, not Hispanic Female	13	15%	5	5%	45	7%
Black, not Hispanic Male	26	30%	35	37%	175	28%
Black, not Hispanic Female	8	9%	17	18%	64	10%
Hispanic Male	5	6%	4	4%	16	3%
Hispanic Female	3	3%	-	-	6	1%
Asian/Pacific Islander Male	-	-	-	-	1	<1%
Asian/Pacific Islander Female	-	-	-	-	-	-
American Indian/Alaska Native Male	-	-	-	-	-	-
American Indian/Alaska Native Female	-	-	-	-	-	-
Unknown	-	-	-	-	-	-
Total	86		94		631	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Toledo Regional Advisory Group

Table 58. Reported cases of AIDS, HIV Prevention Toledo Regional Advisory Group, by year of diagnosis and transmission category

Transmission category	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	34	55%	33	46%	362	71%
Injection drug use (IDU)	5	8%	6	8%	31	6%
Male-to-male sexual contact and IDU	6	10%	3	4%	37	7%
Heterosexual contact	8	13%	2	3%	19	4%
Blood product	-	-	-	-	5	1%
Other/unknown	9	15%	27	38%	55	11%
Subtotal	62	100%	71	100%	509	100%
Female adult or adolescent						
Injection drug use	4	17%	1	5%	23	21%
Heterosexual contact	9	39%	7	32%	51	46%
Blood product	-	-	-	-	1	1%
Other/unknown	10	43%	14	64%	35	32%
Subtotal	23	100%	22	100%	110	100%
Child (<13 yrs at diagnosis)						
Perinatal	1	100%	1	100%	11	92%
Other	-	-	-	-	1	8%
Subtotal	1	100%	1	100%	12	100%
Total	86		94		631	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Youngstown Regional Advisory Group

Table 59. Reported persons living with HIV/AIDS as of Dec. 31, 2003, HIV Prevention Youngstown Regional Advisory Group, by current disease status and selected characteristics

	HIV/AIDS			Current Disease Status			
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Gender							
Male	121.0	348	69%	191	63%	157	79%
Female	50.5	155	31%	112	37%	43	22%
Unknown	*	-	-	-	-	-	-
Age (as of 12/2003)							
<13	8.9	9	2%	9	3%	-	-
13-19	*	4	1%	4	1%	-	-
20-24	15.3	5	1%	4	1%	1	1%
25-29	78.9	27	5%	24	8%	3	2%
30-39	193.7	154	31%	101	33%	53	27%
40-49	205.5	193	38%	102	34%	91	46%
50-64	100.9	99	20%	51	17%	48	24%
65+	12.2	12	2%	8	3%	4	2%
Race/Ethnicity							
White, not Hispanic	47.3	245	49%	129	43%	116	58%
Black, not Hispanic	313.8	196	39%	123	41%	73	37%
Hispanic	316.5	34	7%	23	8%	11	6%
Asian/Pacific Islander	*	-	-	-	-	-	-
American Indian/Alaska Native	*	1	<1%	1	<1%	-	-
Unknown	*	27	5%	27	9%	-	-
Race/Gender							
White, not Hispanic Male	76.2	190	38%	90	30%	100	50%
White, not Hispanic Female	20.5	55	11%	39	13%	16	8%
Black, not Hispanic Male	392.5	120	24%	70	23%	50	25%
Black, not Hispanic Female	238.3	76	15%	53	17%	23	12%
Hispanic Male	333.5	19	4%	12	4%	7	4%
Hispanic Female	297.3	15	3%	11	4%	4	2%
Asian/Pacific Islander Male	*	-	-	-	-	-	-
Asian/Pacific Islander Female	*	-	-	-	-	-	-
American Indian/Alaska Native Male	*	1	<1%	1	<1%	-	-
American Indian/Alaska Native Female	*	-	-	-	-	-	-
Unknown	*	27	5%	27	9%	-	-
Total	84.6	503		303		200	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with HIV/AIDS per 100,000 population calculated using 2000 census data.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Youngstown Regional Advisory Group

Table 60. Reported persons living with HIV/AIDS as of Dec. 31, 2003, HIV Prevention Youngstown Regional Advisory Group, by current disease status and transmission category

Transmission category	HIV/AIDS		Current Disease Status			
			HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	139	41%	57	31%	82	52%
Injection drug use (IDU)	33	10%	19	10%	14	9%
Male-to-male sexual contact and IDU	23	7%	7	4%	16	10%
Heterosexual contact	11	3%	5	3%	6	4%
Blood product	6	2%	1	1%	5	3%
Other/unknown	131	38%	97	52%	34	22%
Subtotal	343	100%	186	100%	157	100%
Female adult or adolescent						
Injection drug use	19	13%	12	11%	7	16%
Heterosexual contact	37	25%	22	21%	15	35%
Blood product	-	-	-	-	-	-
Other/unknown	93	62%	72	68%	21	49%
Subtotal	149	100%	106	100%	43	100%
Child (<13 yrs at diagnosis)						
Perinatal	9	82%	9	82%	-	-
Other	2	18%	2	18%	-	-
Subtotal	11	100%	11	100%	-	-
Total	503		303		200	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Youngstown Regional Advisory Group

Table 61. Reported cases of HIV infection, HIV Prevention Youngstown Regional Advisory Group, by year of diagnosis and selected characteristics

	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
Gender				
Male	70	80%	67	55%
Female	17	20%	55	45%
Unknown	-	-	-	-
Age				
<13	1	1%	1	1%
13-19	2	2%	3	2%
20-24	4	5%	7	6%
25-29	9	10%	19	16%
30-39	28	32%	39	32%
40-49	32	37%	42	34%
50-64	11	13%	11	9%
65+	-	-	-	-
Race/Ethnicity				
White, not Hispanic	45	52%	47	39%
Black, not Hispanic	34	39%	65	53%
Hispanic	4	5%	6	5%
Asian/Pacific Islander	-	-	-	-
American Indian/Alaska Native	1	1%	1	1%
Unknown	3	3%	3	2%
Race/Gender				
White, not Hispanic Male	40	46%	27	22%
White, not Hispanic Female	5	6%	20	16%
Black, not Hispanic Male	23	26%	35	29%
Black, not Hispanic Female	11	13%	30	25%
Hispanic Male	3	3%	2	2%
Hispanic Female	1	1%	4	3%
Asian/Pacific Islander Male	-	-	-	-
Asian/Pacific Islander Female	-	-	-	-
American Indian/Alaska Native Male	1	1%	1	1%
American Indian/Alaska Native Female	-	-	-	-
Unknown	3	3%	3	2%
Total	87		122	

Note: HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Youngstown Regional Advisory Group

Table 62. Reported cases of HIV infection, HIV Prevention Youngstown Regional Advisory Group, by year of diagnosis and transmission category

Transmission category	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
Male adult or adolescent				
Male-to-male sexual contact	34	49%	17	25%
Injection drug use (IDU)	10	14%	3	4%
Male-to-male sexual contact and IDU	3	4%	1	1%
Heterosexual contact	3	4%	4	6%
Blood product	-	-	-	-
Other/unknown	19	28%	42	63%
Subtotal	69	100%	67	100%
Female adult or adolescent				
Injection drug use	2	12%	1	2%
Heterosexual contact	6	35%	4	7%
Blood product	-	-	-	-
Other/unknown	9	53%	49	91%
Subtotal	17	100%	54	100%
Child (<13 yrs at diagnosis)				
Perinatal	1	100%	1	100%
Other	-	-	-	-
Subtotal	1	-	1	-
Total	87		122	

Note: HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Youngstown Regional Advisory Group

Table 63. Reported cases of AIDS, HIV Prevention Youngstown Regional Advisory Group, by year of diagnosis and selected characteristics

	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
Gender						
Male	34	79%	40	66%	320	81%
Female	9	21%	21	34%	74	19%
Unknown	-	-	-	-	-	-
Age						
<13	-	-	-	-	-	-
13-19	-	-	-	-	7	2%
20-24	-	-	-	-	11	3%
25-29	4	9%	7	11%	52	13%
30-39	13	30%	25	41%	174	44%
40-49	20	47%	24	39%	112	28%
50-64	6	14%	5	8%	33	8%
65+	-	-	-	-	5	1%
Race/Ethnicity						
White, not Hispanic	27	63%	28	46%	240	61%
Black, not Hispanic	16	37%	30	49%	137	35%
Hispanic	-	-	3	5%	17	4%
Asian/Pacific Islander	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-
Unknown	-	-	-	-	-	-
Race/Gender						
White, not Hispanic Male	23	53%	20	33%	211	54%
White, not Hispanic Female	4	9%	8	13%	29	7%
Black, not Hispanic Male	11	26%	18	30%	99	25%
Black, not Hispanic Female	5	12%	12	20%	38	10%
Hispanic Male	-	-	2	3%	10	3%
Hispanic Female	-	-	1	2%	7	2%
Asian/Pacific Islander Male	-	-	-	-	-	-
Asian/Pacific Islander Female	-	-	-	-	-	-
American Indian/Alaska Native Male	-	-	-	-	-	-
American Indian/Alaska Native Female	-	-	-	-	-	-
Unknown	-	-	-	-	-	-
Total	43		61		394	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Youngstown Regional Advisory Group

Table 64. Reported cases of AIDS, HIV Prevention Youngstown Regional Advisory Group, by year of diagnosis and transmission category

Transmission category	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	24	71%	9	23%	183	57%
Injection drug use (IDU)	3	9%	5	13%	29	9%
Male-to-male sexual contact and IDU	2	6%	2	5%	25	8%
Heterosexual contact	1	3%	2	5%	13	4%
Blood product	-	-	1	3%	25	8%
Other/unknown	4	12%	21	53%	44	14%
Subtotal	34	100%	40	100%	319	100%
Female adult or adolescent						
Injection drug use	3	33%	-	-	20	27%
Heterosexual contact	2	22%	5	24%	24	32%
Blood product	-	-	-	-	4	5%
Other/unknown	4	44%	16	76%	26	35%
Subtotal	9	100%	21	100%	74	100%
Child (<13 yrs at diagnosis)						
Perinatal	-	-	-	-	-	-
Other	-	-	-	-	1	100%
Subtotal	-	-	-	-	1	100%
Total	43		61		394	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Rural Regional Advisory Group

Table 65. Reported persons living with HIV/AIDS as of Dec. 31, 2003, HIV Prevention Rural Regional Advisory Group, by current disease status and selected characteristics

	HIV/AIDS		Current Disease Status				
	Rate ^a	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
Gender							
Male	74.4	1,768	78%	809	73%	959	83%
Female	20.1	493	22%	303	27%	190	17%
Unknown	*	1	<1%	1	<1%	-	-
Age (as of 12/2003)							
<13	3.0	26	1%	14	1%	12	1%
13-19	3.4	17	1%	11	1%	6	1%
20-24	17.0	50	2%	43	4%	7	1%
25-29	42.0	121	5%	88	8%	33	3%
30-39	102.1	712	31%	349	31%	363	32%
40-49	117.7	888	39%	422	38%	466	41%
50-64	53.3	410	18%	171	15%	239	21%
65+	6.0	38	2%	15	1%	23	2%
Race/Ethnicity							
White, not Hispanic	38.5	1,743	77%	839	75%	904	79%
Black, not Hispanic	197.3	333	15%	164	15%	169	15%
Hispanic	145.3	119	5%	53	5%	66	6%
Asian/Pacific Islander	22.1	7	<1%	2	<1%	5	<1%
American Indian/Alaska Native	56.6	7	<1%	4	<1%	3	<1%
Unknown	*	53	2%	51	5%	2	<1%
Race/Gender							
White, not Hispanic Male	64.0	1,420	63%	641	58%	779	68%
White, not Hispanic Female	14.0	323	14%	198	18%	125	11%
Black, not Hispanic Male	240.9	227	10%	103	9%	124	11%
Black, not Hispanic Female	142.2	106	5%	61	5%	45	4%
Hispanic Male	175.7	74	3%	25	2%	49	4%
Hispanic Female	113.1	45	2%	28	3%	17	1%
Asian/Pacific Islander Male	34.3	5	<1%	2	<1%	3	<1%
Asian/Pacific Islander Female	*	2	<1%	-	-	2	<1%
American Indian/Alaska Native Male	*	4	<1%	2	<1%	2	<1%
American Indian/Alaska Native Female	*	3	<1%	2	<1%	1	<1%
Unknown	*	53	2%	51	5%	2	<1%
Total	46.9	2,262		1,113		1,149	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

^a The rate is the number of persons living with HIV/AIDS per 100,000 population calculated using 2000 census data.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Rural Regional Advisory Group

Table 66. Reported persons living with HIV/AIDS as of Dec. 31, 2003, HIV Prevention Rural Regional Advisory Group, by current disease status and transmission category

Transmission category	HIV/AIDS		Current Disease Status			
			HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	885	51%	323	40%	562	59%
Injection drug use (IDU)	110	6%	40	5%	70	7%
Male-to-male sexual contact and IDU	72	4%	32	4%	40	4%
Heterosexual contact	67	4%	26	3%	41	4%
Blood product	35	2%	11	1%	24	3%
Other/unknown	579	33%	370	46%	209	22%
Subtotal	1,748	100%	802	100%	946	100%
Female adult or adolescent						
Injection drug use	37	8%	19	7%	18	10%
Heterosexual contact	223	47%	123	42%	100	55%
Blood product	5	1%	-	-	5	3%
Other/unknown	208	44%	150	51%	58	32%
Subtotal	473	100%	292	100%	181	100%
Child (<13 yrs at diagnosis)						
Perinatal	32	80%	16	89%	16	73%
Other	8	20%	2	11%	6	27%
Subtotal	40	100%	18	100%	22	100%
Total	2,262		1,113		1,149	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Rural Regional Advisory Group

Table 67. Reported cases of HIV infection, HIV Prevention Rural Regional Advisory Group, by year of diagnosis and selected characteristics

	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
Gender				
Male	389	77%	394	79%
Female	117	23%	106	21%
Unknown	1	<1%	-	-
Age				
<13	2	<1%	2	<1%
13-19	15	3%	14	3%
20-24	42	8%	29	6%
25-29	57	11%	59	12%
30-39	193	38%	173	35%
40-49	149	29%	145	29%
50-64	43	8%	69	14%
65+	6	1%	9	2%
Race/Ethnicity				
White, not Hispanic	374	74%	365	73%
Black, not Hispanic	95	19%	84	17%
Hispanic	26	5%	25	5%
Asian/Pacific Islander	2	<1%	1	<1%
American Indian/Alaska Native	3	1%	1	<1%
Unknown	7	1%	24	5%
Race/Gender				
White, not Hispanic Male	300	59%	295	59%
White, not Hispanic Female	74	15%	70	14%
Black, not Hispanic Male	69	14%	63	13%
Black, not Hispanic Female	26	5%	21	4%
Hispanic Male	15	3%	16	3%
Hispanic Female	11	2%	9	2%
Asian/Pacific Islander Male	1	<1%	1	<1%
Asian/Pacific Islander Female	1	<1%	-	-
American Indian/Alaska Native Male	-	-	1	<1%
American Indian/Alaska Native Female	3	1%	-	-
Unknown	7	1%	24	5%
Total	507		500	

Note: HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Rural Regional Advisory Group

Table 68. Reported cases of HIV infection, HIV Prevention Rural Regional Advisory Group, by year of diagnosis and transmission category

Transmission category	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
Male adult or adolescent				
Male-to-male sexual contact	191	49%	129	33%
Injection drug use (IDU)	20	5%	17	4%
Male-to-male sexual contact and IDU	13	3%	8	2%
Heterosexual contact	19	5%	10	3%
Blood product	1	<1%	4	1%
Other/unknown	144	37%	225	57%
Subtotal	388	100%	393	100%
Female adult or adolescent				
Injection drug use	9	8%	4	4%
Heterosexual contact	45	39%	30	29%
Blood product	1	1%	-	-
Other/unknown	61	53%	71	68%
Subtotal	116	100%	105	100%
Child (<13 yrs at diagnosis)				
Perinatal	2	100%	2	100%
Other	-	-	-	-
Subtotal	2	-	2	-
Total	507		500	

Note: HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Rural Regional Advisory Group

Table 69. Reported cases of AIDS, HIV Prevention Rural Regional Advisory Group, by year of diagnosis and selected characteristics

	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
Gender						
Male	253	83%	255	84%	2,037	87%
Female	50	17%	47	16%	294	13%
Unknown	-	-	-	-	-	-
Age						
<13	1	<1%	1	<1%	36	2%
13-19	1	<1%	2	1%	23	1%
20-24	5	2%	4	1%	90	4%
25-29	32	11%	21	7%	335	14%
30-39	122	40%	123	41%	958	41%
40-49	103	34%	103	34%	624	27%
50-64	34	11%	44	15%	227	10%
65+	5	2%	4	1%	38	2%
Race/Ethnicity						
White, not Hispanic	232	77%	229	76%	1,979	85%
Black, not Hispanic	53	17%	54	18%	254	11%
Hispanic	15	5%	17	6%	88	4%
Asian/Pacific Islander	2	1%	1	<1%	6	<1%
American Indian/Alaska Native	1	<1%	-	-	3	<1%
Unknown	-	-	1	<1%	1	<1%
Race/Gender						
White, not Hispanic Male	202	67%	194	64%	1,765	76%
White, not Hispanic Female	30	10%	35	12%	214	9%
Black, not Hispanic Male	39	13%	47	16%	200	9%
Black, not Hispanic Female	14	5%	7	2%	54	2%
Hispanic Male	11	4%	12	4%	65	3%
Hispanic Female	4	1%	5	2%	23	1%
Asian/Pacific Islander Male	1	<1%	1	<1%	4	<1%
Asian/Pacific Islander Female	1	<1%	-	-	2	<1%
American Indian/Alaska Native Male	-	-	-	-	2	<1%
American Indian/Alaska Native Female	1	<1%	-	-	1	<1%
Unknown	-	-	1	<1%	1	<1%
Total	303		302		2,331	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

HIV Prevention Rural Regional Advisory Group

Table 70. Reported cases of AIDS, HIV Prevention Rural Regional Advisory Group, by year of diagnosis and transmission category

Transmission category	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
Male adult or adolescent						
Male-to-male sexual contact	148	58%	107	42%	1,340	67%
Injection drug use (IDU)	16	6%	15	6%	121	6%
Male-to-male sexual contact and IDU	9	4%	5	2%	93	5%
Heterosexual contact	14	6%	10	4%	75	4%
Blood product	3	1%	2	1%	94	5%
Other/unknown	63	25%	115	45%	288	14%
Subtotal	253	100%	254	100%	2,011	100%
Female adult or adolescent						
Injection drug use	7	14%	6	13%	39	14%
Heterosexual contact	23	47%	15	33%	151	55%
Blood product	1	2%	-	-	18	7%
Other/unknown	18	37%	25	54%	66	24%
Subtotal	49	100%	46	100%	274	100%
Child (<13 yrs at diagnosis)						
Perinatal	1	100%	2	100%	29	63%
Other	-	-	-	-	17	37%
Subtotal	1	100%	2	100%	46	100%
Total	303		302		2,331	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.