

## **HIV/AIDS Data by Ryan White Title I Cleveland Planning Region and Ryan White Title II Consortia Areas**

The following tables present HIV/AIDS data for Ryan White Title I Cleveland Planning Region and for the 11 Ryan White Title II Consortia Areas. For a listing and a map depiction of the counties associated with each of these regions, please refer to section 3 starting on page 274. Additional data sources (i.e. Census, STDs) are presented by planning regions in section 3.

There are six tables of data for Title I planning region and for each consortium. The first table is reported persons living with HIV/AIDS as of Dec. 31, 2003, by current disease status and selected characteristics. The second table is reported persons living with HIV/AIDS by current disease status and transmission category. The third table is reported cases of HIV infection (including persons diagnosed with HIV and later progressed to AIDS and cases concurrently diagnosed with HIV and AIDS) by year of diagnosis and selected characteristics for time periods 1998 to 2000 and 2001 to 2003. The fourth table is reported cases of HIV infection by year of diagnosis and transmission category. The fifth table is reported cases of AIDS by year of diagnosis and selected characteristics for time periods 1998 to 2000 and 2001 to 2003 as well as cumulative AIDS diagnoses. The sixth table is AIDS diagnosis by year of diagnosis and transmission category.

When viewing these tables it is important to take into account the limitations of these data. These data represent confidential reports of HIV infection and AIDS diagnoses and not all persons with HIV infection. The HIV diagnosis data represent the earliest date of diagnosis reported to the HIV/AIDS Surveillance Program. They do not always reflect the earliest date the individual became aware of his or her HIV-infection status. The possibility exists that individuals might have been previously tested anonymously and not confidentially reported or might have been previously diagnosed out-of-state and not reported to Ohio until a later test was performed or diagnosis made.



## Ryan White Title I Cleveland Planning Region

**Table 71. Reported persons living with HIV/AIDS as of Dec. 31, 2003, Ryan White Title I Cleveland Planning Region, by current disease status and selected characteristics**

	HIV/AIDS			Current Disease Status			
	Rate <sup>a</sup>	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
<b>Gender</b>							
Male	259.7	2,799	77%	1,221	74%	1,578	80%
Female	71.3	836	23%	439	26%	397	20%
Unknown	*	1	<1%	1	<1%	-	-
<b>Age (as of 12/2003)</b>							
<13	7.5	31	1%	16	1%	15	1%
13-19	13.6	29	1%	17	1%	12	1%
20-24	59.1	72	2%	52	3%	20	1%
25-29	115.5	161	4%	107	6%	54	3%
30-39	333.4	1,109	31%	565	34%	544	28%
40-49	426.6	1,512	42%	623	38%	889	45%
50-64	188.0	657	18%	257	15%	400	20%
65+	19.9	65	2%	24	1%	41	2%
<b>Race/Ethnicity</b>							
White, not Hispanic	77.1	1,322	36%	564	34%	758	38%
Black, not Hispanic	433.1	1,829	50%	834	50%	995	50%
Hispanic	487.6	365	10%	158	10%	207	10%
Asian/Pacific Islander	38.7	13	<1%	5	<1%	8	<1%
American Indian/Alaska Native	226.4	11	<1%	5	<1%	6	<1%
Unknown	*	96	3%	95	6%	1	<1%
<b>Race/Gender</b>							
White, not Hispanic Male	138.3	1,145	31%	472	28%	673	34%
White, not Hispanic Female	19.9	177	5%	92	6%	85	4%
Black, not Hispanic Male	686.8	1,328	37%	580	35%	748	38%
Black, not Hispanic Female	218.4	500	14%	253	15%	247	13%
Hispanic Male	623.3	236	6%	92	6%	144	7%
Hispanic Female	348.6	129	4%	66	4%	63	3%
Asian/Pacific Islander Male	61.1	10	<1%	3	<1%	7	<1%
Asian/Pacific Islander Female	*	3	<1%	2	<1%	1	<1%
American Indian/Alaska Native Male	299.1	7	<1%	2	<1%	5	<1%
American Indian/Alaska Native Female	*	4	<1%	3	<1%	1	<1%
Unknown	*	97	3%	96	6%	1	<1%
<b>Total</b>	<b>161.5</b>	<b>3,636</b>		<b>1,661</b>		<b>1,975</b>	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (\*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> The rate is the number of persons living with HIV/AIDS per 100,000 population calculated using 2000 census data.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title I Cleveland Planning Region

**Table 72. Reported persons living with HIV/AIDS as of Dec. 31, 2003, Ryan White Title I Cleveland Planning Region, by current disease status and transmission category**

Transmission category	HIV/AIDS		Current Disease Status			
			HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	1,542	56%	566	47%	976	62%
Injection drug use (IDU)	280	10%	91	8%	189	12%
Male-to-male sexual contact and IDU	162	6%	56	5%	106	7%
Heterosexual contact	113	4%	43	4%	70	4%
Blood product	17	1%	6	<1%	11	1%
Other/unknown	661	24%	447	37%	214	14%
Subtotal	2,775	100%	1,209	100%	1,566	100%
<b>Female adult or adolescent</b>						
Injection drug use	150	19%	53	13%	97	25%
Heterosexual contact	316	39%	143	34%	173	45%
Blood product	6	1%	2	<1%	4	1%
Other/unknown	333	41%	226	53%	107	28%
Subtotal	805	100%	424	100%	381	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	42	76%	20	74%	22	79%
Other	13	24%	7	26%	6	21%
Subtotal	55	100%	27	100%	28	100%
<b>Total</b>	<b>3,636</b>		<b>1,661</b>		<b>1,975</b>	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (\*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title I Cleveland Planning Region

**Table 73. Reported cases of HIV infection, Ryan White Title I Cleveland Planning Region, by year of diagnosis and selected characteristics**

	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
<b>Gender</b>				
Male	533	75%	432	73%
Female	175	25%	162	27%
Unknown	-	-	-	-
<b>Age</b>				
<13	11	2%	8	1%
13-19	19	3%	10	2%
20-24	55	8%	55	9%
25-29	82	12%	78	13%
30-39	287	41%	217	37%
40-49	187	26%	168	28%
50-64	60	8%	57	10%
65+	7	1%	1	<1%
<b>Race/Ethnicity</b>				
White, not Hispanic	243	34%	197	33%
Black, not Hispanic	393	56%	316	53%
Hispanic	56	8%	55	9%
Asian/Pacific Islander	4	1%	2	<1%
American Indian/Alaska Native	4	1%	2	<1%
Unknown	8	1%	22	4%
<b>Race/Gender</b>				
White, not Hispanic Male	203	29%	169	28%
White, not Hispanic Female	40	6%	28	5%
Black, not Hispanic Male	279	39%	208	35%
Black, not Hispanic Female	114	16%	108	18%
Hispanic Male	40	6%	37	6%
Hispanic Female	16	2%	18	3%
Asian/Pacific Islander Male	3	<1%	-	-
Asian/Pacific Islander Female	1	<1%	2	<1%
American Indian/Alaska Native Male	2	<1%	1	<1%
American Indian/Alaska Native Female	2	<1%	1	<1%
Unknown	8	1%	22	4%
<b>Total</b>	<b>708</b>		<b>594</b>	

*Note:* HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title I Cleveland Planning Region

**Table 74. Reported cases of HIV infection, Ryan White Title I Cleveland Planning Region, by year of diagnosis and transmission category**

Transmission category	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
<b>Male adult or adolescent</b>				
Male-to-male sexual contact	277	53%	185	43%
Injection drug use (IDU)	47	9%	23	5%
Male-to-male sexual contact and IDU	15	3%	7	2%
Heterosexual contact	21	4%	14	3%
Blood product	-	-	-	-
Other/unknown	166	32%	200	47%
Subtotal	526	100%	429	100%
<b>Female adult or adolescent</b>				
Injection drug use	14	8%	10	6%
Heterosexual contact	61	36%	32	20%
Blood product	2	1%	1	1%
Other/unknown	94	55%	114	73%
Subtotal	171	100%	157	100%
<b>Child (&lt;13 yrs at diagnosis)</b>				
Perinatal	8	73%	7	88%
Other	3	27%	1	13%
Subtotal	11	-	8	-
<b>Total</b>	<b>708</b>		<b>594</b>	

*Note:* HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title I Cleveland Planning Region

**Table 75. Reported cases of AIDS, Ryan White Title I Cleveland Planning Region, by year of diagnosis and selected characteristics**

	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
<b>Gender</b>						
Male	379	76%	345	74%	3,132	84%
Female	117	24%	124	26%	608	16%
Unknown	-	-	-	-	-	-
<b>Age</b>						
<13	3	1%	5	1%	46	1%
13-19	2	<1%	4	1%	24	1%
20-24	16	3%	22	5%	150	4%
25-29	43	9%	38	8%	500	13%
30-39	222	45%	196	42%	1,698	45%
40-49	163	33%	149	32%	940	25%
50-64	41	8%	53	11%	335	9%
65+	6	1%	2	<1%	47	1%
<b>Race/Ethnicity</b>						
White, not Hispanic	157	32%	158	34%	1,590	43%
Black, not Hispanic	302	61%	264	56%	1,852	50%
Hispanic	30	6%	45	10%	283	8%
Asian/Pacific Islander	3	1%	-	-	8	<1%
American Indian/Alaska Native	4	1%	1	<1%	6	<1%
Unknown	-	-	1	<1%	1	<1%
<b>Race/Gender</b>						
White, not Hispanic Male	137	28%	134	29%	1,469	39%
White, not Hispanic Female	20	4%	24	5%	121	3%
Black, not Hispanic Male	214	43%	182	39%	1,451	39%
Black, not Hispanic Female	88	18%	82	17%	401	11%
Hispanic Male	23	5%	27	6%	200	5%
Hispanic Female	7	1%	18	4%	83	2%
Asian/Pacific Islander Male	2	<1%	-	-	7	<1%
Asian/Pacific Islander Female	1	<1%	-	-	1	<1%
American Indian/Alaska Native Male	3	1%	1	<1%	4	<1%
American Indian/Alaska Native Female	1	<1%	-	-	2	<1%
Unknown	-	-	1	<1%	1	<1%
<b>Total</b>	<b>496</b>		<b>469</b>		<b>3,740</b>	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title I Cleveland Planning Region

**Table 76. Reported cases of AIDS, Ryan White Title I Cleveland Planning Region, by year of diagnosis and transmission category**

Transmission category	Year of AIDS Diagnosis				Cumulative	
	1998 - 2000		2001 - 2003			
	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	218	58%	193	56%	2,091	67%
Injection drug use (IDU)	39	10%	30	9%	353	11%
Male-to-male sexual contact and IDU	19	5%	17	5%	238	8%
Heterosexual contact	20	5%	14	4%	111	4%
Blood product	3	1%	1	<1%	44	1%
Other/unknown	79	21%	87	25%	272	9%
Subtotal	378	100%	342	100%	3,109	100%
<b>Female adult or adolescent</b>						
Injection drug use	21	18%	21	18%	156	27%
Heterosexual contact	48	42%	42	35%	282	49%
Blood product	1	1%	1	1%	8	1%
Other/unknown	44	39%	56	47%	132	23%
Subtotal	114	100%	120	100%	578	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	3	75%	5	71%	42	79%
Other	1	25%	2	29%	11	21%
Subtotal	4	100%	7	100%	53	100%
<b>Total</b>	<b>496</b>		<b>469</b>		<b>3,740</b>	

*Note:* Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Cleveland Consortium 1

**Table 77. Reported persons living with HIV/AIDS as of Dec. 31, 2003, Ryan White Title II Cleveland Consortium 1, by current disease status and selected characteristics**

	HIV/AIDS			Current Disease Status			
	Rate <sup>a</sup>	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
<b>Gender</b>							
Male	277.0	2,779	77%	1,214	73%	1,565	80%
Female	75.9	832	23%	438	26%	394	20%
Unknown	*	1	<1%	1	<1%	-	-
<b>Age (as of 12/2003)</b>							
<13	8.1	31	1%	16	1%	15	1%
13-19	14.6	29	1%	17	1%	12	1%
20-24	62.7	72	2%	52	3%	20	1%
25-29	122.1	160	4%	106	6%	54	3%
30-39	357.5	1,102	31%	561	34%	541	28%
40-49	456.5	1,501	42%	621	38%	880	45%
50-64	201.2	653	18%	256	15%	397	20%
65+	20.6	64	2%	24	1%	40	2%
<b>Race/Ethnicity</b>							
White, not Hispanic	83.1	1,303	36%	558	34%	745	38%
Black, not Hispanic	433.5	1,824	50%	832	50%	992	51%
Hispanic	496.8	365	10%	158	10%	207	11%
Asian/Pacific Islander	40.0	13	<1%	5	<1%	8	<1%
American Indian/Alaska Native	239.5	11	<1%	5	<1%	6	<1%
Unknown	*	96	3%	95	6%	1	<1%
<b>Race/Gender</b>							
White, not Hispanic Male	149.4	1,129	31%	467	28%	662	34%
White, not Hispanic Female	21.4	174	5%	91	6%	83	4%
Black, not Hispanic Male	687.4	1,324	37%	578	35%	746	38%
Black, not Hispanic Female	218.7	499	14%	253	15%	246	13%
Hispanic Male	634.9	236	7%	92	6%	144	7%
Hispanic Female	355.5	129	4%	66	4%	63	3%
Asian/Pacific Islander Male	63.0	10	<1%	3	<1%	7	<1%
Asian/Pacific Islander Female	*	3	<1%	2	<1%	1	<1%
American Indian/Alaska Native Male	318.6	7	<1%	2	<1%	5	<1%
American Indian/Alaska Native Female	*	4	<1%	3	<1%	1	<1%
Unknown	*	97	3%	96	6%	1	<1%
<b>Total</b>	<b>172.0</b>	<b>3,612</b>		<b>1,653</b>		<b>1,959</b>	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (\*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> The rate is the number of persons living with HIV/AIDS per 100,000 population calculated using 2000 census data.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Cleveland Consortium 1

**Table 78. Reported persons living with HIV/AIDS as of Dec. 31, 2003, Ryan White Title II Cleveland Consortium 1, by current disease status and transmission category**

Transmission category	HIV/AIDS		Current Disease Status			
			HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	1,533	56%	564	47%	969	62%
Injection drug use (IDU)	280	10%	91	8%	189	12%
Male-to-male sexual contact and IDU	161	6%	56	5%	105	7%
Heterosexual contact	111	4%	43	4%	68	4%
Blood product	15	1%	5	<1%	10	1%
Other/unknown	655	24%	443	37%	212	14%
Subtotal	2,755	100%	1,202	100%	1,553	100%
<b>Female adult or adolescent</b>						
Injection drug use	150	19%	53	13%	97	26%
Heterosexual contact	313	39%	142	34%	171	45%
Blood product	6	1%	2	<1%	4	1%
Other/unknown	332	41%	226	53%	106	28%
Subtotal	801	100%	423	100%	378	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	42	76%	20	74%	22	79%
Other	13	24%	7	26%	6	21%
Subtotal	55	100%	27	100%	28	100%
<b>Total</b>	<b>3,612</b>		<b>1,653</b>		<b>1,959</b>	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (\*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Cleveland Consortium 1

**Table 79. Reported cases of HIV infection, Ryan White Title II Cleveland Consortium 1, by year of diagnosis and selected characteristics**

	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
<b>Gender</b>				
Male	530	75%	429	73%
Female	175	25%	162	27%
Unknown	-	-	-	-
<b>Age</b>				
<13	11	2%	8	1%
13-19	19	3%	10	2%
20-24	54	8%	55	9%
25-29	81	11%	77	13%
30-39	287	41%	215	36%
40-49	187	27%	168	28%
50-64	59	8%	57	10%
65+	7	1%	1	<1%
<b>Race/Ethnicity</b>				
White, not Hispanic	241	34%	194	33%
Black, not Hispanic	392	56%	316	53%
Hispanic	56	8%	55	9%
Asian/Pacific Islander	4	1%	2	<1%
American Indian/Alaska Native	4	1%	2	<1%
Unknown	8	1%	22	4%
<b>Race/Gender</b>				
White, not Hispanic Male	201	29%	166	28%
White, not Hispanic Female	40	6%	28	5%
Black, not Hispanic Male	278	39%	208	35%
Black, not Hispanic Female	114	16%	108	18%
Hispanic Male	40	6%	37	6%
Hispanic Female	16	2%	18	3%
Asian/Pacific Islander Male	3	<1%	-	-
Asian/Pacific Islander Female	1	<1%	2	<1%
American Indian/Alaska Native Male	2	<1%	1	<1%
American Indian/Alaska Native Female	2	<1%	1	<1%
Unknown	8	1%	22	4%
<b>Total</b>	<b>705</b>		<b>591</b>	

*Note:* HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Cleveland Consortium 1

**Table 80. Reported cases of HIV infection, Ryan White Title II Cleveland Consortium 1, by year of diagnosis and transmission category**

Transmission category	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
<b>Male adult or adolescent</b>				
Male-to-male sexual contact	276	53%	183	43%
Injection drug use (IDU)	47	9%	23	5%
Male-to-male sexual contact and IDU	15	3%	7	2%
Heterosexual contact	21	4%	14	3%
Blood product	-	-	-	-
Other/unknown	164	31%	199	47%
Subtotal	523	100%	426	100%
<b>Female adult or adolescent</b>				
Injection drug use	14	8%	10	6%
Heterosexual contact	61	36%	32	20%
Blood product	2	1%	1	1%
Other/unknown	94	55%	114	73%
Subtotal	171	100%	157	100%
<b>Child (&lt;13 yrs at diagnosis)</b>				
Perinatal	8	73%	7	88%
Other	3	27%	1	13%
Subtotal	11	-	8	-
<b>Total</b>	<b>705</b>		<b>591</b>	

*Note:* HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Cleveland Consortium 1

**Table 81. Reported cases of AIDS, Ryan White Title II Cleveland Consortium 1, by year of diagnosis and selected characteristics**

	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
<b>Gender</b>						
Male	377	76%	344	74%	3,103	84%
Female	117	24%	124	26%	605	16%
Unknown	-	-	-	-	-	-
<b>Age</b>						
<13	3	1%	5	1%	46	1%
13-19	2	<1%	4	1%	24	1%
20-24	16	3%	22	5%	148	4%
25-29	42	9%	38	8%	495	13%
30-39	222	45%	195	42%	1,680	45%
40-49	163	33%	149	32%	937	25%
50-64	40	8%	53	11%	332	9%
65+	6	1%	2	<1%	46	1%
<b>Race/Ethnicity</b>						
White, not Hispanic	156	32%	157	34%	1,561	42%
Black, not Hispanic	301	61%	264	56%	1,851	50%
Hispanic	30	6%	45	10%	281	8%
Asian/Pacific Islander	3	1%	-	-	8	<1%
American Indian/Alaska Native	4	1%	1	<1%	6	<1%
Unknown	-	-	1	<1%	1	<1%
<b>Race/Gender</b>						
White, not Hispanic Male	136	28%	133	28%	1,442	39%
White, not Hispanic Female	20	4%	24	5%	119	3%
Black, not Hispanic Male	213	43%	182	39%	1,450	39%
Black, not Hispanic Female	88	18%	82	18%	401	11%
Hispanic Male	23	5%	27	6%	199	5%
Hispanic Female	7	1%	18	4%	82	2%
Asian/Pacific Islander Male	2	<1%	-	-	7	<1%
Asian/Pacific Islander Female	1	<1%	-	-	1	<1%
American Indian/Alaska Native Male	3	1%	1	<1%	4	<1%
American Indian/Alaska Native Female	1	<1%	-	-	2	<1%
Unknown	-	-	1	<1%	1	<1%
<b>Total</b>	<b>494</b>		<b>468</b>		<b>3,708</b>	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Cleveland Consortium 1

**Table 82. Reported cases of AIDS, Ryan White Title II Cleveland Consortium 1, by year of diagnosis and transmission category**

Transmission category	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	216	57%	192	56%	2,070	67%
Injection drug use (IDU)	39	10%	30	9%	352	11%
Male-to-male sexual contact and IDU	19	5%	17	5%	237	8%
Heterosexual contact	20	5%	14	4%	109	4%
Blood product	3	1%	1	<1%	42	1%
Other/unknown	79	21%	87	26%	270	9%
Subtotal	376	100%	341	100%	3,080	100%
<b>Female adult or adolescent</b>						
Injection drug use	21	18%	21	18%	156	27%
Heterosexual contact	48	42%	42	35%	279	49%
Blood product	1	1%	1	1%	8	1%
Other/unknown	44	39%	56	47%	132	23%
Subtotal	114	100%	120	100%	575	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	3	75%	5	71%	42	79%
Other	1	25%	2	29%	11	21%
Subtotal	4	100%	7	100%	53	100%
<b>Total</b>	<b>494</b>		<b>468</b>		<b>3,708</b>	

*Note:* Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Columbus Consortium 2

**Table 83. Reported persons living with HIV/AIDS as of Dec. 31, 2003, Ryan White Title II Columbus Consortium 2, by current disease status and selected characteristics**

	Current Disease Status						
	HIV/AIDS			HIV (not AIDS)		AIDS	
	Rate <sup>a</sup>	No.	%	No.	%	No.	%
<b>Gender</b>							
Male	298.1	2,313	83%	1,359	81%	954	85%
Female	60.0	483	17%	311	19%	172	15%
Unknown	*	1	<1%	1	<1%	-	-
<b>Age (as of 12/2003)</b>							
<13	4.7	14	1%	10	1%	4	<1%
13-19	7.8	12	<1%	9	1%	3	<1%
20-24	56.5	68	2%	58	3%	10	1%
25-29	126.0	159	6%	123	7%	36	3%
30-39	371.4	967	35%	605	36%	362	32%
40-49	446.0	1,081	39%	612	37%	469	42%
50-64	201.9	448	16%	228	14%	220	20%
65+	30.3	48	2%	26	2%	22	2%
<b>Race/Ethnicity</b>							
White, not Hispanic	130.0	1,679	60%	938	56%	741	66%
Black, not Hispanic	410.1	887	32%	549	33%	338	30%
Hispanic	288.5	82	3%	48	3%	34	3%
Asian/Pacific Islander	35.0	14	1%	5	<1%	9	1%
American Indian/Alaska Native	182.7	9	<1%	7	<1%	2	<1%
Unknown	*	126	5%	124	7%	2	<1%
<b>Race/Gender</b>							
White, not Hispanic Male	239.5	1,517	54%	836	50%	681	60%
White, not Hispanic Female	24.6	162	6%	102	6%	60	5%
Black, not Hispanic Male	590.5	616	22%	376	23%	240	21%
Black, not Hispanic Female	242.0	271	10%	173	10%	98	9%
Hispanic Male	368.4	58	2%	35	2%	23	2%
Hispanic Female	189.3	24	1%	13	1%	11	1%
Asian/Pacific Islander Male	60.0	12	<1%	4	<1%	8	1%
Asian/Pacific Islander Female	*	2	<1%	1	<1%	1	<1%
American Indian/Alaska Native Male	244.4	6	<1%	6	<1%	-	-
American Indian/Alaska Native Female	*	3	<1%	1	<1%	2	<1%
Unknown	*	126	5%	124	7%	2	<1%
<b>Total</b>	<b>176.9</b>	<b>2,797</b>		<b>1,671</b>		<b>1,126</b>	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (\*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> The rate is the number of persons living with HIV/AIDS per 100,000 population calculated using 2000 census data.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Columbus Consortium 2

**Table 84. Reported persons living with HIV/AIDS as of Dec. 31, 2003, Ryan White Title II Columbus Consortium 2, by current disease status and transmission category**

Transmission category	HIV/AIDS		Current Disease Status			
			HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	1,028	45%	497	37%	531	56%
Injection drug use (IDU)	94	4%	34	3%	60	6%
Male-to-male sexual contact and IDU	60	3%	27	2%	33	3%
Heterosexual contact	53	2%	22	2%	31	3%
Blood product	15	1%	7	1%	8	1%
Other/unknown	1,051	46%	764	57%	287	30%
Subtotal	2,301	100%	1,351	100%	950	100%
<b>Female adult or adolescent</b>						
Injection drug use	46	10%	21	7%	25	15%
Heterosexual contact	166	35%	90	30%	76	45%
Blood product	5	1%	4	1%	1	1%
Other/unknown	255	54%	188	62%	67	40%
Subtotal	472	100%	303	100%	169	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	18	78%	12	75%	6	86%
Other	5	22%	4	25%	1	14%
Subtotal	23	100%	16	100%	7	100%
<b>Total</b>	<b>2,797</b>		<b>1,671</b>		<b>1,126</b>	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (\*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Columbus Consortium 2

**Table 85. Reported cases of HIV infection, Ryan White Title II Columbus Consortium 2, by year of diagnosis and selected characteristics**

	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
<b>Gender</b>				
Male	465	83%	596	82%
Female	97	17%	135	18%
Unknown	1	<1%	-	-
<b>Age</b>				
<13	2	<1%	5	1%
13-19	10	2%	12	2%
20-24	40	7%	60	8%
25-29	86	15%	79	11%
30-39	254	45%	284	39%
40-49	126	22%	213	29%
50-64	41	7%	73	10%
65+	4	1%	5	1%
<b>Race/Ethnicity</b>				
White, not Hispanic	340	60%	385	53%
Black, not Hispanic	177	31%	249	34%
Hispanic	12	2%	26	4%
Asian/Pacific Islander	5	1%	4	1%
American Indian/Alaska Native	2	<1%	2	<1%
Unknown	27	5%	65	9%
<b>Race/Gender</b>				
White, not Hispanic Male	315	56%	347	47%
White, not Hispanic Female	25	4%	38	5%
Black, not Hispanic Male	118	21%	166	23%
Black, not Hispanic Female	59	10%	83	11%
Hispanic Male	6	1%	22	3%
Hispanic Female	6	1%	4	1%
Asian/Pacific Islander Male	4	1%	3	<1%
Asian/Pacific Islander Female	1	<1%	1	<1%
American Indian/Alaska Native Male	1	<1%	2	<1%
American Indian/Alaska Native Female	1	<1%	-	-
Unknown	27	5%	65	9%
<b>Total</b>	<b>563</b>		<b>731</b>	

*Note:* HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Columbus Consortium 2

**Table 86. Reported cases of HIV infection, Ryan White Title II Columbus Consortium 2, by year of diagnosis and transmission category**

Transmission category	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
<b>Male adult or adolescent</b>				
Male-to-male sexual contact	201	43%	138	23%
Injection drug use (IDU)	10	2%	8	1%
Male-to-male sexual contact and IDU	12	3%	3	1%
Heterosexual contact	13	3%	7	1%
Blood product	-	-	1	<1%
Other/unknown	228	49%	437	74%
Subtotal	464	100%	594	100%
<b>Female adult or adolescent</b>				
Injection drug use	8	8%	3	2%
Heterosexual contact	31	32%	19	14%
Blood product	2	2%	1	1%
Other/unknown	55	57%	109	83%
Subtotal	96	100%	132	100%
<b>Child (&lt;13 yrs at diagnosis)</b>				
Perinatal	2	100%	4	80%
Other	-	-	1	20%
Subtotal	2	-	5	-
<b>Total</b>	<b>563</b>		<b>731</b>	

*Note:* HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Columbus Consortium 2

**Table 87. Reported cases of AIDS, Ryan White Title II Columbus Consortium 2, by year of diagnosis and selected characteristics**

	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
<b>Gender</b>						
Male	219	80%	313	84%	2,257	90%
Female	56	20%	61	16%	263	10%
Unknown	-	-	-	-	-	-
<b>Age</b>						
<13	1	<1%	1	<1%	14	1%
13-19	1	<1%	1	<1%	11	<1%
20-24	4	1%	12	3%	99	4%
25-29	25	9%	22	6%	368	15%
30-39	152	55%	154	41%	1,183	47%
40-49	65	24%	130	35%	618	25%
50-64	24	9%	52	14%	213	8%
65+	3	1%	2	1%	14	1%
<b>Race/Ethnicity</b>						
White, not Hispanic	163	59%	226	60%	1,844	73%
Black, not Hispanic	99	36%	133	36%	617	24%
Hispanic	7	3%	11	3%	41	2%
Asian/Pacific Islander	2	1%	2	1%	11	<1%
American Indian/Alaska Native	3	1%	-	-	4	<1%
Unknown	1	<1%	2	1%	3	<1%
<b>Race/Gender</b>						
White, not Hispanic Male	149	54%	202	54%	1,734	69%
White, not Hispanic Female	14	5%	24	6%	110	4%
Black, not Hispanic Male	63	23%	98	26%	476	19%
Black, not Hispanic Female	36	13%	35	9%	141	6%
Hispanic Male	4	1%	9	2%	32	1%
Hispanic Female	3	1%	2	1%	9	<1%
Asian/Pacific Islander Male	1	<1%	2	1%	10	<1%
Asian/Pacific Islander Female	1	<1%	-	-	1	<1%
American Indian/Alaska Native Male	1	<1%	-	-	2	<1%
American Indian/Alaska Native Female	2	1%	-	-	2	<1%
Unknown	1	<1%	2	1%	3	<1%
<b>Total</b>	<b>275</b>		<b>374</b>		<b>2,520</b>	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Columbus Consortium 2

**Table 88. Reported cases of AIDS, Ryan White Title II Columbus Consortium 2, by year of diagnosis and transmission category**

Transmission category	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	117	54%	113	36%	1,619	72%
Injection drug use (IDU)	12	6%	11	4%	128	6%
Male-to-male sexual contact and IDU	8	4%	9	3%	81	4%
Heterosexual contact	8	4%	9	3%	55	2%
Blood product	-	-	1	<1%	28	1%
Other/unknown	73	33%	170	54%	337	15%
Subtotal	218	100%	313	100%	2,248	100%
<b>Female adult or adolescent</b>						
Injection drug use	11	20%	5	8%	44	17%
Heterosexual contact	24	43%	16	27%	121	47%
Blood product	2	4%	-	-	9	4%
Other/unknown	19	34%	39	65%	81	32%
Subtotal	56	100%	60	100%	255	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	1	100%	1	100%	13	76%
Other	-	-	-	-	4	24%
Subtotal	1	100%	1	100%	17	100%
<b>Total</b>	<b>275</b>		<b>374</b>		<b>2,520</b>	

*Note:* Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Cincinnati Consortium 3

**Table 89. Reported persons living with HIV/AIDS as of Dec. 31, 2003, Ryan White Title II Cincinnati Consortium 3, by current disease status and selected characteristics**

	Current Disease Status						
	HIV/AIDS			HIV (not AIDS)		AIDS	
	Rate <sup>a</sup>	No.	%	No.	%	No.	%
<b>Gender</b>							
Male	221.3	1,786	82%	923	78%	863	87%
Female	46.0	395	18%	263	22%	132	13%
Unknown	*	-	-	-	-	-	-
<b>Age (as of 12/2003)</b>							
<13	4.1	13	1%	10	1%	3	<1%
13-19	4.1	7	<1%	3	<1%	4	<1%
20-24	54.9	61	3%	54	5%	7	1%
25-29	126.5	140	6%	103	9%	37	4%
30-39	268.6	686	31%	400	34%	286	29%
40-49	341.6	884	41%	437	37%	447	45%
50-64	145.2	351	16%	162	14%	189	19%
65+	19.5	39	2%	17	1%	22	2%
<b>Race/Ethnicity</b>							
White, not Hispanic	79.8	1,110	51%	577	49%	533	54%
Black, not Hispanic	424.8	970	44%	537	45%	433	44%
Hispanic	180.3	33	2%	17	1%	16	2%
Asian/Pacific Islander	41.1	10	<1%	4	<1%	6	1%
American Indian/Alaska Native	*	4	<1%	2	<1%	2	<1%
Unknown	*	54	2%	49	4%	5	1%
<b>Race/Gender</b>							
White, not Hispanic Male	144.5	978	45%	488	41%	490	49%
White, not Hispanic Female	18.5	132	6%	89	8%	43	4%
Black, not Hispanic Male	681.6	725	33%	377	32%	348	35%
Black, not Hispanic Female	200.8	245	11%	160	13%	85	9%
Hispanic Male	264.3	26	1%	12	1%	14	1%
Hispanic Female	82.7	7	<1%	5	<1%	2	<1%
Asian/Pacific Islander Male	74.8	9	<1%	4	<1%	5	1%
Asian/Pacific Islander Female	*	1	<1%	-	-	1	<1%
American Indian/Alaska Native Male	*	4	<1%	2	<1%	2	<1%
American Indian/Alaska Native Female	*	-	-	-	-	-	-
Unknown	*	54	2%	49	4%	5	1%
<b>Total</b>	<b>131.0</b>	<b>2,181</b>		<b>1,186</b>		<b>995</b>	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (\*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> The rate is the number of persons living with HIV/AIDS per 100,000 population calculated using 2000 census data.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Cincinnati Consortium 3

**Table 90. Reported persons living with HIV/AIDS as of Dec. 31, 2003, Ryan White Title II Cincinnati Consortium 3, by current disease status and transmission category**

Transmission category	HIV/AIDS		Current Disease Status			
			HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	944	53%	394	43%	550	64%
Injection drug use (IDU)	105	6%	39	4%	66	8%
Male-to-male sexual contact and IDU	82	5%	29	3%	53	6%
Heterosexual contact	42	2%	15	2%	27	3%
Blood product	15	1%	7	1%	8	1%
Other/unknown	585	33%	433	47%	152	18%
Subtotal	1,773	100%	917	100%	856	100%
<b>Female adult or adolescent</b>						
Injection drug use	42	11%	19	7%	23	18%
Heterosexual contact	133	34%	77	30%	56	43%
Blood product	3	1%	2	1%	1	1%
Other/unknown	208	54%	159	62%	49	38%
Subtotal	386	100%	257	100%	129	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	15	68%	10	83%	5	50%
Other	7	32%	2	17%	5	50%
Subtotal	22	100%	12	100%	10	100%
<b>Total</b>	<b>2,181</b>		<b>1,186</b>		<b>995</b>	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (\*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Cincinnati Consortium 3

**Table 91. Reported cases of HIV infection, Ryan White Title II Cincinnati Consortium 3, by year of diagnosis and selected characteristics**

	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
<b>Gender</b>				
Male	335	80%	338	78%
Female	85	20%	94	22%
Unknown	-	-	-	-
<b>Age</b>				
<13	3	1%	-	-
13-19	14	3%	13	3%
20-24	41	10%	45	10%
25-29	63	15%	54	13%
30-39	172	41%	143	33%
40-49	88	21%	129	30%
50-64	35	8%	41	9%
65+	4	1%	7	2%
<b>Race/Ethnicity</b>				
White, not Hispanic	203	48%	194	45%
Black, not Hispanic	191	45%	206	48%
Hispanic	7	2%	9	2%
Asian/Pacific Islander	1	<1%	-	-
American Indian/Alaska Native	-	-	1	<1%
Unknown	18	4%	22	5%
<b>Race/Gender</b>				
White, not Hispanic Male	173	41%	165	38%
White, not Hispanic Female	30	7%	29	7%
Black, not Hispanic Male	141	34%	150	35%
Black, not Hispanic Female	50	12%	56	13%
Hispanic Male	5	1%	6	1%
Hispanic Female	2	<1%	3	1%
Asian/Pacific Islander Male	1	<1%	-	-
Asian/Pacific Islander Female	-	-	-	-
American Indian/Alaska Native Male	-	-	1	<1%
American Indian/Alaska Native Female	-	-	-	-
Unknown	18	4%	22	5%
<b>Total</b>	<b>420</b>		<b>432</b>	

*Note:* HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Cincinnati Consortium 3

**Table 92. Reported cases of HIV infection, Ryan White Title II Cincinnati Consortium 3, by year of diagnosis and transmission category**

Transmission category	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
<b>Male adult or adolescent</b>				
Male-to-male sexual contact	144	43%	124	37%
Injection drug use (IDU)	13	4%	14	4%
Male-to-male sexual contact and IDU	5	1%	4	1%
Heterosexual contact	3	1%	14	4%
Blood product	1	<1%	2	1%
Other/unknown	168	50%	180	53%
Subtotal	334	100%	338	100%
<b>Female adult or adolescent</b>				
Injection drug use	3	4%	4	4%
Heterosexual contact	27	33%	20	21%
Blood product	-	-	-	-
Other/unknown	53	64%	70	74%
Subtotal	83	100%	94	100%
<b>Child (&lt;13 yrs at diagnosis)</b>				
Perinatal	1	33%	-	-
Other	2	67%	-	-
Subtotal	3	-	-	-
<b>Total</b>	<b>420</b>		<b>432</b>	

*Note:* HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Cincinnati Consortium 3

**Table 93. Reported cases of AIDS, Ryan White Title II Cincinnati Consortium 3, by year of diagnosis and selected characteristics**

	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
<b>Gender</b>						
Male	194	87%	176	84%	1,844	90%
Female	30	13%	33	16%	194	10%
Unknown	-	-	-	-	-	-
<b>Age</b>						
<13	1	<1%	-	-	15	1%
13-19	1	<1%	4	2%	13	1%
20-24	6	3%	9	4%	70	3%
25-29	32	14%	16	8%	310	15%
30-39	105	47%	79	38%	954	47%
40-49	57	25%	73	35%	482	24%
50-64	20	9%	23	11%	170	8%
65+	2	1%	5	2%	24	1%
<b>Race/Ethnicity</b>						
White, not Hispanic	112	50%	95	45%	1,231	60%
Black, not Hispanic	106	47%	106	51%	775	38%
Hispanic	2	1%	4	2%	16	1%
Asian/Pacific Islander	2	1%	1	<1%	8	<1%
American Indian/Alaska Native	-	-	-	-	2	<1%
Unknown	2	1%	3	1%	6	<1%
<b>Race/Gender</b>						
White, not Hispanic Male	103	46%	81	39%	1,160	57%
White, not Hispanic Female	9	4%	14	7%	71	3%
Black, not Hispanic Male	86	38%	88	42%	656	32%
Black, not Hispanic Female	20	9%	18	9%	119	6%
Hispanic Male	2	1%	4	2%	14	1%
Hispanic Female	-	-	-	-	2	<1%
Asian/Pacific Islander Male	1	<1%	1	<1%	7	<1%
Asian/Pacific Islander Female	1	<1%	-	-	1	<1%
American Indian/Alaska Native Male	-	-	-	-	2	<1%
American Indian/Alaska Native Female	-	-	-	-	-	-
Unknown	2	1%	3	1%	6	<1%
<b>Total</b>	<b>224</b>		<b>209</b>		<b>2,038</b>	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Cincinnati Consortium 3

**Table 94. Reported cases of AIDS, Ryan White Title II Cincinnati Consortium 3, by year of diagnosis and transmission category**

Transmission category	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	117	60%	83	47%	1,319	72%
Injection drug use (IDU)	10	5%	13	7%	97	5%
Male-to-male sexual contact and IDU	7	4%	6	3%	116	6%
Heterosexual contact	1	1%	6	3%	39	2%
Blood product	1	1%	-	-	40	2%
Other/unknown	58	30%	67	38%	220	12%
Subtotal	194	100%	175	100%	1,831	100%
<b>Female adult or adolescent</b>						
Injection drug use	3	10%	3	9%	41	22%
Heterosexual contact	11	38%	8	25%	80	43%
Blood product	-	-	-	-	4	2%
Other/unknown	15	52%	21	66%	63	34%
Subtotal	29	100%	32	100%	188	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	-	-	1	50%	13	68%
Other	1	100%	1	50%	6	32%
Subtotal	1	100%	2	100%	19	100%
<b>Total</b>	<b>224</b>		<b>209</b>		<b>2,038</b>	

*Note:* Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Dayton Consortium 4

**Table 95. Reported persons living with HIV/AIDS as of Dec. 31, 2003, Ryan White Title II Dayton Consortium 4, by current disease status and selected characteristics**

	Current Disease Status						
	HIV/AIDS			HIV (not AIDS)		AIDS	
	Rate <sup>a</sup>	No.	%	No.	%	No.	%
<b>Gender</b>							
Male	206.5	1,044	80%	484	76%	560	83%
Female	48.7	263	20%	152	24%	111	17%
Unknown	*	-	-	-	-	-	-
<b>Age (as of 12/2003)</b>							
<13	3.2	6	<1%	3	<1%	3	<1%
13-19	13.1	14	1%	8	1%	6	1%
20-24	58.7	41	3%	33	5%	8	1%
25-29	139.1	91	7%	62	10%	29	4%
30-39	269.8	401	31%	197	31%	204	30%
40-49	308.1	487	37%	226	36%	261	39%
50-64	147.8	250	19%	101	16%	149	22%
65+	12.0	17	1%	6	1%	11	2%
<b>Race/Ethnicity</b>							
White, not Hispanic	82.4	723	55%	337	53%	386	58%
Black, not Hispanic	390.8	547	42%	274	43%	273	41%
Hispanic	142.1	17	1%	8	1%	9	1%
Asian/Pacific Islander	36.9	5	<1%	3	<1%	2	<1%
American Indian/Alaska Native	*	1	<1%	1	<1%	-	-
Unknown	*	14	1%	13	2%	1	<1%
<b>Race/Gender</b>							
White, not Hispanic Male	145.2	619	47%	279	44%	340	51%
White, not Hispanic Female	23.0	104	8%	58	9%	46	7%
Black, not Hispanic Male	608.6	398	30%	189	30%	209	31%
Black, not Hispanic Female	199.8	149	11%	85	13%	64	10%
Hispanic Male	212.2	13	1%	4	1%	9	1%
Hispanic Female	*	4	<1%	4	1%	-	-
Asian/Pacific Islander Male	*	4	<1%	3	<1%	1	<1%
Asian/Pacific Islander Female	*	1	<1%	-	-	1	<1%
American Indian/Alaska Native Male	*	-	-	-	-	-	-
American Indian/Alaska Native Female	*	1	<1%	1	<1%	-	-
Unknown	*	14	1%	13	2%	1	<1%
<b>Total</b>	<b>124.9</b>	<b>1,307</b>		<b>636</b>		<b>671</b>	

*Note:* Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (\*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> The rate is the number of persons living with HIV/AIDS per 100,000 population calculated using 2000 census data.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Dayton Consortium 4

**Table 96. Reported persons living with HIV/AIDS as of Dec. 31, 2003, Ryan White Title II Dayton Consortium 4, by current disease status and transmission category**

Transmission category	HIV/AIDS		Current Disease Status			
			HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	644	62%	280	58%	364	65%
Injection drug use (IDU)	48	5%	16	3%	32	6%
Male-to-male sexual contact and IDU	60	6%	17	4%	43	8%
Heterosexual contact	40	4%	13	3%	27	5%
Blood product	15	1%	5	1%	10	2%
Other/unknown	231	22%	151	31%	80	14%
Subtotal	1,038	100%	482	100%	556	100%
<b>Female adult or adolescent</b>						
Injection drug use	32	13%	13	9%	19	18%
Heterosexual contact	116	45%	71	48%	45	42%
Blood product	4	2%	-	-	4	4%
Other/unknown	103	40%	65	44%	38	36%
Subtotal	255	100%	149	100%	106	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	10	71%	3	60%	7	78%
Other	4	29%	2	40%	2	22%
Subtotal	14	100%	5	100%	9	100%
<b>Total</b>	<b>1,307</b>		<b>636</b>		<b>671</b>	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (\*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Dayton Consortium 4

**Table 97. Reported cases of HIV infection, Ryan White Title II Dayton Consortium 4, by year of diagnosis and selected characteristics**

	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
<b>Gender</b>				
Male	230	78%	195	73%
Female	63	22%	73	27%
Unknown	-	-	-	-
<b>Age</b>				
<13	-	-	1	<1%
13-19	7	2%	20	7%
20-24	29	10%	20	7%
25-29	41	14%	37	14%
30-39	110	38%	99	37%
40-49	78	27%	69	26%
50-64	23	8%	20	7%
65+	5	2%	2	1%
<b>Race/Ethnicity</b>				
White, not Hispanic	150	51%	127	47%
Black, not Hispanic	134	46%	132	49%
Hispanic	4	1%	2	1%
Asian/Pacific Islander	1	<1%	3	1%
American Indian/Alaska Native	1	<1%	-	-
Unknown	3	1%	4	1%
<b>Race/Gender</b>				
White, not Hispanic Male	127	43%	106	40%
White, not Hispanic Female	23	8%	21	8%
Black, not Hispanic Male	98	33%	82	31%
Black, not Hispanic Female	36	12%	50	19%
Hispanic Male	3	1%	1	<1%
Hispanic Female	1	<1%	1	<1%
Asian/Pacific Islander Male	1	<1%	3	1%
Asian/Pacific Islander Female	-	-	-	-
American Indian/Alaska Native Male	-	-	-	-
American Indian/Alaska Native Female	1	<1%	-	-
Unknown	3	1%	4	1%
<b>Total</b>	<b>293</b>		<b>268</b>	

*Note:* HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Dayton Consortium 4

**Table 98. Reported cases of HIV infection, Ryan White Title II Dayton Consortium 4, by year of diagnosis and transmission category**

Transmission category	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
<b>Male adult or adolescent</b>				
Male-to-male sexual contact	153	67%	84	43%
Injection drug use (IDU)	7	3%	4	2%
Male-to-male sexual contact and IDU	6	3%	5	3%
Heterosexual contact	12	5%	7	4%
Blood product	-	-	1	1%
Other/unknown	52	23%	94	48%
Subtotal	230	100%	195	100%
<b>Female adult or adolescent</b>				
Injection drug use	2	3%	5	7%
Heterosexual contact	23	37%	21	29%
Blood product	2	3%	1	1%
Other/unknown	36	57%	45	63%
Subtotal	63	100%	72	100%
<b>Child (&lt;13 yrs at diagnosis)</b>				
Perinatal	-	-	1	100%
Other	-	-	-	-
Subtotal	-	-	1	-
<b>Total</b>	<b>293</b>		<b>268</b>	

*Note:* HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Dayton Consortium 4

**Table 99. Reported cases of AIDS, Ryan White Title II Dayton Consortium 4, by year of diagnosis and selected characteristics**

	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
<b>Gender</b>						
Male	147	87%	137	75%	1,076	87%
Female	22	13%	45	25%	161	13%
Unknown	-	-	-	-	-	-
<b>Age</b>						
<13	-	-	-	-	16	1%
13-19	2	1%	3	2%	12	1%
20-24	4	2%	7	4%	47	4%
25-29	16	9%	18	10%	161	13%
30-39	79	47%	79	43%	539	44%
40-49	48	28%	51	28%	342	28%
50-64	16	9%	21	12%	102	8%
65+	4	2%	3	2%	18	1%
<b>Race/Ethnicity</b>						
White, not Hispanic	96	57%	96	53%	816	66%
Black, not Hispanic	70	41%	84	46%	405	33%
Hispanic	3	2%	1	1%	13	1%
Asian/Pacific Islander	-	-	1	1%	3	<1%
American Indian/Alaska Native	-	-	-	-	-	-
Unknown	-	-	-	-	-	-
<b>Race/Gender</b>						
White, not Hispanic Male	88	52%	80	44%	740	60%
White, not Hispanic Female	8	5%	16	9%	76	6%
Black, not Hispanic Male	56	33%	55	30%	321	26%
Black, not Hispanic Female	14	8%	29	16%	84	7%
Hispanic Male	3	2%	1	1%	13	1%
Hispanic Female	-	-	-	-	-	-
Asian/Pacific Islander Male	-	-	1	1%	2	<1%
Asian/Pacific Islander Female	-	-	-	-	1	<1%
American Indian/Alaska Native Male	-	-	-	-	-	-
American Indian/Alaska Native Female	-	-	-	-	-	-
Unknown	-	-	-	-	-	-
<b>Total</b>	<b>169</b>		<b>182</b>		<b>1,237</b>	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Dayton Consortium 4

**Table 100. Reported cases of AIDS, Ryan White Title II Dayton Consortium 4, by year of diagnosis and transmission category**

Transmission category	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	99	67%	73	54%	742	70%
Injection drug use (IDU)	6	4%	3	2%	53	5%
Male-to-male sexual contact and IDU	7	5%	6	4%	76	7%
Heterosexual contact	13	9%	2	1%	44	4%
Blood product	2	1%	-	-	29	3%
Other/unknown	20	14%	52	38%	121	11%
Subtotal	147	100%	136	100%	1,065	100%
<b>Female adult or adolescent</b>						
Injection drug use	4	18%	6	14%	34	22%
Heterosexual contact	6	27%	13	30%	60	39%
Blood product	2	9%	1	2%	9	6%
Other/unknown	10	45%	24	55%	49	32%
Subtotal	22	100%	44	100%	152	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	-	-	1	50%	14	70%
Other	-	-	1	50%	6	30%
Subtotal	-	-	2	100%	20	100%
<b>Total</b>	<b>169</b>		<b>182</b>		<b>1,237</b>	

*Note:* Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Toledo Consortium 5

**Table 101. Reported persons living with HIV/AIDS as of Dec. 31, 2003, Ryan White Title II Toledo Consortium 5, by current disease status and selected characteristics**

	Current Disease Status						
	HIV/AIDS			HIV (not AIDS)		AIDS	
	Rate <sup>a</sup>	No.	%	No.	%	No.	%
<b>Gender</b>							
Male	180.2	724	77%	363	73%	361	82%
Female	49.9	213	23%	135	27%	78	18%
Unknown	*	-	-	-	-	-	-
<b>Age (as of 12/2003)</b>							
<13	5.2	8	1%	4	1%	4	1%
13-19	5.6	5	1%	4	1%	1	<1%
20-24	42.3	25	3%	20	4%	5	1%
25-29	134.2	73	8%	56	11%	17	4%
30-39	268.9	310	33%	179	36%	131	30%
40-49	280.5	356	38%	167	34%	189	43%
50-64	116.7	144	15%	63	13%	81	18%
65+	14.8	16	2%	5	1%	11	3%
<b>Race/Ethnicity</b>							
White, not Hispanic	76.2	530	57%	283	57%	247	56%
Black, not Hispanic	398.7	338	36%	173	35%	165	38%
Hispanic	132.7	51	5%	25	5%	26	6%
Asian/Pacific Islander	70.3	6	1%	5	1%	1	<1%
American Indian/Alaska Native	*	-	-	-	-	-	-
Unknown	*	12	1%	12	2%	-	-
<b>Race/Gender</b>							
White, not Hispanic Male	131.9	445	47%	227	46%	218	50%
White, not Hispanic Female	23.8	85	9%	56	11%	29	7%
Black, not Hispanic Male	574.9	229	24%	107	21%	122	28%
Black, not Hispanic Female	242.6	109	12%	66	13%	43	10%
Hispanic Male	185.9	36	4%	16	3%	20	5%
Hispanic Female	78.6	15	2%	9	2%	6	1%
Asian/Pacific Islander Male	145.2	6	1%	5	1%	1	<1%
Asian/Pacific Islander Female	*	-	-	-	-	-	-
American Indian/Alaska Native Male	*	-	-	-	-	-	-
American Indian/Alaska Native Female	*	-	-	-	-	-	-
Unknown	*	12	1%	12	2%	-	-
<b>Total</b>	<b>113.0</b>	<b>937</b>		<b>498</b>		<b>439</b>	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (\*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> The rate is the number of persons living with HIV/AIDS per 100,000 population calculated using 2000 census data.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Toledo Consortium 5

**Table 102. Reported persons living with HIV/AIDS as of Dec. 31, 2003, Ryan White Title II Toledo Consortium 5, by current disease status and transmission category**

Transmission category	HIV/AIDS		Current Disease Status			
			HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	429	60%	207	57%	222	62%
Injection drug use (IDU)	51	7%	30	8%	21	6%
Male-to-male sexual contact and IDU	42	6%	13	4%	29	8%
Heterosexual contact	38	5%	18	5%	20	6%
Blood product	10	1%	3	1%	7	2%
Other/unknown	149	21%	90	25%	59	16%
Subtotal	719	100%	361	100%	358	100%
<b>Female adult or adolescent</b>						
Injection drug use	28	14%	13	10%	15	20%
Heterosexual contact	90	44%	60	46%	30	40%
Blood product	1	<1%	-	-	1	1%
Other/unknown	87	42%	58	44%	29	39%
Subtotal	206	100%	131	100%	75	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	11	92%	6	100%	5	83%
Other	1	8%	-	-	1	17%
Subtotal	12	100%	6	100%	6	100%
<b>Total</b>	<b>937</b>		<b>498</b>		<b>439</b>	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (\*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Toledo Consortium 5

**Table 103. Reported cases of HIV infection, Ryan White Title II Toledo Consortium 5, by year of diagnosis and selected characteristics**

	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
<b>Gender</b>				
Male	121	69%	157	77%
Female	54	31%	48	23%
Unknown	-	-	-	-
<b>Age</b>				
<13	3	2%	2	1%
13-19	4	2%	1	<1%
20-24	19	11%	18	9%
25-29	29	17%	29	14%
30-39	76	43%	72	35%
40-49	30	17%	59	29%
50-64	12	7%	22	11%
65+	2	1%	2	1%
<b>Race/Ethnicity</b>				
White, not Hispanic	96	55%	101	49%
Black, not Hispanic	62	35%	88	43%
Hispanic	14	8%	10	5%
Asian/Pacific Islander	1	1%	3	1%
American Indian/Alaska Native	-	-	-	-
Unknown	2	1%	3	1%
<b>Race/Gender</b>				
White, not Hispanic Male	74	42%	87	42%
White, not Hispanic Female	22	13%	14	7%
Black, not Hispanic Male	35	20%	56	27%
Black, not Hispanic Female	27	15%	32	16%
Hispanic Male	9	5%	10	5%
Hispanic Female	5	3%	-	-
Asian/Pacific Islander Male	1	1%	3	1%
Asian/Pacific Islander Female	-	-	-	-
American Indian/Alaska Native Male	-	-	-	-
American Indian/Alaska Native Female	-	-	-	-
Unknown	2	1%	3	1%
<b>Total</b>	<b>175</b>		<b>205</b>	

*Note:* HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Toledo Consortium 5

**Table 104. Reported cases of HIV infection, Ryan White Title II Toledo Consortium 5, by year of diagnosis and transmission category**

Transmission category	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
<b>Male adult or adolescent</b>				
Male-to-male sexual contact	68	56%	73	47%
Injection drug use (IDU)	5	4%	10	6%
Male-to-male sexual contact and IDU	4	3%	5	3%
Heterosexual contact	11	9%	10	6%
Blood product	-	-	-	-
Other/unknown	33	27%	58	37%
Subtotal	121	100%	156	100%
<b>Female adult or adolescent</b>				
Injection drug use	5	10%	1	2%
Heterosexual contact	19	37%	16	34%
Blood product	-	-	-	-
Other/unknown	27	53%	30	64%
Subtotal	51	100%	47	100%
<b>Child (&lt;13 yrs at diagnosis)</b>				
Perinatal	3	100%	2	100%
Other	-	-	-	-
Subtotal	3	-	2	-
<b>Total</b>	<b>175</b>		<b>205</b>	

*Note:* HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Toledo Consortium 5

**Table 105. Reported cases of AIDS, Ryan White Title II Toledo Consortium 5, by year of diagnosis and selected characteristics**

	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
<b>Gender</b>						
Male	80	74%	96	79%	655	83%
Female	28	26%	25	21%	133	17%
Unknown	-	-	-	-	-	-
<b>Age</b>						
<13	1	1%	1	1%	13	2%
13-19	1	1%	-	-	8	1%
20-24	5	5%	4	3%	53	7%
25-29	9	8%	11	9%	108	14%
30-39	49	45%	53	44%	338	43%
40-49	32	30%	37	31%	191	24%
50-64	9	8%	14	12%	68	9%
65+	2	2%	1	1%	9	1%
<b>Race/Ethnicity</b>						
White, not Hispanic	63	58%	58	48%	504	64%
Black, not Hispanic	36	33%	57	47%	252	32%
Hispanic	9	8%	6	5%	30	4%
Asian/Pacific Islander	-	-	-	-	1	<1%
American Indian/Alaska Native	-	-	-	-	1	<1%
Unknown	-	-	-	-	-	-
<b>Race/Gender</b>						
White, not Hispanic Male	46	43%	51	42%	444	56%
White, not Hispanic Female	17	16%	7	6%	60	8%
Black, not Hispanic Male	28	26%	39	32%	186	24%
Black, not Hispanic Female	8	7%	18	15%	66	8%
Hispanic Male	6	6%	6	5%	23	3%
Hispanic Female	3	3%	-	-	7	1%
Asian/Pacific Islander Male	-	-	-	-	1	<1%
Asian/Pacific Islander Female	-	-	-	-	-	-
American Indian/Alaska Native Male	-	-	-	-	1	<1%
American Indian/Alaska Native Female	-	-	-	-	-	-
Unknown	-	-	-	-	-	-
<b>Total</b>	<b>108</b>		<b>121</b>		<b>788</b>	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Toledo Consortium 5

**Table 106. Reported cases of AIDS, Ryan White Title II Toledo Consortium 5, by year of diagnosis and transmission category**

Transmission category	Year of AIDS Diagnosis				Cumulative	
	1998 - 2000		2001 - 2003			
	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	47	59%	47	49%	458	71%
Injection drug use (IDU)	5	6%	7	7%	35	5%
Male-to-male sexual contact and IDU	8	10%	4	4%	43	7%
Heterosexual contact	9	11%	4	4%	26	4%
Blood product	-	-	-	-	12	2%
Other/unknown	11	14%	33	35%	72	11%
Subtotal	80	100%	95	100%	646	100%
<b>Female adult or adolescent</b>						
Injection drug use	5	19%	1	4%	24	19%
Heterosexual contact	11	41%	8	32%	60	47%
Blood product	-	-	-	-	4	3%
Other/unknown	11	41%	16	64%	39	31%
Subtotal	27	100%	25	100%	127	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	1	100%	1	100%	12	80%
Other	-	-	-	-	3	20%
Subtotal	1	100%	1	100%	15	100%
<b>Total</b>	<b>108</b>		<b>121</b>		<b>788</b>	

*Note:* Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Akron Consortium 6

**Table 107. Reported persons living with HIV/AIDS as of Dec. 31, 2003, Ryan White Title II Akron Consortium 6, by current disease status and selected characteristics**

	HIV/AIDS			Current Disease Status			
	Rate <sup>a</sup>	No.	%	HIV (not AIDS)		AIDS	
				No.	%	No.	%
<b>Gender</b>							
Male	130.9	537	81%	239	78%	298	84%
Female	29.1	127	19%	69	22%	58	16%
Unknown	*	-	-	-	-	-	-
<b>Age (as of 12/2003)</b>							
<13	*	4	1%	3	1%	1	<1%
13-19	*	3	<1%	2	1%	1	<1%
20-24	22.8	12	2%	11	4%	1	<1%
25-29	78.7	41	6%	25	8%	16	4%
30-39	150.5	192	29%	97	31%	95	27%
40-49	192.1	260	39%	113	37%	147	41%
50-64	106.4	140	21%	54	18%	86	24%
65+	11.0	12	2%	3	1%	9	3%
<b>Race/Ethnicity</b>							
White, not Hispanic	52.3	390	59%	160	52%	230	65%
Black, not Hispanic	305.2	246	37%	126	41%	120	34%
Hispanic	165.0	12	2%	7	2%	5	1%
Asian/Pacific Islander	*	4	1%	3	1%	1	<1%
American Indian/Alaska Native	*	1	<1%	1	<1%	-	-
Unknown	*	11	2%	11	4%	-	-
<b>Race/Gender</b>							
White, not Hispanic Male	94.3	342	52%	139	45%	203	57%
White, not Hispanic Female	12.5	48	7%	21	7%	27	8%
Black, not Hispanic Male	456.5	171	26%	81	26%	90	25%
Black, not Hispanic Female	173.8	75	11%	45	15%	30	8%
Hispanic Male	272.6	10	2%	6	2%	4	1%
Hispanic Female	*	2	<1%	1	<1%	1	<1%
Asian/Pacific Islander Male	*	3	<1%	2	1%	1	<1%
Asian/Pacific Islander Female	*	1	<1%	1	<1%	-	-
American Indian/Alaska Native Male	*	1	<1%	1	<1%	-	-
American Indian/Alaska Native Female	*	-	-	-	-	-	-
Unknown	*	11	2%	11	4%	-	-
<b>Total</b>	<b>78.5</b>	<b>664</b>		<b>308</b>		<b>356</b>	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (\*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> The rate is the number of persons living with HIV/AIDS per 100,000 population calculated using 2000 census data.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Akron Consortium 6

**Table 108. Reported persons living with HIV/AIDS as of Dec. 31, 2003, Ryan White Title II Akron Consortium 6, by current disease status and transmission category**

Transmission category	HIV/AIDS		Current Disease Status			
			HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	257	48%	70	30%	187	63%
Injection drug use (IDU)	29	5%	9	4%	20	7%
Male-to-male sexual contact and IDU	26	5%	7	3%	19	6%
Heterosexual contact	12	2%	4	2%	8	3%
Blood product	3	1%	1	<1%	2	1%
Other/unknown	206	39%	146	62%	60	20%
Subtotal	533	100%	237	100%	296	100%
<b>Female adult or adolescent</b>						
Injection drug use	13	10%	3	5%	10	17%
Heterosexual contact	43	35%	15	23%	28	48%
Blood product	1	1%	1	2%	-	-
Other/unknown	67	54%	47	71%	20	34%
Subtotal	124	100%	66	100%	58	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	7	100%	5	100%	2	100%
Other	-	-	-	-	-	-
Subtotal	7	100%	5	100%	2	100%
<b>Total</b>	<b>664</b>		<b>308</b>		<b>356</b>	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (\*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Akron Consortium 6

**Table 109. Reported cases of HIV infection, Ryan White Title II Akron Consortium 6, by year of diagnosis and selected characteristics**

	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
<b>Gender</b>				
Male	121	82%	119	78%
Female	26	18%	34	22%
Unknown	-	-	-	-
<b>Age</b>				
<13	2	1%	2	1%
13-19	3	2%	1	1%
20-24	12	8%	19	12%
25-29	14	10%	13	8%
30-39	58	39%	62	41%
40-49	43	29%	33	22%
50-64	14	10%	21	14%
65+	1	1%	2	1%
<b>Race/Ethnicity</b>				
White, not Hispanic	79	54%	86	56%
Black, not Hispanic	62	42%	55	36%
Hispanic	4	3%	4	3%
Asian/Pacific Islander	1	1%	3	2%
American Indian/Alaska Native	-	-	1	1%
Unknown	1	1%	4	3%
<b>Race/Gender</b>				
White, not Hispanic Male	70	48%	73	48%
White, not Hispanic Female	9	6%	13	8%
Black, not Hispanic Male	45	31%	36	24%
Black, not Hispanic Female	17	12%	19	12%
Hispanic Male	4	3%	3	2%
Hispanic Female	-	-	1	1%
Asian/Pacific Islander Male	1	1%	2	1%
Asian/Pacific Islander Female	-	-	1	1%
American Indian/Alaska Native Male	-	-	1	1%
American Indian/Alaska Native Female	-	-	-	-
Unknown	1	1%	4	3%
<b>Total</b>	<b>147</b>		<b>153</b>	

*Note:* HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Akron Consortium 6

**Table 110. Reported cases of HIV infection, Ryan White Title II Akron Consortium 6, by year of diagnosis and transmission category**

Transmission category	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
<b>Male adult or adolescent</b>				
Male-to-male sexual contact	59	49%	34	29%
Injection drug use (IDU)	5	4%	4	3%
Male-to-male sexual contact and IDU	2	2%	1	1%
Heterosexual contact	1	1%	4	3%
Blood product	-	-	-	-
Other/unknown	53	44%	75	64%
Subtotal	120	100%	118	100%
<b>Female adult or adolescent</b>				
Injection drug use	-	-	2	6%
Heterosexual contact	10	40%	5	15%
Blood product	-	-	-	-
Other/unknown	15	60%	26	79%
Subtotal	25	100%	33	100%
<b>Child (&lt;13 yrs at diagnosis)</b>				
Perinatal	2	100%	2	100%
Other	-	-	-	-
Subtotal	2	-	2	-
<b>Total</b>	<b>147</b>		<b>153</b>	

*Note:* HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Akron Consortium 6

**Table 111. Reported cases of AIDS, Ryan White Title II Akron Consortium 6, by year of diagnosis and selected characteristics**

	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
<b>Gender</b>						
Male	89	84%	50	76%	599	88%
Female	17	16%	16	24%	78	12%
Unknown	-	-	-	-	-	-
<b>Age</b>						
<13	1	1%	-	-	1	<1%
13-19	1	1%	-	-	2	<1%
20-24	2	2%	3	5%	29	4%
25-29	14	13%	7	11%	96	14%
30-39	45	42%	27	41%	297	44%
40-49	34	32%	21	32%	185	27%
50-64	8	8%	7	11%	62	9%
65+	1	1%	1	2%	5	1%
<b>Race/Ethnicity</b>						
White, not Hispanic	64	60%	41	62%	443	65%
Black, not Hispanic	39	37%	24	36%	224	33%
Hispanic	2	2%	1	2%	9	1%
Asian/Pacific Islander	1	1%	-	-	1	<1%
American Indian/Alaska Native	-	-	-	-	-	-
Unknown	-	-	-	-	-	-
<b>Race/Gender</b>						
White, not Hispanic Male	55	52%	33	50%	408	60%
White, not Hispanic Female	9	8%	8	12%	35	5%
Black, not Hispanic Male	31	29%	16	24%	183	27%
Black, not Hispanic Female	8	8%	8	12%	41	6%
Hispanic Male	2	2%	1	2%	7	1%
Hispanic Female	-	-	-	-	2	<1%
Asian/Pacific Islander Male	1	1%	-	-	1	<1%
Asian/Pacific Islander Female	-	-	-	-	-	-
American Indian/Alaska Native Male	-	-	-	-	-	-
American Indian/Alaska Native Female	-	-	-	-	-	-
Unknown	-	-	-	-	-	-
<b>Total</b>	<b>106</b>		<b>66</b>		<b>677</b>	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Akron Consortium 6

**Table 112. Reported cases of AIDS, Ryan White Title II Akron Consortium 6, by year of diagnosis and transmission category**

Transmission category	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	64	73%	15	30%	426	71%
Injection drug use (IDU)	4	5%	2	4%	33	6%
Male-to-male sexual contact and IDU	2	2%	3	6%	39	7%
Heterosexual contact	-	-	1	2%	13	2%
Blood product	-	-	-	-	15	3%
Other/unknown	18	20%	29	58%	72	12%
Subtotal	88	100%	50	100%	598	100%
<b>Female adult or adolescent</b>						
Injection drug use	1	6%	1	6%	11	14%
Heterosexual contact	7	41%	7	44%	39	50%
Blood product	-	-	-	-	4	5%
Other/unknown	9	53%	8	50%	24	31%
Subtotal	17	100%	16	100%	78	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	1	100%	-	-	1	100%
Other	-	-	-	-	-	-
Subtotal	1	100%	-	-	1	100%
<b>Total</b>	<b>106</b>		<b>66</b>		<b>677</b>	

*Note:* Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Youngstown Consortium 7

**Table 113. Reported persons living with HIV/AIDS as of Dec. 31, 2003, Ryan White Title II Youngstown Consortium 7, by current disease status and selected characteristics**

	Current Disease Status						
	HIV/AIDS			HIV (not AIDS)		AIDS	
	Rate <sup>a</sup>	No.	%	No.	%	No.	%
<b>Gender</b>							
Male	117.4	379	70%	212	65%	167	77%
Female	47.7	165	30%	116	35%	49	23%
Unknown	*	-	-	-	-	-	-
<b>Age (as of 12/2003)</b>							
<13	8.0	9	2%	9	3%	-	-
13-19	7.8	5	1%	4	1%	1	<1%
20-24	16.3	6	1%	5	2%	1	<1%
25-29	76.0	29	5%	26	8%	3	1%
30-39	189.7	168	31%	110	34%	58	27%
40-49	195.9	207	38%	110	34%	97	45%
50-64	97.0	108	20%	56	17%	52	24%
65+	10.7	12	2%	8	2%	4	2%
<b>Race/Ethnicity</b>							
White, not Hispanic	46.9	275	51%	147	45%	128	59%
Black, not Hispanic	304.8	204	38%	127	39%	77	36%
Hispanic	303.5	34	6%	23	7%	11	5%
Asian/Pacific Islander	*	-	-	-	-	-	-
American Indian/Alaska Native	*	1	<1%	1	<1%	-	-
Unknown	*	30	6%	30	9%	-	-
<b>Race/Gender</b>							
White, not Hispanic Male	75.1	212	39%	104	32%	108	50%
White, not Hispanic Female	20.7	63	12%	43	13%	20	9%
Black, not Hispanic Male	385.1	126	23%	74	23%	52	24%
Black, not Hispanic Female	228.0	78	14%	53	16%	25	12%
Hispanic Male	321.4	19	3%	12	4%	7	3%
Hispanic Female	283.5	15	3%	11	3%	4	2%
Asian/Pacific Islander Male	*	-	-	-	-	-	-
Asian/Pacific Islander Female	*	-	-	-	-	-	-
American Indian/Alaska Native Male	*	1	<1%	1	<1%	-	-
American Indian/Alaska Native Female	*	-	-	-	-	-	-
Unknown	*	30	6%	30	9%	-	-
<b>Total</b>	<b>81.4</b>	<b>544</b>		<b>328</b>		<b>216</b>	

*Note:* Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (\*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> The rate is the number of persons living with HIV/AIDS per 100,000 population calculated using 2000 census data.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Youngstown Consortium 7

**Table 114. Reported persons living with HIV/AIDS as of Dec. 31, 2003, Ryan White Title II Youngstown Consortium 7, by current disease status and transmission category**

Transmission category	HIV/AIDS		Current Disease Status			
			HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	154	41%	65	31%	89	53%
Injection drug use (IDU)	34	9%	19	9%	15	9%
Male-to-male sexual contact and IDU	23	6%	7	3%	16	10%
Heterosexual contact	12	3%	6	3%	6	4%
Blood product	7	2%	2	1%	5	3%
Other/unknown	144	39%	108	52%	36	22%
Subtotal	374	100%	207	100%	167	100%
<b>Female adult or adolescent</b>						
Injection drug use	20	13%	12	11%	8	17%
Heterosexual contact	42	27%	24	22%	18	38%
Blood product	-	-	-	-	-	-
Other/unknown	96	61%	74	67%	22	46%
Subtotal	158	100%	110	100%	48	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	10	83%	9	82%	1	100%
Other	2	17%	2	18%	-	-
Subtotal	12	100%	11	100%	1	100%
<b>Total</b>	<b>544</b>		<b>328</b>		<b>216</b>	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (\*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Youngstown Consortium 7

**Table 115. Reported cases of HIV infection, Ryan White Title II Youngstown Consortium 7, by year of diagnosis and selected characteristics**

	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
<b>Gender</b>				
Male	74	80%	76	58%
Female	19	20%	56	42%
Unknown	-	-	-	-
<b>Age</b>				
<13	1	1%	1	1%
13-19	3	3%	3	2%
20-24	4	4%	7	5%
25-29	10	11%	21	16%
30-39	30	32%	42	32%
40-49	34	37%	45	34%
50-64	11	12%	13	10%
65+	-	-	-	-
<b>Race/Ethnicity</b>				
White, not Hispanic	51	55%	54	41%
Black, not Hispanic	34	37%	67	51%
Hispanic	4	4%	6	5%
Asian/Pacific Islander	-	-	-	-
American Indian/Alaska Native	1	1%	1	1%
Unknown	3	3%	4	3%
<b>Race/Gender</b>				
White, not Hispanic Male	44	47%	33	25%
White, not Hispanic Female	7	8%	21	16%
Black, not Hispanic Male	23	25%	37	28%
Black, not Hispanic Female	11	12%	30	23%
Hispanic Male	3	3%	2	2%
Hispanic Female	1	1%	4	3%
Asian/Pacific Islander Male	-	-	-	-
Asian/Pacific Islander Female	-	-	-	-
American Indian/Alaska Native Male	1	1%	1	1%
American Indian/Alaska Native Female	-	-	-	-
Unknown	3	3%	4	3%
<b>Total</b>	<b>93</b>		<b>132</b>	

*Note:* HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Youngstown Consortium 7

**Table 116. Reported cases of HIV infection, Ryan White Title II Youngstown Consortium 7, by year of diagnosis and transmission category**

Transmission category	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
<b>Male adult or adolescent</b>				
Male-to-male sexual contact	36	49%	18	24%
Injection drug use (IDU)	10	14%	3	4%
Male-to-male sexual contact and IDU	3	4%	1	1%
Heterosexual contact	3	4%	4	5%
Blood product	-	-	1	1%
Other/unknown	21	29%	49	64%
Subtotal	73	100%	76	100%
<b>Female adult or adolescent</b>				
Injection drug use	2	11%	1	2%
Heterosexual contact	6	32%	4	7%
Blood product	-	-	-	-
Other/unknown	11	58%	50	91%
Subtotal	19	100%	55	100%
<b>Child (&lt;13 yrs at diagnosis)</b>				
Perinatal	1	100%	1	100%
Other	-	-	-	-
Subtotal	1	-	1	-
<b>Total</b>	<b>93</b>		<b>132</b>	

*Note:* HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Youngstown Consortium 7

**Table 117. Reported cases of AIDS, Ryan White Title II Youngstown Consortium 7, by year of diagnosis and selected characteristics**

	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
<b>Gender</b>						
Male	37	79%	41	65%	359	81%
Female	10	21%	22	35%	82	19%
Unknown	-	-	-	-	-	-
<b>Age</b>						
<13	-	-	-	-	2	<1%
13-19	-	-	-	-	7	2%
20-24	-	-	-	-	14	3%
25-29	5	11%	8	13%	60	14%
30-39	16	34%	25	40%	197	45%
40-49	20	43%	25	40%	121	27%
50-64	6	13%	5	8%	35	8%
65+	-	-	-	-	5	1%
<b>Race/Ethnicity</b>						
White, not Hispanic	30	64%	29	46%	277	63%
Black, not Hispanic	17	36%	31	49%	147	33%
Hispanic	-	-	3	5%	17	4%
Asian/Pacific Islander	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-
Unknown	-	-	-	-	-	-
<b>Race/Gender</b>						
White, not Hispanic Male	26	55%	20	32%	243	55%
White, not Hispanic Female	4	9%	9	14%	34	8%
Black, not Hispanic Male	11	23%	19	30%	106	24%
Black, not Hispanic Female	6	13%	12	19%	41	9%
Hispanic Male	-	-	2	3%	10	2%
Hispanic Female	-	-	1	2%	7	2%
Asian/Pacific Islander Male	-	-	-	-	-	-
Asian/Pacific Islander Female	-	-	-	-	-	-
American Indian/Alaska Native Male	-	-	-	-	-	-
American Indian/Alaska Native Female	-	-	-	-	-	-
Unknown	-	-	-	-	-	-
<b>Total</b>	<b>47</b>		<b>63</b>		<b>441</b>	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Youngstown Consortium 7

**Table 118. Reported cases of AIDS, Ryan White Title II Youngstown Consortium 7, by year of diagnosis and transmission category**

Transmission category	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	27	73%	9	22%	206	58%
Injection drug use (IDU)	3	8%	5	12%	32	9%
Male-to-male sexual contact and IDU	2	5%	2	5%	28	8%
Heterosexual contact	1	3%	2	5%	14	4%
Blood product	-	-	1	2%	29	8%
Other/unknown	4	11%	22	54%	48	13%
Subtotal	37	100%	41	100%	357	100%
<b>Female adult or adolescent</b>						
Injection drug use	3	30%	-	-	21	26%
Heterosexual contact	2	20%	5	23%	27	33%
Blood product	-	-	-	-	4	5%
Other/unknown	5	50%	17	77%	29	36%
Subtotal	10	100%	22	100%	81	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	-	-	-	-	2	67%
Other	-	-	-	-	1	33%
Subtotal	-	-	-	-	3	100%
<b>Total</b>	<b>47</b>		<b>63</b>		<b>441</b>	

*Note:* Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Canton Consortium 8

**Table 119. Reported persons living with HIV/AIDS as of Dec. 31, 2003, Ryan White Title II Canton Consortium 8, by current disease status and selected characteristics**

	Current Disease Status						
	HIV/AIDS			HIV (not AIDS)		AIDS	
	Rate <sup>a</sup>	No.	%	No.	%	No.	%
<b>Gender</b>							
Male	81.3	262	79%	146	77%	116	81%
Female	20.8	71	21%	44	23%	27	19%
Unknown	*	-	-	-	-	-	-
<b>Age (as of 12/2003)</b>							
<13	*	3	1%	1	1%	2	1%
13-19	*	-	-	-	-	-	-
20-24	*	4	1%	4	2%	-	-
25-29	46.2	18	5%	12	6%	6	4%
30-39	127.5	117	35%	79	42%	38	27%
40-49	119.8	123	37%	59	31%	64	45%
50-64	61.1	65	20%	33	17%	32	22%
65+	*	3	1%	2	1%	1	1%
<b>Race/Ethnicity</b>							
White, not Hispanic	39.4	245	74%	135	71%	110	77%
Black, not Hispanic	224.2	73	22%	46	24%	27	19%
Hispanic	145.8	8	2%	2	1%	6	4%
Asian/Pacific Islander	*	-	-	-	-	-	-
American Indian/Alaska Native	*	-	-	-	-	-	-
Unknown	*	7	2%	7	4%	-	-
<b>Race/Gender</b>							
White, not Hispanic Male	67.3	203	61%	112	59%	91	64%
White, not Hispanic Female	13.1	42	13%	23	12%	19	13%
Black, not Hispanic Male	306.7	48	14%	28	15%	20	14%
Black, not Hispanic Female	147.8	25	8%	18	9%	7	5%
Hispanic Male	243.6	7	2%	2	1%	5	3%
Hispanic Female	*	1	<1%	-	-	1	1%
Asian/Pacific Islander Male	*	-	-	-	-	-	-
Asian/Pacific Islander Female	*	-	-	-	-	-	-
American Indian/Alaska Native Male	*	-	-	-	-	-	-
American Indian/Alaska Native Female	*	-	-	-	-	-	-
Unknown	*	7	2%	7	4%	-	-
<b>Total</b>	<b>50.1</b>	<b>333</b>		<b>190</b>		<b>143</b>	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (\*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> The rate is the number of persons living with HIV/AIDS per 100,000 population calculated using 2000 census data.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Canton Consortium 8

**Table 120. Reported persons living with HIV/AIDS as of Dec. 31, 2003, Ryan White Title II Canton Consortium 8, by current disease status and transmission category**

Transmission category	HIV/AIDS		Current Disease Status			
			HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	134	51%	56	38%	78	67%
Injection drug use (IDU)	11	4%	5	3%	6	5%
Male-to-male sexual contact and IDU	10	4%	4	3%	6	5%
Heterosexual contact	9	3%	5	3%	4	3%
Blood product	4	2%	1	1%	3	3%
Other/unknown	94	36%	75	51%	19	16%
Subtotal	262	100%	146	100%	116	100%
<b>Female adult or adolescent</b>						
Injection drug use	3	4%	3	7%	-	-
Heterosexual contact	25	37%	11	26%	14	56%
Blood product	2	3%	1	2%	1	4%
Other/unknown	38	56%	28	65%	10	40%
Subtotal	68	100%	43	100%	25	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	3	100%	1	100%	2	100%
Other	-	-	-	-	-	-
Subtotal	3	100%	1	100%	2	100%
<b>Total</b>	<b>333</b>		<b>190</b>		<b>143</b>	

*Note:* Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (\*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Canton Consortium 8

**Table 121. Reported cases of HIV infection, Ryan White Title II Canton Consortium 8, by year of diagnosis and selected characteristics**

	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
<b>Gender</b>				
Male	45	65%	49	80%
Female	24	35%	12	20%
Unknown	-	-	-	-
<b>Age</b>				
<13	2	3%	-	-
13-19	1	1%	1	2%
20-24	7	10%	3	5%
25-29	10	14%	10	16%
30-39	26	38%	27	44%
40-49	17	25%	10	16%
50-64	6	9%	10	16%
65+	-	-	-	-
<b>Race/Ethnicity</b>				
White, not Hispanic	54	78%	39	64%
Black, not Hispanic	12	17%	15	25%
Hispanic	1	1%	3	5%
Asian/Pacific Islander	-	-	-	-
American Indian/Alaska Native	-	-	-	-
Unknown	2	3%	4	7%
<b>Race/Gender</b>				
White, not Hispanic Male	36	52%	34	56%
White, not Hispanic Female	18	26%	5	8%
Black, not Hispanic Male	8	12%	10	16%
Black, not Hispanic Female	4	6%	5	8%
Hispanic Male	1	1%	2	3%
Hispanic Female	-	-	1	2%
Asian/Pacific Islander Male	-	-	-	-
Asian/Pacific Islander Female	-	-	-	-
American Indian/Alaska Native Male	-	-	-	-
American Indian/Alaska Native Female	-	-	-	-
Unknown	2	3%	4	7%
<b>Total</b>	<b>69</b>		<b>61</b>	

*Note:* HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Canton Consortium 8

**Table 122. Reported cases of HIV infection, Ryan White Title II Canton Consortium 8, by year of diagnosis and transmission category**

Transmission category	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
<b>Male adult or adolescent</b>				
Male-to-male sexual contact	28	62%	12	24%
Injection drug use (IDU)	1	2%	1	2%
Male-to-male sexual contact and IDU	1	2%	-	-
Heterosexual contact	1	2%	1	2%
Blood product	-	-	-	-
Other/unknown	14	31%	35	71%
Subtotal	45	100%	49	100%
<b>Female adult or adolescent</b>				
Injection drug use	2	9%	-	-
Heterosexual contact	11	50%	2	17%
Blood product	-	-	-	-
Other/unknown	9	41%	10	83%
Subtotal	22	100%	12	100%
<b>Child (&lt;13 yrs at diagnosis)</b>				
Perinatal	2	100%	-	-
Other	-	-	-	-
Subtotal	2	-	-	-
<b>Total</b>	<b>69</b>		<b>61</b>	

*Note:* HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Canton Consortium 8

**Table 123. Reported cases of AIDS, Ryan White Title II Canton Consortium 8, by year of diagnosis and selected characteristics**

	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
<b>Gender</b>						
Male	31	78%	22	76%	265	85%
Female	9	23%	7	24%	45	15%
Unknown	-	-	-	-	-	-
<b>Age</b>						
<13	1	3%	-	-	6	2%
13-19	-	-	-	-	-	-
20-24	2	5%	-	-	15	5%
25-29	2	5%	3	10%	38	12%
30-39	17	43%	13	45%	137	44%
40-49	11	28%	9	31%	78	25%
50-64	7	18%	3	10%	31	10%
65+	-	-	1	3%	5	2%
<b>Race/Ethnicity</b>						
White, not Hispanic	31	78%	19	66%	250	81%
Black, not Hispanic	8	20%	7	24%	53	17%
Hispanic	1	3%	3	10%	7	2%
Asian/Pacific Islander	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	-	-
Unknown	-	-	-	-	-	-
<b>Race/Gender</b>						
White, not Hispanic Male	24	60%	14	48%	218	70%
White, not Hispanic Female	7	18%	5	17%	32	10%
Black, not Hispanic Male	6	15%	6	21%	41	13%
Black, not Hispanic Female	2	5%	1	3%	12	4%
Hispanic Male	1	3%	2	7%	6	2%
Hispanic Female	-	-	1	3%	1	<1%
Asian/Pacific Islander Male	-	-	-	-	-	-
Asian/Pacific Islander Female	-	-	-	-	-	-
American Indian/Alaska Native Male	-	-	-	-	-	-
American Indian/Alaska Native Female	-	-	-	-	-	-
Unknown	-	-	-	-	-	-
<b>Total</b>	<b>40</b>		<b>29</b>		<b>310</b>	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Canton Consortium 8

**Table 124. Reported cases of AIDS, Ryan White Title II Canton Consortium 8, by year of diagnosis and transmission category**

Transmission category	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	24	77%	9	41%	185	70%
Injection drug use (IDU)	-	-	-	-	12	5%
Male-to-male sexual contact and IDU	-	-	-	-	12	5%
Heterosexual contact	-	-	-	-	6	2%
Blood product	-	-	-	-	12	5%
Other/unknown	7	23%	13	59%	36	14%
Subtotal	31	100%	22	100%	263	100%
<b>Female adult or adolescent</b>						
Injection drug use	1	13%	-	-	8	20%
Heterosexual contact	5	63%	2	29%	17	41%
Blood product	-	-	-	-	2	5%
Other/unknown	2	25%	5	71%	14	34%
Subtotal	8	100%	7	100%	41	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	1	100%	-	-	6	100%
Other	-	-	-	-	-	-
Subtotal	1	100%	-	-	6	100%
<b>Total</b>	<b>40</b>		<b>29</b>		<b>310</b>	

*Note:* Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Lima Consortium 9A

**Table 125. Reported persons living with HIV/AIDS as of Dec. 31, 2003, Ryan White Title II Lima Consortium 9A, by current disease status and selected characteristics**

	Current Disease Status						
	HIV/AIDS			HIV (not AIDS)		AIDS	
	Rate <sup>a</sup>	No.	%	No.	%	No.	%
<b>Gender</b>							
Male	71.0	181	70%	85	61%	96	81%
Female	29.4	77	30%	54	39%	23	19%
Unknown	*	-	-	-	-	-	-
<b>Age (as of 12/2003)</b>							
<13	*	2	1%	-	-	2	2%
13-19	*	2	1%	1	1%	1	1%
20-24	22.2	7	3%	6	4%	1	1%
25-29	45.1	14	5%	11	8%	3	3%
30-39	119.1	86	33%	43	31%	43	36%
40-49	124.6	98	38%	58	42%	40	34%
50-64	56.3	44	17%	18	13%	26	22%
65+	7.1	5	2%	2	1%	3	3%
<b>Race/Ethnicity</b>							
White, not Hispanic	40.2	195	76%	103	74%	92	77%
Black, not Hispanic	260.9	49	19%	29	21%	20	17%
Hispanic	107.9	9	3%	3	2%	6	5%
Asian/Pacific Islander	*	1	<1%	-	-	1	1%
American Indian/Alaska Native	*	-	-	-	-	-	-
Unknown	*	4	2%	4	3%	-	-
<b>Race/Gender</b>							
White, not Hispanic Male	60.8	145	56%	67	48%	78	66%
White, not Hispanic Female	20.2	50	19%	36	26%	14	12%
Black, not Hispanic Male	238.0	24	9%	12	9%	12	10%
Black, not Hispanic Female	287.4	25	10%	17	12%	8	7%
Hispanic Male	163.6	7	3%	2	1%	5	4%
Hispanic Female	*	2	1%	1	1%	1	1%
Asian/Pacific Islander Male	*	1	<1%	-	-	1	1%
Asian/Pacific Islander Female	*	-	-	-	-	-	-
American Indian/Alaska Native Male	*	-	-	-	-	-	-
American Indian/Alaska Native Female	*	-	-	-	-	-	-
Unknown	*	4	2%	4	3%	-	-
<b>Total</b>	<b>49.9</b>	<b>258</b>		<b>139</b>		<b>119</b>	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (\*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> The rate is the number of persons living with HIV/AIDS per 100,000 population calculated using 2000 census data.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Lima Consortium 9A

**Table 126. Reported persons living with HIV/AIDS as of Dec. 31, 2003, Ryan White Title II Lima Consortium 9A, by current disease status and transmission category**

Transmission category	HIV/AIDS		Current Disease Status			
			HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	87	48%	37	44%	50	53%
Injection drug use (IDU)	10	6%	4	5%	6	6%
Male-to-male sexual contact and IDU	9	5%	4	5%	5	5%
Heterosexual contact	10	6%	3	4%	7	7%
Blood product	7	4%	2	2%	5	5%
Other/unknown	57	32%	35	41%	22	23%
Subtotal	180	100%	85	100%	95	100%
<b>Female adult or adolescent</b>						
Injection drug use	12	16%	8	15%	4	19%
Heterosexual contact	29	39%	17	31%	12	57%
Blood product	-	-	-	-	-	-
Other/unknown	34	45%	29	54%	5	24%
Subtotal	75	100%	54	100%	21	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	3	100%	-	-	3	100%
Other	-	-	-	-	-	-
Subtotal	3	100%	-	-	3	100%
<b>Total</b>	<b>258</b>		<b>139</b>		<b>119</b>	

*Note:* Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (\*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Lima Consortium 9A

**Table 127. Reported cases of HIV infection, Ryan White Title II Lima Consortium 9A, by year of diagnosis and selected characteristics**

	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
<b>Gender</b>				
Male	48	68%	48	77%
Female	23	32%	14	23%
Unknown	-	-	-	-
<b>Age</b>				
<13	-	-	-	-
13-19	4	6%	1	2%
20-24	7	10%	4	6%
25-29	9	13%	5	8%
30-39	21	30%	24	39%
40-49	19	27%	18	29%
50-64	10	14%	9	15%
65+	1	1%	1	2%
<b>Race/Ethnicity</b>				
White, not Hispanic	52	73%	47	76%
Black, not Hispanic	18	25%	6	10%
Hispanic	1	1%	5	8%
Asian/Pacific Islander	-	-	-	-
American Indian/Alaska Native	-	-	-	-
Unknown	-	-	4	6%
<b>Race/Gender</b>				
White, not Hispanic Male	39	55%	34	55%
White, not Hispanic Female	13	18%	13	21%
Black, not Hispanic Male	9	13%	6	10%
Black, not Hispanic Female	9	13%	-	-
Hispanic Male	-	-	4	6%
Hispanic Female	1	1%	1	2%
Asian/Pacific Islander Male	-	-	-	-
Asian/Pacific Islander Female	-	-	-	-
American Indian/Alaska Native Male	-	-	-	-
American Indian/Alaska Native Female	-	-	-	-
Unknown	-	-	4	6%
<b>Total</b>	<b>71</b>		<b>62</b>	

*Note:* HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Lima Consortium 9A

**Table 128. Reported cases of HIV infection, Ryan White Title II Lima Consortium 9A, by year of diagnosis and transmission category**

Transmission category	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
<b>Male adult or adolescent</b>				
Male-to-male sexual contact	21	44%	17	35%
Injection drug use (IDU)	1	2%	-	-
Male-to-male sexual contact and IDU	3	6%	1	2%
Heterosexual contact	2	4%	2	4%
Blood product	1	2%	-	-
Other/unknown	20	42%	28	58%
Subtotal	48	100%	48	100%
<b>Female adult or adolescent</b>				
Injection drug use	2	9%	-	-
Heterosexual contact	9	39%	5	36%
Blood product	-	-	-	-
Other/unknown	12	52%	9	64%
Subtotal	23	100%	14	100%
<b>Child (&lt;13 yrs at diagnosis)</b>				
Perinatal	-	-	-	-
Other	-	-	-	-
Subtotal	-	-	-	-
<b>Total</b>	<b>71</b>		<b>62</b>	

*Note:* HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Lima Consortium 9A

**Table 129. Reported cases of AIDS, Ryan White Title II Lima Consortium 9A, by year of diagnosis and selected characteristics**

	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
<b>Gender</b>						
Male	29	74%	25	89%	206	85%
Female	10	26%	3	11%	35	15%
Unknown	-	-	-	-	-	-
<b>Age</b>						
<13	-	-	-	-	5	2%
13-19	-	-	-	-	2	1%
20-24	1	3%	2	7%	8	3%
25-29	8	21%	2	7%	45	19%
30-39	12	31%	14	50%	88	37%
40-49	8	21%	6	21%	55	23%
50-64	10	26%	3	11%	32	13%
65+	-	-	1	4%	6	2%
<b>Race/Ethnicity</b>						
White, not Hispanic	30	77%	18	64%	201	83%
Black, not Hispanic	9	23%	5	18%	31	13%
Hispanic	-	-	4	14%	8	3%
Asian/Pacific Islander	-	-	1	4%	1	<1%
American Indian/Alaska Native	-	-	-	-	-	-
Unknown	-	-	-	-	-	-
<b>Race/Gender</b>						
White, not Hispanic Male	24	62%	16	57%	177	73%
White, not Hispanic Female	6	15%	2	7%	24	10%
Black, not Hispanic Male	5	13%	5	18%	21	9%
Black, not Hispanic Female	4	10%	-	-	10	4%
Hispanic Male	-	-	3	11%	7	3%
Hispanic Female	-	-	1	4%	1	<1%
Asian/Pacific Islander Male	-	-	1	4%	1	<1%
Asian/Pacific Islander Female	-	-	-	-	-	-
American Indian/Alaska Native Male	-	-	-	-	-	-
American Indian/Alaska Native Female	-	-	-	-	-	-
Unknown	-	-	-	-	-	-
<b>Total</b>	<b>39</b>		<b>28</b>		<b>241</b>	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Lima Consortium 9A

**Table 130. Reported cases of AIDS, Ryan White Title II Lima Consortium 9A, by year of diagnosis and transmission category**

Transmission category	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	16	55%	6	24%	135	66%
Injection drug use (IDU)	-	-	-	-	7	3%
Male-to-male sexual contact and IDU	1	3%	1	4%	8	4%
Heterosexual contact	1	3%	2	8%	7	3%
Blood product	1	3%	-	-	15	7%
Other/unknown	10	34%	16	64%	33	16%
Subtotal	29	100%	25	100%	205	100%
<b>Female adult or adolescent</b>						
Injection drug use	1	10%	1	33%	6	19%
Heterosexual contact	6	60%	-	-	19	61%
Blood product	-	-	-	-	-	-
Other/unknown	3	30%	2	67%	6	19%
Subtotal	10	100%	3	100%	31	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	-	-	-	-	4	80%
Other	-	-	-	-	1	20%
Subtotal	-	-	-	-	5	100%
<b>Total</b>	<b>39</b>		<b>28</b>		<b>241</b>	

*Note:* Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Mansfield Consortium 9B

**Table 131. Reported persons living with HIV/AIDS as of Dec. 31, 2003, Ryan White Title II Mansfield Consortium 9B, by current disease status and selected characteristics**

	Current Disease Status						
	HIV/AIDS			HIV (not AIDS)		AIDS	
	Rate <sup>a</sup>	No.	%	No.	%	No.	%
<b>Gender</b>							
Male	73.8	220	81%	109	78%	111	83%
Female	17.5	53	19%	31	22%	22	17%
Unknown	*	-	-	-	-	-	-
<b>Age (as of 12/2003)</b>							
<13	7.4	8	3%	5	4%	3	2%
13-19	*	3	1%	3	2%	-	-
20-24	*	4	1%	3	2%	1	1%
25-29	50.2	18	7%	13	9%	5	4%
30-39	92.8	78	29%	40	29%	38	29%
40-49	117.7	110	40%	46	33%	64	48%
50-64	49.4	48	18%	28	20%	20	15%
65+	*	4	1%	2	1%	2	2%
<b>Race/Ethnicity</b>							
White, not Hispanic	34.8	195	71%	101	72%	94	71%
Black, not Hispanic	237.5	65	24%	29	21%	36	27%
Hispanic	*	3	1%	1	1%	2	2%
Asian/Pacific Islander	*	-	-	-	-	-	-
American Indian/Alaska Native	*	1	<1%	-	-	1	1%
Unknown	*	9	3%	9	6%	-	-
<b>Race/Gender</b>							
White, not Hispanic Male	60.0	165	60%	85	61%	80	60%
White, not Hispanic Female	10.5	30	11%	16	11%	14	11%
Black, not Hispanic Male	275.1	44	16%	16	11%	28	21%
Black, not Hispanic Female	184.6	21	8%	13	9%	8	6%
Hispanic Male	*	2	1%	-	-	2	2%
Hispanic Female	*	1	<1%	1	1%	-	-
Asian/Pacific Islander Male	*	-	-	-	-	-	-
Asian/Pacific Islander Female	*	-	-	-	-	-	-
American Indian/Alaska Native Male	*	1	<1%	-	-	1	1%
American Indian/Alaska Native Female	*	-	-	-	-	-	-
Unknown	*	9	3%	9	6%	-	-
<b>Total</b>	<b>45.4</b>	<b>273</b>		<b>140</b>		<b>133</b>	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (\*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> The rate is the number of persons living with HIV/AIDS per 100,000 population calculated using 2000 census data.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Mansfield Consortium 9B

**Table 132. Reported persons living with HIV/AIDS as of Dec. 31, 2003, Ryan White Title II Mansfield Consortium 9B, by current disease status and transmission category**

Transmission category	HIV/AIDS		Current Disease Status			
			HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	104	49%	40	38%	64	59%
Injection drug use (IDU)	12	6%	6	6%	6	6%
Male-to-male sexual contact and IDU	15	7%	7	7%	8	7%
Heterosexual contact	5	2%	2	2%	3	3%
Blood product	3	1%	2	2%	1	1%
Other/unknown	75	35%	48	46%	27	25%
Subtotal	214	100%	105	100%	109	100%
<b>Female adult or adolescent</b>						
Injection drug use	3	6%	1	3%	2	10%
Heterosexual contact	23	46%	13	45%	10	48%
Blood product	-	-	-	-	-	-
Other/unknown	24	48%	15	52%	9	43%
Subtotal	50	100%	29	100%	21	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	8	89%	5	83%	3	100%
Other	1	11%	1	17%	-	-
Subtotal	9	100%	6	100%	3	100%
<b>Total</b>	<b>273</b>		<b>140</b>		<b>133</b>	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (\*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Mansfield Consortium 9B

**Table 133. Reported cases of HIV infection, Ryan White Title II Mansfield Consortium 9B, by year of diagnosis and selected characteristics**

	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
<b>Gender</b>				
Male	45	80%	45	79%
Female	11	20%	12	21%
Unknown	-	-	-	-
<b>Age</b>				
<13	1	2%	-	-
13-19	3	5%	2	4%
20-24	6	11%	4	7%
25-29	5	9%	7	12%
30-39	25	45%	18	32%
40-49	15	27%	17	30%
50-64	-	-	8	14%
65+	1	2%	1	2%
<b>Race/Ethnicity</b>				
White, not Hispanic	38	68%	36	63%
Black, not Hispanic	16	29%	17	30%
Hispanic	-	-	1	2%
Asian/Pacific Islander	-	-	-	-
American Indian/Alaska Native	-	-	-	-
Unknown	2	4%	3	5%
<b>Race/Gender</b>				
White, not Hispanic Male	31	55%	29	51%
White, not Hispanic Female	7	13%	7	12%
Black, not Hispanic Male	12	21%	13	23%
Black, not Hispanic Female	4	7%	4	7%
Hispanic Male	-	-	-	-
Hispanic Female	-	-	1	2%
Asian/Pacific Islander Male	-	-	-	-
Asian/Pacific Islander Female	-	-	-	-
American Indian/Alaska Native Male	-	-	-	-
American Indian/Alaska Native Female	-	-	-	-
Unknown	2	4%	3	5%
<b>Total</b>	<b>56</b>		<b>57</b>	

*Note:* HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Mansfield Consortium 9B

**Table 134. Reported cases of HIV infection, Ryan White Title II Mansfield Consortium 9B, by year of diagnosis and transmission category**

Transmission category	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
<b>Male adult or adolescent</b>				
Male-to-male sexual contact	17	39%	12	27%
Injection drug use (IDU)	4	9%	3	7%
Male-to-male sexual contact and IDU	2	5%	2	4%
Heterosexual contact	5	11%	-	-
Blood product	-	-	1	2%
Other/unknown	16	36%	27	60%
Subtotal	44	100%	45	100%
<b>Female adult or adolescent</b>				
Injection drug use	2	18%	-	-
Heterosexual contact	3	27%	5	42%
Blood product	-	-	-	-
Other/unknown	6	55%	7	58%
Subtotal	11	100%	12	100%
<b>Child (&lt;13 yrs at diagnosis)</b>				
Perinatal	1	100%	-	-
Other	-	-	-	-
Subtotal	1	-	-	-
<b>Total</b>	<b>56</b>		<b>57</b>	

*Note:* HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Mansfield Consortium 9B

**Table 135. Reported cases of AIDS, Ryan White Title II Mansfield Consortium 9B, by year of diagnosis and selected characteristics**

	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
<b>Gender</b>						
Male	28	85%	33	87%	276	88%
Female	5	15%	5	13%	37	12%
Unknown	-	-	-	-	-	-
<b>Age</b>						
<13	-	-	-	-	7	2%
13-19	-	-	-	-	2	1%
20-24	1	3%	-	-	9	3%
25-29	6	18%	5	13%	57	18%
30-39	13	39%	11	29%	128	41%
40-49	12	36%	15	39%	85	27%
50-64	-	-	7	18%	21	7%
65+	1	3%	-	-	4	1%
<b>Race/Ethnicity</b>						
White, not Hispanic	25	76%	26	68%	267	85%
Black, not Hispanic	8	24%	12	32%	42	13%
Hispanic	-	-	-	-	3	1%
Asian/Pacific Islander	-	-	-	-	-	-
American Indian/Alaska Native	-	-	-	-	1	<1%
Unknown	-	-	-	-	-	-
<b>Race/Gender</b>						
White, not Hispanic Male	22	67%	23	61%	240	77%
White, not Hispanic Female	3	9%	3	8%	27	9%
Black, not Hispanic Male	6	18%	10	26%	32	10%
Black, not Hispanic Female	2	6%	2	5%	10	3%
Hispanic Male	-	-	-	-	3	1%
Hispanic Female	-	-	-	-	-	-
Asian/Pacific Islander Male	-	-	-	-	-	-
Asian/Pacific Islander Female	-	-	-	-	-	-
American Indian/Alaska Native Male	-	-	-	-	1	<1%
American Indian/Alaska Native Female	-	-	-	-	-	-
Unknown	-	-	-	-	-	-
<b>Total</b>	<b>33</b>		<b>38</b>		<b>313</b>	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Mansfield Consortium 9B

**Table 136. Reported cases of AIDS, Ryan White Title II Mansfield Consortium 9B, by year of diagnosis and transmission category**

Transmission category	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	15	54%	11	33%	172	63%
Injection drug use (IDU)	3	11%	2	6%	20	7%
Male-to-male sexual contact and IDU	1	4%	2	6%	23	8%
Heterosexual contact	3	11%	-	-	7	3%
Blood product	-	-	-	-	13	5%
Other/unknown	6	21%	18	55%	37	14%
Subtotal	28	100%	33	100%	272	100%
<b>Female adult or adolescent</b>						
Injection drug use	2	40%	-	-	6	18%
Heterosexual contact	2	40%	2	40%	19	56%
Blood product	-	-	-	-	1	3%
Other/unknown	1	20%	3	60%	8	24%
Subtotal	5	100%	5	100%	34	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	-	-	-	-	5	71%
Other	-	-	-	-	2	29%
Subtotal	-	-	-	-	7	100%
<b>Total</b>	<b>33</b>		<b>38</b>		<b>313</b>	

*Note:* Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Southeast Consortium 9C

**Table 137. Reported persons living with HIV/AIDS as of Dec. 31, 2003, Ryan White Title II Southeast Consortium 9C, by current disease status and selected characteristics**

	Current Disease Status						
	HIV/AIDS			HIV (not AIDS)		AIDS	
	Rate <sup>a</sup>	No.	%	No.	%	No.	%
<b>Gender</b>							
Male	77.5	318	80%	130	69%	188	90%
Female	18.6	79	20%	58	31%	21	10%
Unknown	*	-	-	-	-	-	-
<b>Age (as of 12/2003)</b>							
<13	*	4	1%	1	1%	3	1%
13-19	*	3	1%	2	1%	1	<1%
20-24	18.8	11	3%	9	5%	2	1%
25-29	15.5	8	2%	5	3%	3	1%
30-39	111.2	129	32%	68	36%	61	29%
40-49	130.5	164	41%	75	40%	89	43%
50-64	52.6	71	18%	25	13%	46	22%
65+	6.0	7	2%	3	2%	4	2%
<b>Race/Ethnicity</b>							
White, not Hispanic	42.6	341	86%	159	85%	182	87%
Black, not Hispanic	153.9	36	9%	14	7%	22	11%
Hispanic	*	4	1%	1	1%	3	1%
Asian/Pacific Islander	*	-	-	-	-	-	-
American Indian/Alaska Native	*	2	1%	1	1%	1	<1%
Unknown	*	14	4%	13	7%	1	<1%
<b>Race/Gender</b>							
White, not Hispanic Male	71.5	279	70%	113	60%	166	79%
White, not Hispanic Female	15.1	62	16%	46	24%	16	8%
Black, not Hispanic Male	164.5	24	6%	7	4%	17	8%
Black, not Hispanic Female	136.4	12	3%	7	4%	5	2%
Hispanic Male	*	4	1%	1	1%	3	1%
Hispanic Female	*	-	-	-	-	-	-
Asian/Pacific Islander Male	*	-	-	-	-	-	-
Asian/Pacific Islander Female	*	-	-	-	-	-	-
American Indian/Alaska Native Male	*	2	1%	1	1%	1	<1%
American Indian/Alaska Native Female	*	-	-	-	-	-	-
Unknown	*	14	4%	13	7%	1	<1%
<b>Total</b>	<b>47.6</b>	<b>397</b>		<b>188</b>		<b>209</b>	

Note: Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (\*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

<sup>a</sup> The rate is the number of persons living with HIV/AIDS per 100,000 population calculated using 2000 census data.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Southeast Consortium 9C

**Table 138. Reported persons living with HIV/AIDS as of Dec. 31, 2003, Ryan White Title II Southeast Consortium 9C, by current disease status and transmission category**

Transmission category	HIV/AIDS		Current Disease Status			
			HIV (not AIDS)		AIDS	
	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	153	49%	43	33%	110	60%
Injection drug use (IDU)	20	6%	6	5%	14	8%
Male-to-male sexual contact and IDU	6	2%	2	2%	4	2%
Heterosexual contact	17	5%	9	7%	8	4%
Blood product	6	2%	2	2%	4	2%
Other/unknown	110	35%	68	52%	42	23%
Subtotal	312	100%	130	100%	182	100%
<b>Female adult or adolescent</b>						
Injection drug use	3	4%	2	4%	1	5%
Heterosexual contact	30	39%	19	34%	11	55%
Blood product	2	3%	-	-	2	10%
Other/unknown	41	54%	35	63%	6	30%
Subtotal	76	100%	56	100%	20	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	5	56%	2	100%	3	43%
Other	4	44%	-	-	4	57%
Subtotal	9	100%	2	100%	7	100%
<b>Total</b>	<b>397</b>		<b>188</b>		<b>209</b>	

*Note:* Living with HIV/AIDS represents all persons ever diagnosed and reported with HIV or AIDS who have not been reported as having died as of Dec. 31, 2003. Living with HIV/AIDS by current disease status breaks out the number of persons living with HIV/AIDS by cases that were HIV (not AIDS) and those who have been diagnosed with AIDS as of Dec. 31, 2003.

Asterisk (\*) indicates rates were not calculated for case counts < 5.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Southeast Consortium 9C

**Table 139. Reported cases of HIV infection, Ryan White Title II Southeast Consortium 9C, by year of diagnosis and selected characteristics**

	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
<b>Gender</b>				
Male	81	84%	63	76%
Female	16	16%	20	24%
Unknown	-	-	-	-
<b>Age</b>				
<13	-	-	-	-
13-19	3	3%	2	2%
20-24	3	3%	4	5%
25-29	9	9%	5	6%
30-39	45	46%	28	34%
40-49	25	26%	29	35%
50-64	12	12%	13	16%
65+	-	-	2	2%
<b>Race/Ethnicity</b>				
White, not Hispanic	81	84%	68	82%
Black, not Hispanic	13	13%	9	11%
Hispanic	2	2%	1	1%
Asian/Pacific Islander	-	-	-	-
American Indian/Alaska Native	-	-	-	-
Unknown	1	1%	5	6%
<b>Race/Gender</b>				
White, not Hispanic Male	68	70%	52	63%
White, not Hispanic Female	13	13%	16	19%
Black, not Hispanic Male	10	10%	7	8%
Black, not Hispanic Female	3	3%	2	2%
Hispanic Male	2	2%	1	1%
Hispanic Female	-	-	-	-
Asian/Pacific Islander Male	-	-	-	-
Asian/Pacific Islander Female	-	-	-	-
American Indian/Alaska Native Male	-	-	-	-
American Indian/Alaska Native Female	-	-	-	-
Unknown	1	1%	5	6%
<b>Total</b>	<b>97</b>		<b>83</b>	

*Note:* HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Southeast Consortium 9C

**Table 140. Reported cases of HIV infection, Ryan White Title II Southeast Consortium 9C, by year of diagnosis and transmission category**

Transmission category	Year of HIV Diagnosis			
	1998 - 2000		2001 - 2003	
	No.	%	No.	%
<b>Male adult or adolescent</b>				
Male-to-male sexual contact	39	48%	14	22%
Injection drug use (IDU)	6	7%	3	5%
Male-to-male sexual contact and IDU	1	1%	1	2%
Heterosexual contact	4	5%	2	3%
Blood product	-	-	-	-
Other/unknown	31	38%	43	68%
Subtotal	81	100%	63	100%
<b>Female adult or adolescent</b>				
Injection drug use	1	6%	-	-
Heterosexual contact	5	31%	5	25%
Blood product	-	-	-	-
Other/unknown	10	63%	15	75%
Subtotal	16	100%	20	100%
<b>Child (&lt;13 yrs at diagnosis)</b>				
Perinatal	-	-	-	-
Other	-	-	-	-
Subtotal	-	-	-	-
<b>Total</b>	<b>97</b>		<b>83</b>	

*Note:* HIV diagnoses reflect all cases of HIV infection diagnosed for the first time, including cases that are diagnosed with HIV infection, not AIDS; cases that are diagnosed with HIV and later progressed to AIDS; and cases concurrently diagnosed with HIV and AIDS.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Southeast Consortium 9C

**Table 141. Reported cases of AIDS, Ryan White Title II Southeast Consortium 9C, by year of diagnosis and selected characteristics**

	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
<b>Gender</b>						
Male	51	89%	51	93%	355	90%
Female	6	11%	4	7%	41	10%
Unknown	-	-	-	-	-	-
<b>Age</b>						
<13	-	-	-	-	3	1%
13-19	1	2%	-	-	7	2%
20-24	-	-	-	-	5	1%
25-29	3	5%	-	-	53	13%
30-39	25	44%	22	40%	166	42%
40-49	20	35%	24	44%	115	29%
50-64	8	14%	8	15%	44	11%
65+	-	-	1	2%	3	1%
<b>Race/Ethnicity</b>						
White, not Hispanic	48	84%	42	76%	359	91%
Black, not Hispanic	9	16%	11	20%	31	8%
Hispanic	-	-	1	2%	4	1%
Asian/Pacific Islander	-	-	-	-	1	<1%
American Indian/Alaska Native	-	-	-	-	-	-
Unknown	-	-	1	2%	1	<1%
<b>Race/Gender</b>						
White, not Hispanic Male	45	79%	38	69%	325	82%
White, not Hispanic Female	3	5%	4	7%	34	9%
Black, not Hispanic Male	6	11%	11	20%	24	6%
Black, not Hispanic Female	3	5%	-	-	7	2%
Hispanic Male	-	-	1	2%	4	1%
Hispanic Female	-	-	-	-	-	-
Asian/Pacific Islander Male	-	-	-	-	1	<1%
Asian/Pacific Islander Female	-	-	-	-	-	-
American Indian/Alaska Native Male	-	-	-	-	-	-
American Indian/Alaska Native Female	-	-	-	-	-	-
Unknown	-	-	1	2%	1	<1%
<b>Total</b>	<b>57</b>		<b>55</b>		<b>396</b>	

Note: Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

Source: Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.

## Ryan White Title II Southeast Consortium 9C

**Table 142. Reported cases of AIDS, Ryan White Title II Southeast Consortium 9C, by year of diagnosis and transmission category**

Transmission category	Year of AIDS Diagnosis					
	1998 - 2000		2001 - 2003		Cumulative	
	No.	%	No.	%	No.	%
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	30	59%	22	43%	238	68%
Injection drug use (IDU)	3	6%	4	8%	16	5%
Male-to-male sexual contact and IDU	-	-	-	-	13	4%
Heterosexual contact	3	6%	4	8%	13	4%
Blood product	-	-	-	-	15	4%
Other/unknown	15	29%	21	41%	53	15%
Subtotal	51	100%	51	100%	348	100%
<b>Female adult or adolescent</b>						
Injection drug use	2	33%	-	-	7	18%
Heterosexual contact	2	33%	2	50%	20	50%
Blood product	-	-	-	-	8	20%
Other/unknown	2	33%	2	50%	5	13%
Subtotal	6	100%	4	100%	40	100%
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	-	-	-	-	2	25%
Other	-	-	-	-	6	75%
Subtotal	-	-	-	-	8	100%
<b>Total</b>	<b>57</b>		<b>55</b>		<b>396</b>	

*Note:* Cumulative AIDS diagnoses include persons with a diagnosis of AIDS, reported from the beginning of the epidemic (1981) through 2003.

Dash (-) indicates no cases were reported for the given category.

*Source:* Ohio Department of Health HIV/AIDS Surveillance Program. Data reported through Dec. 31, 2004.